



IRELAND

NATIONAL INVENTORY REPORT 2007

GREENHOUSE GAS EMISSIONS 1990 - 2005 REPORTED TO THE UNITED NATIONS FRAMEWORK CONVENTION ON CLIMATE CHANGE

M. McGettigan, P. Duffy, N. Connolly, B. Hyde and P. O'Brien

Environmental Protection Agency
Johnstown Castle Estate, Wexford, Ireland
Telephone : +353 53 60600 Fax : +353 53 60699

CONTENTS

Acknowledgements	v
List of Tables	vi
List of Figures	vii

EXECUTIVE SUMMARY 1

ES.1 Background	1
ES.2 Emissions Profile and Key Categories.....	1
ES.3 Overview of Emissions Estimates and Trends.....	2
ES.4 Indirect Greenhouse Gases	3

Chapter One INTRODUCTION 4

1.1 Background and Context	4
1.2 Scope of GHG Inventories.....	5
1.2.1 Gases and Global Warming Potential	5
1.2.2 IPCC Reporting Format	5
1.3 Institutional and Procedural Arrangements	6
1.3.1 Current Practice	6
1.3.2 National System Development and Implementation	7
1.4 Overview of Methodologies	11
1.5 Overview of Key Categories.....	11
1.5.1 Key Categories at IPCC Level 2.....	11
1.5.2 Disaggregated Key Categories	13
1.5.3 Application of Results.....	14
1.6 Quality Assurance and Quality Control	14
1.7 Uncertainty Assessment	15
1.8 Completeness and Time-Series Consistency.....	17

Chapter Two EMISSION TRENDS 24

2.1 Trends in Total Emissions	24
2.2 Trends by Sector and Gas.....	24
2.3 Emissions of Indirect Greenhouse Gases	26

Chapter Three ENERGY..... 33

3.1 Overview of <i>Energy</i> Sector	33
3.2 Sectoral Approach for Emissions from Energy Use	35
3.2.1 Combustion Sources	35
3.2.2 Fugitive Emissions	38
3.3 IPCC Reference Approach for CO₂ Emissions from Energy Use.....	40
3.4 Comparison CO₂ Estimates from the Sectoral and Reference Approach Results	40
3.5 Memo Items	41
3.6 QA/QC in <i>Energy</i>	41

3.7	Recalculations in <i>Energy</i>	42
Chapter Four INDUSTRIAL PROCESSES		44
4.1	Overview of <i>Industrial Processes</i> Sector.....	44
4.2	Emissions from 2.A Mineral Products.....	45
4.2.1	2.A.1 Cement Production	46
4.2.2	2.A.2 Lime Production	46
4.2.3	2.A.3 Limestone and Dolomite Use	48
4.3	Emissions of HFC, PFC and SF₆ from <i>Industrial Processes</i>	48
4.3.1	Special Studies	48
4.3.2	HFC, PFC and SF ₆ Time Series 1990-2005.....	49
4.4	Recalculations for <i>Industrial Processes</i>	51
4.5	Improvements in <i>Industrial Processes</i>.....	51
Chapter Five SOLVENT AND OTHER PRODUCT USE.....		54
5.1	Overview of <i>Solvent and Other Product Use</i> Sector	54
5.2	NM VOC and CO₂ Inventory Time-Series	54
5.3	Recalculations for <i>Solvent and Other Product Use</i>	55
Chapter Six AGRICULTURE.....		58
6.1	Overview of <i>Agriculture</i> Sector	58
6.2	CH₄ Emissions from Enteric Fermentation	59
6.2.1	Overall Approach	59
6.2.2	Enteric Fermentation in Dairy Cattle	60
6.2.3	Enteric Fermentation in Beef Cattle	61
6.2.4	Enteric Fermentation in Other Cattle.....	62
6.2.5	Summary of Tier 2 Emission Factors for Cattle.....	62
6.2.6	Enteric Fermentation in Other Livestock	63
6.3	CH₄ Emissions from Manure Management	63
6.4	N₂O Emissions from Manure Management.....	64
6.5	N₂O Emissions from Agricultural Soils	65
6.5.1	4.D.1 Direct Soil Emissions.....	65
6.5.2	4.D.2 Pasture Range and Paddock Manure.....	67
6.5.3	4.D.3 Indirect Emissions	67
6.6	QA/QC in <i>Agriculture</i>	68
6.7	Recalculations in <i>Agriculture</i>.....	68
6.8	Improvements in <i>Agriculture</i>	69
Chapter Seven LAND USE LAND-USE CHANGE AND FORESTRY		71
7.1	Introduction	71
7.2	Overview of <i>LULUCF</i> Sector	71
7.2.1	Sector Coverage	71
7.2.2	Land Use Definitions and Land-Use Change Matrices.....	72
7.2.3	Soil Type and Soil Organic Carbon	73
7.2.4	Estimation of Emissions from Soils	74

7.3	5.A Forest Land	80
	7.3.1 Carbon Stock Change in Living Biomass	80
	7.3.2 Carbon Stock Change in Dead Organic Matter	82
	7.3.3 Net Carbon Stock Change in Soils	82
	7.3.4 Emissions of Non-CO ₂ Gases	83
7.4	5.B Cropland	83
	7.4.1 Cropland Areas	83
	7.4.2 Carbon Stock Change in Biomass	84
	7.4.3 Carbon Stock Change in Soils	84
	7.4.4 N ₂ O Emissions in Cropland.....	85
7.5	5.C Grassland	85
	7.5.1 Grassland Areas	85
	7.5.2 Carbon Stock Changes in Grassland	86
	7.5.3 Agricultural Lime Application	87
7.6	5.D Wetlands	87
	7.6.1 Wetland Areas.....	87
	7.6.2 Carbon Stock Changes in Wetland	88
7.7	5.E Settlements	89
	7.7.1 Areas of Settlements	89
	7.7.2 Carbon Stock Changes in Settlements.....	89
7.8	5.F Other Land	89
	7.8.1 Areas of Other Land	89
	7.8.2 Carbon Stock Changes in Other Land.....	90
7.9	Uncertainties in LULUCF	90
7.10	Recalculations in LULUCF	90
7.11	Improvements in LULUCF	96
	 Chapter Eight WASTE	 97
8.1	Overview of Waste Sector	97
8.2	Solid Waste Disposal	98
	8.2.1 Methodology for CH ₄ Emissions from Solid Waste Disposal.....	98
	8.2.2 CH ₄ Production Potential of Solid Waste.....	98
	8.2.3 CH ₄ Emissions from Solid Waste in 2005	100
8.3	Emissions from Wastewater Handling	102
	8.3.1 CH ₄ Emissions from Wastewater handling	102
	8.3.2 N ₂ O Emissions from Human Sewage	102
8.4	Recalculations in Waste	103
8.5	Improvements in Waste	103
	 Chapter Nine RECALCULATIONS AND IMPROVEMENTS	 106
9.1	Recalculation Issues	106
9.2	Recalculations in 1990-2004 Time-Series	106
9.3	Effect of Recalculations in the 1990-2004 Time Series	107
9.4	Inventory Improvements to Date	108
9.5	Inventory Improvements Going Forward	108
	 References	 113

Glossary	117
-----------------------	------------

<i>Annex A Greenhouse Gases and IPCC Reporting Formats</i>	<i>120</i>
<i>Annex B Energy Balance Sheets 2005</i>	<i>123</i>
<i>Annex C Calculation Sheets for Energy 2005</i>	<i>126</i>
<i>Annex D Activity Data for Agriculture</i>	<i>133</i>
<i>Annex E Activity Data and Carbon Stock Changes for 5.A Forest Land.....</i>	<i>139</i>
<i>Annex F Activity Data for 6.A Solid Waste Disposal on Land</i>	<i>143</i>

Acknowledgements

The authors wish to express their appreciation to the various Government Departments, Sustainable Energy Ireland, the Central Statistics Office, the Forest Service, COFORD, Teagasc, the Electricity Supply Board and the many other data suppliers and individuals who have contributed to the subject matter of this inventory report.

List of Tables

Table	Page
1.1 Summary of Methods	9
1.2 Summary of Emission Factors	10
1.3 Key Categories at IPCC Level 2 in 1990	12
1.4 Key Categories at IPCC Level 2 in 2005	13
1.5 Disaggregated Key Categories 1990	18
1.6 Disaggregated Key Categories 2005	19
1.7 Key Category Trend Assessment	20
1.8 Tier 1 Uncertainty Estimates 2005	21
1.9 Summary of Completeness	23
2.1 Greenhouse Gas Emissions 1990-2005	26
2.2 Emissions of SO ₂ , NO _x , NMVOC and CO 1990-2005	28
3.1 Level 3 Category Coverage for <i>Energy</i>	34
3.2 Percentage Change in Emissions from Energy due to Recalculations	43
4.1 Level 3 Category Coverage for <i>Industrial Processes</i>	45
4.2 Emissions from <i>Industrial Processes</i> 1990-2005	47
4.3 Emissions of HFC, PFC and SF ₆ from <i>Industrial Processes</i> 1990-2005	52
4.4 Recalculated Emission Estimates for <i>Industrial Processes</i> 1990-2004	53
5.1 NMVOC and CO ₂ Emissions from <i>Solvent and Other Product Use</i> 1990-2005	56
5.2 Recalculations for <i>Solvent and Other Product Use</i> 1990-2005	57
6.1 Level 3 Category Coverage for <i>Agriculture</i>	59
6.2 Animal Classification for Cattle Population	60
6.3 Tier 2 CH ₄ Emission Factors for 1990, 2004 and 2005	63
6.4 Percentage Change in Emissions from <i>Agriculture</i> due to Recalculations	70
7.1 Level 3 Category Coverage for <i>Land Use Land-Use Change and Forestry</i>	73
7.2 Emissions and Removals from LULUCF 1990-2005	75
7.3 Land Use Categories	76
7.4 Soil Class Coverage and Soil Organic Carbon	74
7.5 Land Use Matrices	77

7.6	Adjustment Factors for SOC	79
7.7	Area Statistics for Peatlands	88
7.8	Mechanism for Calculation of Land Area Converted to Settlements	93
8.1	Level 3 Source Category and Gas Coverage for <i>Waste</i>	97
8.2	Parameters for Estimation of CH ₄ Emissions from Wastewater and Sludge 2005	103
8.3	Recalculations for <i>Waste</i> 1996-2000 and 2002-2004	106
9.1	Recalculations by Gas 1990-2004	111
9.2	Recalculations by IPCC Sector 1990-2004	112

List of Figures

Figure		Page
1.1	Inventory Institutional and Procedural Arrangements	8
2.1	Trend in Emissions from Energy 1990-2005	29
2.2	Trend in Emissions from <i>Industrial Processes</i> 1990-2005	29
2.3	Trend in Emissions from <i>Solvent and Other Product Use</i> 1990-2005	30
2.4	Trend in Emissions from <i>Agriculture</i> 1990-2005	30
2.5	Trend in Emissions and Removals from <i>LULUCF</i> 1990-2005	31
2.6	Trend in Emissions from <i>Waste</i> 1990-2005	31
2.7	Total Primary Energy Requirement 1990-2005	32
2.8	Principal Drivers of Emissions from <i>Agriculture</i> 1990-2005	32
3.1	Trend in Emissions from 1.A.1 Energy Industries 1990-2005	37
3.2	Trend in Emissions from 1.A.4 Other Sectors 1990-2005	39
7.1	Estimated Area of Agricultural land in Ireland 1991-2005	91
7.2	Time-series of Land Area converted for Housing	95
7.3	Time-series of Land Area converted for Settlements	95
8.1	Typical CH ₄ Production Pattern in Solid Waste Disposal Sites	99
8.2	Methane Emissions from Solid Waste Disposal Sites 1990-2005	101

EXECUTIVE SUMMARY

ES.1 Background

The reporting guidelines on annual inventories adopted by the United Nations Framework Convention on Climate Change (UNFCCC), describe the scope and reporting of greenhouse gas emission inventories by Parties included in Annex I to the Convention. The guidelines set out the methodologies and procedures to be followed for submitting consistent and comparable data on an annual basis in a timely, efficient and transparent manner to meet the needs of the Convention. The UNFCCC guidelines require that Parties prepare a National Inventory Report (NIR) as one of the key components of their annual submissions to the UNFCCC secretariat. The purpose of the NIR is to describe the input data, methodologies, emission factors, quality assurance and quality control procedures and other information underlying the inventory compilation for greenhouse gases and to give details of any recalculations of inventories previously submitted. It is needed to assess the transparency, completeness and overall quality of the inventories as part of the rigorous ongoing technical review of submissions from Annex I Parties.

The present report constitutes Ireland's NIR for 2007 and refers to the inventory time-series for the years 1990-2005. It is an update of the 2006 report and is compiled according to the structure adopted by Decision 18/CP.9. As such, it includes sections describing emission trends, key emission categories, recalculations and ongoing improvements, in addition to the detailed documentation of methods, activity data and emission factors used for each of the source categories as defined by the Intergovernmental Panel on Climate Change (IPCC). A number of Annexes are part of the report, which include calculation sheets and other appropriate reference material to support the description of methods and to provide adequate transparency, as required by the UNFCCC Reporting Guidelines.

The Environmental Protection Agency performs the role of inventory agency in Ireland and has for many years been responsible for compiling the annual inventories and for submitting the results to meet UNFCCC and EU reporting requirements. Further progress was achieved during 2006 and 2007 to consolidate the Agency's role in this regard and its arrangements with data providers to develop a formal national system for implementation of the Kyoto Protocol. In addition to complying with the UNFCCC reporting guidelines, the 2007 NIR is intended to inform Irish Government departments and institutions involved in the national system, as well as other stakeholders in Ireland, of the level of emissions and the state-of-the-art of Irish greenhouse gas inventories as they address the challenges to comply with commitments under the Kyoto Protocol. The in-depth analysis of key sources and the up-to-date trend data provides useful support for the implementation of the Government's strategy to limit the increase in emissions in some key sectors. An informative NIR, together with activities provided for in the national system, allows data suppliers to become fully aware of the importance of their contributions to the inventory process and it serves to identify areas where improvements in input data can be achieved.

ES.2 Emissions Profile and Key Categories

In 2005, total emissions of greenhouse gases (excluding the *LULUCF* sector) in Ireland were 69,945.48 Gigagrams (Gg) CO₂ equivalent. The *Energy* sector accounted for 66.3 percent of total emissions, *Agriculture* contributed 26.4 percent while a further 4.7 percent emanated from *Industrial Processes* and 2.5 percent was due to *Waste*. Emissions of CO₂ accounted for 67.6 percent of the national total in 2005, with CH₄ and N₂O contributing 18.7 percent and 12.7 percent, respectively. The combined emissions of HFC, PFC and SF₆ accounted for 1.0 percent of total emissions in 2005.

Tier 1 level assessment of emission source categories (ranking on the basis of their contribution to total emissions) taken at the level at which they could be targeted on an individual basis identified 26 key categories in 2005 (excluding the *LULUCF* sector). There were 14 key categories of CO₂, accounting for 66.3 percent of total emissions. There were seven key categories of CH₄, four key categories of N₂O and 1 key category of HFC in level assessment, which accounted for 18.2 percent, 10.3 percent and 0.6 percent, respectively, of total emissions. The results of the Tier 1 key category analysis clearly show the impact of CO₂ emissions from energy consumption on total emissions in Ireland. These combustion sources of CO₂ emissions accounted for 14 out of 26 key categories identified by level assessment in 2005 and for 62.9 percent of total emissions. The top ten key categories contributed almost 70 percent of total emissions in 2005 with emissions of CO₂ from the combustion of petrol and diesel by road traffic being the single largest source, accounting for 17.8 percent of the total national emissions.

The application of uncertainty analysis for Irish greenhouse gas inventories using the IPCC approach indicates an overall level uncertainty of 6.2 percent in the 2005 inventory and a trend uncertainty of 3.6 percent for the period 1990 to 2005. These values represent some reductions on those in previous submissions due to improved methodologies for some major sources of CH₄. The overall outcome is determined largely by the high uncertainty in the estimate of N₂O emissions from agricultural soils, which is a major source in Ireland and for which the IPCC methodology remains very simplified. Two-thirds of total Irish emissions, i.e. the proportion contributed by CO₂, are estimated to have an uncertainty of just over one percent. The impact of HFC, PFC and SF₆ on inventory uncertainty in the year 2005 is negligible because they account for only 1 percent of total emissions.

ES.3 Overview of Emissions Estimates and Trends

A consistent time-series of greenhouse gas inventories is available for the years 1990 through 2005. The results are compiled as a complete set of Common Reporting Format files, generated by the CRF Reporter Tool, the electronic reporting protocol adopted for annual data submissions to the UNFCCC. The annual inventories are substantially complete with respect to both the coverage of the six greenhouse gases for which information is required and the coverage of IPCC source categories. Some recalculations have again been undertaken for the purposes of the 2007 submission and the latest inventories for the years 1990-2005 reflect these revisions.

The latest time-series of estimates show that total GHG emissions in Ireland increased from 55,374.43 Gg CO₂ equivalent in 1990 to 70,922.59 Gg CO₂ equivalent in 2001. Following this long period of sustained increase, the emissions decreased to 68,659.31 Gg CO₂ equivalent in 2004, a reduction of approximately three percent on their highest level in 2001. The emissions increased again in 2005 and the total of 69,945.48 Gg CO₂ equivalent for the year is 1.9 percent higher than that for 2004 and is 26.3 percent higher than emissions in 1990.

Greenhouse gas emissions associated with road traffic grew by 7.1 percent in 2005, making this source category the main contributor to the increase in the total emissions for Ireland. There was an increase of 2.8 percent in emissions from electricity generation in 2005 while the emissions from agriculture, which account for a large proportion of Ireland's emissions, decreased by about 2 percent. Emissions of the fluorinated gases increased by almost 10 percent in 2005 but they represent only one percent of the total.

ES.4 Indirect Greenhouse Gases

The inventory reporting process requires the inclusion of a number of gases whose indirect effects are also relevant to the assessment of human-induced impacts on climate. They include sulphur dioxide (SO₂), nitrogen oxides (NO_x), carbon monoxide (CO) and volatile organic compounds (VOC). Emissions of SO₂ contribute to the formation of aerosols, which may offset the effects of greenhouse gases, while CO, NO_x and VOC are precursors of ozone, another naturally occurring greenhouse gas. This NIR does not describe the methods used to estimate emissions of SO₂, NO_x, CO and VOC but their annual emissions over the period 1990-2005 are included. The emissions of most of these gases have decreased substantially in this period under various forms of control legislation emanating from the European Commission and the Convention on Long Range Transboundary Air Pollution (UNECE, 1999). The reductions achieved in Ireland are of the order of 60 percent for SO₂ and 40 percent for CO and NMVOC. However, in the case of NO_x, the emissions reductions have been more difficult to achieve, due mainly to the large increase in road traffic, with the result that emissions in 2005 were only 4 percent below their 1990 level, following a decrease of approximately 12 percent on their highest levels around 2001.

Chapter One

Introduction

1.1 Background and Context

Under Articles 4 and 12 of the United Nations Framework Convention on Climate Change (UNFCCC), hereafter referred to as the Convention, Annex I Parties must develop, publish and make available to the Conference of the Parties (COP), the Convention's implementation body, their national inventories of emissions and removals of all greenhouse gases not controlled by the Montreal Protocol. The UNFCCC Reporting Guidelines on Annual Inventories (SBSTA, 1999 and SBSTA, 2002), hereafter referred to as the UNFCCC reporting guidelines, describe the scope and reporting of the emissions inventories. They specify the methodologies and procedures to be followed for submitting consistent and comparable data on an annual basis in a timely, efficient and transparent manner to meet the needs of the Convention. Under the UNFCCC reporting guidelines, Parties are required to compile a National Inventory Report (NIR) and up-to-date annual inventories in an electronic Common Reporting Format (CRF) as the key components of their annual submissions. The objective of the NIR is to describe the methodologies, input data, background information and the entire process of inventory compilation for greenhouse gases and to give explanations for any improvements and recalculations of the inventories reported in previous submissions. The report is needed by expert review teams to assess the transparency, completeness and overall quality of the inventories as part of the ongoing review process for the submissions from Annex I Parties.

The present report constitutes Ireland's NIR for 2007 and refers to the inventory time-series for the years 1990-2005. It is an update of the 2006 report and is compiled according to the structure adopted by Decision 18/CP.9. As such, it addresses the full range of reporting requirements related to annual inventories set down in the UNFCCC reporting guidelines. This NIR is designed to capture the cyclical nature of the reporting process and to clarify the chronology of changes and revisions that are part of normal inventory development. In this way, the report continues to improve the basis for technical assessment and expert review of Irish greenhouse gas inventories. An attempt has been made to provide all the information, including calculation sheets as appropriate, to facilitate replication of the emissions estimates for the most recent year of the inventory time-series so that transparency may be fully tested.

In addition to complying with the UNFCCC reporting guidelines, the report is intended to inform Government departments, national institutions and other stakeholders of the state of the art of Irish greenhouse gas inventories as they address the challenges to comply with commitments under the Kyoto Protocol. In this context, it provides some additional background on relevant emission sources in Ireland, the standard reporting format and other issues for the benefit of those not entirely familiar with the agreed content of the NIR or the general reporting requirements under the Convention. The report is also aimed at all data suppliers, with a view to making them fully aware of the importance of their contributions to the inventory process and to provide a means of identifying areas where improvements in input data may be possible.

The NIR is updated annually in accordance with the UNFCCC guidelines and is published on the web site of the EPA <http://coe.epa.ie/>. Such updating is necessary to keep the UNFCCC secretariat and other interested parties informed of the status of Irish greenhouse gas inventories and to document ongoing improvements, recalculations and other developments affecting the estimates of emissions. The structure of the report is designed to facilitate year-on-year revision in a manner that allows for systematic and efficient assessment of progress towards the achievement of greenhouse gas emission inventories that meet the guiding principles of transparency, consistency, comparability, completeness and accuracy.

1.2 Scope of Greenhouse Gas Inventories

1.2.1 Gases and Global Warming Potential

The full range of greenhouse gases for which emissions data are required under the Convention is given in Table A.1 of Annex A. It includes carbon dioxide (CO₂), methane (CH₄) and nitrous oxide (N₂O), the most widely known and most ubiquitous of the anthropogenic greenhouse gases, along with 13 hydrofluorocarbons (HFC), seven perfluorocarbons (PFC) and sulphur hexafluoride (SF₆). The global warming potentials (GWP) of the various greenhouse gases vary enormously, as shown on Table A.1 of Annex A. The GWP of a gas is a measure of the cumulative warming over a specified time period, e.g. 100 years, resulting from a unit mass of the gas emitted at the beginning of that time period, expressed relative to an absolute GWP of 1 for the reference gas carbon dioxide (IUCC, 1998). The mass emission of any gas multiplied by its GWP gives the equivalent emission of the gas as carbon dioxide. Therefore, while CO₂, CH₄ and N₂O are important because they are normally emitted in large amounts, HFC, PFC and SF₆ are included in the inventory process mainly because of their comparatively much larger GWP values.

The inventory reporting process allows for the inclusion of a number of additional gases whose indirect effects are also relevant to the assessment of human-induced impacts on climate. They include sulphur dioxide (SO₂), nitrogen oxides (NO_x), carbon monoxide (CO) and volatile organic compounds (VOC). Emissions of SO₂ contribute to the formation of aerosols, which may offset the effects of greenhouse gases, while CO, NO_x and VOC are precursors of ozone, another naturally occurring greenhouse gas. This NIR does not describe the methods used to estimate emissions of SO₂, NO_x, CO and VOC but up-to-date estimates of total emissions are included for information purposes.

1.2.2 IPCC Reporting Format

The reporting of greenhouse gas emissions under the Convention is done with reference to the multi-level reporting format adopted by the Intergovernmental Panel on Climate Change (IPCC). This is a standard table format that forms the basis of the CRF and it assigns all potential sources of emission and removals making up a Party's national total to six Level 1 broad source categories. A further category is provided for the reporting of any additional sources that may be specific to individual Parties. The Level 1 source categories are each divided into as many as seven sub-categories, giving a total of 36 Level 2 source/sink categories, which in turn are further sub-divided to give the 126 standard sub-categories disaggregated at Level 3 in the CRF. Table A.2 of Annex A lists the Level 1 and Level 2 source/sink categories. The Level 3 categories are detailed in the description of category coverage and inventory methods and data in the respective sectoral chapters of this NIR. The computation of emissions is usually undertaken at Level 3 or lower, using further appropriate disaggregation (for example, by using fuel type in the case of combustion sources under *1.A Energy-Fuel Combustion*) while summary results are normally published at Level 2.

The IPCC reporting format also includes a number of *Memo Item* entries. These items refer to sources of emissions whose contributions are not included in a Party's national total but which are to be reported because of their importance in relation to the overall assessment of emissions and for comparisons among Parties. Much reference is made throughout this report to the IPCC reporting format when describing source category coverage, methods, emissions and key categories. The national total of emissions that is commonly used excludes the estimates for the Land Use Land-Use Change and Forestry (LULUCF) category in Table A.2 of Annex A, this total being consistent with that for the categories included in Annex A of the Kyoto Protocol.

1.3 Institutional and Procedural Arrangements

1.3.1 Current Practice

Under Section 52 of the Environmental Protection Agency Act of 1992 (DOE, 1992), the Agency is required to establish and maintain databases of information on the environment and to disseminate such information to interested parties. Section 55 of the Act states that the Agency must provide, of its own volition or upon request, information and advice to Ministers of the Government in the performance of their duties. This includes making available such data and materials as are necessary to comply with Ireland's reporting obligations and commitments within the framework of international agreements. These requirements are the regulatory basis on which the EPA prepares annual inventories of greenhouse gases and other important emissions to air in Ireland. The activities related to the compilation and reporting of greenhouse gas emissions constitute one specific ongoing project in the Agency's work programme. The inventories team is engaged in two other parallel projects dealing with emissions of other compounds.

The Department of the Environment Heritage and Local Government (DEHLG) has designated the EPA as the inventory agency with responsibility for the submission of emissions data to the UNFCCC Secretariat and to the UNECE Secretariat. The Agency's Office of Environmental Assessment (OEA) compiles the national greenhouse gas emission inventories on behalf of DEHLG for submission under the Framework Convention on Climate Change and Decision 280/2004/EC (EP and CEU, 2004), the latter being the basis for EU Member States' reporting under the Convention and the Kyoto Protocol.

Figure 1.1 gives a schematic overview of the institutions and information flows involved in compiling Irish emission inventories for a variety of compounds emitted into the atmosphere, including greenhouse gases. This institutional framework has become consolidated and formalised during 2006 and early 2007 to serve as Ireland's national system under Article 5 of the Kyoto Protocol. The EPA receives the energy balance statistics from Sustainable Energy Ireland (SEI) while agricultural statistics are obtained from the Central Statistics Office (CSO) and the Department of Agriculture and Food (DAF). These primary inputs are complemented by contributions from specific energy and industrial sub-sectors and by information from some EPA databases. The emissions of SO₂, NO_x and CO₂ from power plants are obtained on a plant-by-plant basis from electricity companies and similar data are available for a number of large industrial sources. Gas production and distribution companies supply estimates of the gas losses associated with natural gas. The Annual Environmental Reports (AER) submitted by licensed companies provide useful information on emissions to air and they may be readily accessed within the Agency for inventory purposes. Information in the National Waste Database maintained by the EPA together with data on landfill gas utilisation and flaring is used as the primary inputs to estimate methane emissions from landfills.

The Emissions Trading Unit (ETU), established under the EPA Office of Licence and


Guidance (OLG) in late 2003 to implement Directive 2003/87/EC (EP and CEU, 2003) in Ireland, is an important new source of activity-specific and company-specific data on emissions of greenhouse gases. Emissions trading covers approximately 110 plants and installations in Ireland with combined CO₂ emissions of approximately 24,000 Gg annually, equal to approximately 35 percent of total greenhouse gas emissions. Guidance provided under the associated Decision on methodologies for estimating and reporting greenhouse gas emissions (CEC, 2004) to support the Directive, together with monitoring and verification mechanisms administered by the ETU, consolidates and improves the information in relation to a substantial proportion of emissions for the purposes of reporting under the Convention. The returns under the scheme for 2005 are fully utilised in the national inventory for 2005.

Various preparatory calculations, conversions and reallocations are generally required for both the emissions estimates reported directly to the inventory agency and the activity data acquired from the different sources before they become part of the annual inventory in each reporting cycle. This is undertaken at the lowest possible level of aggregation compatible with the adopted calculation methodologies and the CRF structure. Suitable emission factors are applied to the activity data to calculate emissions in a top-down manner and the results are combined with those already available from bottom-up approaches from some data suppliers for appropriate aggregation according to the IPCC reporting format. All inventory data, including background information and supporting calculation spreadsheets, are stored at the EPA offices in Dublin.

1.3.2 National System Development and Implementation

In 2005, UK consultants carried out a scoping study to identify the essential elements and structure of a national inventory system for Ireland to meet the needs of Decision 280/2004/EC and to comply with obligations under Articles 5 and 7 of the Kyoto Protocol. The report (Thistlethwaite et al, 2005) describes how institutional arrangements among the EPA, DEHLG and other stakeholders may be reorganised, extended and legally consolidated across all participating institutions to strengthen inventory capacity within the Agency and ensure that more formal and comprehensive mechanisms of data collection and processing are established for long term implementation. The report sets out the extent of institutional participation, resource requirements and the form of legal arrangements necessary to perform the functions prescribed in the guidelines for national systems and enable Ireland to meet the objectives specified in those guidelines. The consultants' proposals for system development were benchmarked on systems in operation in other EU Member States and they prescribed how the arrangements in place could be enhanced within the existing statutory framework. The scoping report also made recommendations on internal inventory review and proposed a database system to facilitate more efficient data management and reporting.

The development of Ireland's national inventory system was largely completed during 2006 and early 2007, building on the framework that has been applied for many years. It establishes formal procedures for the planning, preparation and management of the national atmospheric inventory and identifies the roles and responsibilities of all the organisations involved in its compilation. This was achieved through extensive discussions with the key data providers leading to the adoption of Memoranda of Understanding between them and the inventory agency stipulating the scope, timing and quality of the inputs necessary for inventory compilation in accordance with the guidelines for national systems. The EPA remains as the inventory agency and is designated as the single national entity with overall responsibility for the annual greenhouse gas inventory. As a formal management system, the national system aims for continuous improvement to increase the quality and robustness the national atmospheric inventory over time.



DCMNR : Department of Communications Marine & Natural Resources
 ETU : Emissions Trading Unit
 DAF : Department of Agriculture and Food
 CSO : Central Statistics Office
 IPCC : Intergovernmental Panel on Climate Change

ESB : Electricity Supply Board
 EPA : Environmental Protection Agency
 BGE : Bord Gais Eireann
 EIG : Emissions Inventory Guidebook
 SEI : Sustainable Energy Ireland

Figure 1.1. Inventory Institutional and Procedural Arrangements

Table 1.1. Summary of Methods

IPCC SOURCE AND SINK CATEGORIES	CO ₂	CH ₄	N ₂ O	HFC	PFC	SF ₆
1. Energy						
A. Fuel Combustion (Sectoral Approach)						
1. Energy Industries	Tier 1 & 3	Tier 1 & 3	Tier 1 & 3	NA	NA	NA
2. Manufacturing Industries and Construction	Tier 1	Tier 1	Tier 1	NA	NA	NA
3. Transport	Tier 1 & 2	Tier 1 & 3	Tier 1 & 3	NA	NA	NA
4. Other Sectors	Tier 1	Tier 1	Tier 1	NA	NA	NA
5. Other	NA	NA	NA	NA	NA	NA
B. Fugitive Emissions from Fuels						
1. Solid Fuels	NA	NA	NA	NA	NA	NA
2. Oil and Natural Gas	CS	CS	NA	NA	NA	NA
2. Industrial Processes						
A. Mineral Products	Tier 1&2	NA	NA	NA	NA	NA
B. Chemical Industry	NA	NA	NA	NA	NA	NA
C. Metal Production	NA	NA	NA	NA	NA	NA
D. Other Production	NA	NA	NA	NA	NA	NA
E. Production of Halocarbons and SF ₆	NA	NA	NA	NA	NA	NA
F. Consumption of Halocarbons and SF ₆	NA	NA	NA	Tier 1,2& 3	Tier 2	Tier 1 & 2
G. Other	NA	NA	NA	NA	NA	NA
3. Solvent and Other Product Use	CS, C	NA	NA	NA	NA	NA
4. Agriculture						
A. Enteric Fermentation	NA	Tier 1 & 2	NA	NA	NA	NA
B. Manure Management	NA	Tier 1 & 2	Tier 1	NA	NA	NA
C. Rice Cultivation	NA	NA	NA	NA	NA	NA
D. Agricultural Soils	NA	NA	Tier 1a & 1b	NA	NA	NA
E. Prescribed Burning of Savannas	NA	NA	NA	NA	NA	NA
F. Field Burning of Agricultural Residues	NA	NA	NA	NA	NA	NA
G. Other	NA	NA	NA	NA	NA	NA
5. Land-Use Land-Use Change Change and Forestry						
A. Forest Land	Tier 1 & 3	NA	NA	NA	NA	NA
B. Cropland	Tier 1	NA	NA	NA	NA	NA
C. Grassland	Tier 1	NA	NA	NA	NA	NA
D. Wetlands	Tier 1	NA	NA	NA	NA	NA
E. Settlements	Tier 1	NA	NA	NA	NA	NA
F. Other Land	Tier 1	NA	NA	NA	NA	NA
G. Other	NA	NA	NA	NA	NA	NA
6. Waste						
A. Solid Waste Disposal on Land	NA	Tier 2	NA	NA	NA	NA
B. Wastewater Handling	NA	Tier 1	Tier 1	NA	NA	NA
C. Waste Incineration	NA	NA	NA	NA	NA	NA
D. Other	NA	NA	NA	NA	NA	NA
7. Other	NA	NA	NA	NA	NA	NA
International Bunkers						
Aviation	Tier 1	D	D	NA	NA	NA
Marine	D	D	D	NA	NA	NA
Multilateral Operations	NA	NA	NA	NA	NA	NA
CO₂ Emissions from Biomass	Tier 1	Tier 1	Tier 1	NA	NA	NA

Tier 1 : IPCC Tier 1 or equivalent
Tier 2 : IPCC Tier 2 or equivalent
Tier 3 : IPCC Tier 3 or equivalent

CS : Country specific
C : CORINAIR
D : IPCC Default

Table 1.2. Summary of Emission Factors

IPCC SOURCE AND SINK CATEGORIES	CO ₂	CH ₄	N ₂ O	HFC	PFC	SF ₆
1. Energy						
A. Fuel Combustion (Sectoral Approach)						
1. Energy Industries	PS, CS	C & D	C & D	NA	NA	NA
2. Manufacturing Industries and Construction	C	C & D	C & D	NA	NA	NA
3. Transport	CS	M, C	M, C	NA	NA	NA
4. Other Sectors	CS	C, D	C, D	NA	NA	NA
5. Other	NA	NA	NA	NA	NA	NA
B. Fugitive Emissions from Fuels						
1. Solid Fuels	NA	NA	NA	NA	NA	NA
2. Oil and Natural Gas	CS	CS	NA	NA	NA	NA
2. Industrial Processes						
A. Mineral Products	CS, PS	NA	NA	NA	NA	NA
B. Chemical Industry	NA	NA	NA	NA	NA	NA
C. Metal Production	NA	NA	NA	NA	NA	NA
D. Other Production	NA	NA	NA	NA	NA	NA
E. Production of Halocarbons and SF ₆	NA	NA	NA	NA	NA	NA
F. Consumption of Halocarbons and SF ₆	NA	NA	NA	CS	CS	CS
G. Other	NA	NA	NA	NA	NA	NA
3. Solvent and Other Product Use	C	NA	NA	NA	NA	NA
4. Agriculture						
A. Enteric Fermentation	NA	CS, D	NA	NA	NA	NA
B. Manure Management	NA	CS, D	D	NA	NA	NA
C. Rice Cultivation	NA	NA	NA	NA	NA	NA
D. Agricultural Soils	NA	NA	CS, D	NA	NA	NA
E. Prescribed Burning of Savannas	NA	NA	NA	NA	NA	NA
F. Field Burning of Agricultural Residues	NA	NA	NA	NA	NA	NA
G. Other	NA	NA	NA	NA	NA	NA
5. Land-Use Land-Use Change Change and Forestry						
A. Forest Land	CS, D	NA	NA	NA	NA	NA
B. Cropland	D	NA	NA	NA	NA	NA
C. Grassland	D	NA	NA	NA	NA	NA
D. Wetlands	D	NA	NA	NA	NA	NA
E. Settlements	D	NA	NA	NA	NA	NA
F. Other Land	D	NA	NA	NA	NA	NA
G. Other	NA	NA	NA	NA	NA	NA
6. Waste						
A. Solid Waste Disposal on Land	NA	CS	NA	NA	NA	NA
B. Wastewater Handling	NA	D	D	NA	NA	NA
C. Waste Incineration	NA	NA	NA	NA	NA	NA
D. Other	NA	NA	NA	NA	NA	NA
7. Other	NA	NA	NA	NA	NA	NA
International Bunkers						
Aviation	CS	C	C	NA	NA	NA
Marine	CS	C	C	NA	NA	NA
Multilateral Operations	NA	NA	NA	NA	NA	NA
CO₂ Emissions from Biomass	C	C	C	NA	NA	NA

PS : Plant specific
CS : Country specific
C : CORINAIR

D : Default
M : Model

1.4 Overview of Methodologies

An emissions inventory database normally contains information on measured emission quantities, activity statistics (populations, fuel consumption, vehicle/kilometres of travel, industrial production, forest area), emission factors and the associated emission estimates for a specified list of source categories. In practice, very few measured data are available for greenhouse gases and, consequently, the emissions from most activities are estimated by applying emission factors for each source/gas combination to appropriate activity data for the activity concerned. Virtually all emissions may be ultimately derived on the basis of such simple product of activity data and emission factor. However, a certain amount of data analysis and preparatory calculations are generally needed in order to make available suitable combinations of activity data and emission factors at the level of disaggregation that gives the best estimate of emissions. In the case of some source/gas combinations, it may be necessary to apply sophisticated models to generate the activity data, the emission factors or the emissions. The methods recommended by the IPCC Guidelines and good practice use a tier system to take account of these issues and other factors, such as data availability, technical expertise, inventory capacity and other circumstances, which may vary considerably across countries.

Table 1.1 and Table 1.2 present an overview of the methodologies and emission factors used by Ireland to estimate emissions now reported for the years 1990-2005. The current situation regarding data availability and national circumstances dictates the use of a combination of Tier 1 and Tier 2 methods across the IPCC source categories. These methods range from relatively simple calculations for CO₂ emissions from combustion sources and some industrial processes, where quite basic inputs are required, to much more in-depth analysis in other source categories. Examples of the latter include the estimation of N₂O from agricultural soils and CH₄ from landfills, for which several interdependent steps must be followed and many contributing factors must be taken into account. On a sector/gas basis, there is approximately equal application of country-specific and default emission factors. Source categories for which the estimates are based on country-specific methods and emission factors account for 75 percent of total emissions.

1.5 Overview of Key Categories

The IPCC good practice guidance defines a key category as one that is prioritised within the national inventory system because its emission estimate has a significant influence on the Party's total inventory in terms of the absolute level of emissions, the trend in emissions or both. Information about key categories is considered to be crucial to the choice of methodology for individual sources and to the management and reduction of overall inventory uncertainty. The identification of such categories is recommended in order that inventory agencies can give them priority in the preparation of annual inventories, especially in cases where resources may be limited. Information on key categories is clearly also vital for the development of policies and measures for emissions reduction. The IPCC good practice guidance provides several methods for undertaking the analysis of key categories that can be applied at any appropriate level of source aggregation, depending on the information available. The simplest approach is again used for 2005 to further highlight which sources of emissions are the most important in Ireland.

1.5.1 Key Categories at IPCC Level 2

As inventories of CO₂, CH₄ and N₂O were being developed in Ireland during the 1990s, it was quickly established that CO₂ emissions from fuel combustion made by far the largest contribution to the national total for these three primary greenhouse gases. It was also

evident that CH₄ emissions produced by large cattle herds and the N₂O emissions from agricultural soils, associated with intensive farming practices and large inputs of nitrogen, were also major sources, even if the estimates were more uncertain than those of CO₂. A good first estimate of key categories is therefore provided by considering the emissions aggregated at the IPCC Level 2 source category classification, which clearly indicates the importance of CO₂ emissions from fuel combustion and CH₄ and N₂O emissions from agriculture.

The results at the IPCC Level 2 source category classification may be readily drawn from the CRF Summary 2 and those for 1990 and 2005 are shown in Table 1.3 and Table 1.4, respectively. It can be seen that there are six highly significant key categories of emissions in Ireland. They are the CO₂ combustion sources in *1.A.1 Energy Industries*, *1.A.4 Other Sectors*, *1.A.2 Manufacturing Industries and Construction* and *1.A.3 Transport*, along with the CH₄ emissions from category *4.A Enteric Fermentation* and N₂O emissions from *4.D Agricultural Soils*. These categories accounted for 84 percent and 86 percent of total emissions in 1990 and 2005, respectively. In the case of 2005 emissions, only three additional Level 2 source categories are needed to reach the cumulative 95 percent threshold that defines a key category. The increase in the contribution of CO₂ emissions from category *1.A.3 Transport* from 9.11 percent in 1990 to 18.5 percent in 2005 is notable, along with the corresponding reductions in the contributions from the two categories in *Agriculture*. This simple analysis of key categories continues to prove useful to the formulation of abatement strategies and for prioritising work on inventories in Ireland.

Table 1.3. Key Categories at IPCC Level 2 in 1990

IPCC Level 2 Source Category		GHG	Emissions in 1990 Gg CO ₂ eq	1990 Level Assessment %	Cumulative Total of Level %
1.A.1	Energy Industries	CO ₂	11,158.62	20.15	20.15
1.A.4	Other Sectors	CO ₂	10,064.90	18.18	38.33
4.A	Enteric Fermentation	CH ₄	9,337.80	16.86	55.19
4.D	Agricultural Soils	N ₂ O	7,005.34	12.65	67.84
1.A.3	Transport	CO ₂	5,045.02	9.11	76.95
1.A.2	Manufacturing Industries and Construction	CO ₂	3,969.76	7.17	84.12
4.B	Manure Management	CH ₄	2,313.90	4.18	88.30
6.A	Solid Waste Disposal on Land	CH ₄	1,332.05	2.41	90.70
2.B.2	Nitric Acid Production	N ₂ O	1,035.40	1.87	92.57
2.B.1	Ammonia Production	CO ₂	989.17	1.79	94.36
2.A.1	Cement Production	CO ₂	884.00	1.60	95.96

Table 1.4. Key Categories at IPCC Level 2 in 2005

IPCC Level 2 Source Category		GHG	Emissions in 2005 Gg CO ₂ eq	2005 Level Assessment %	Cumulative Total of Level %
1.A.1	Energy Industries	CO ₂	15,657.29	22.38	22.38
1.A.3	Transport	CO ₂	12,942.05	18.50	40.89
1.A.4	Other Sectors	CO ₂	10,549.80	15.08	55.97
4.A	Enteric Fermentation	CH ₄	9,049.10	12.94	68.91
4.D	Agricultural Soils	N ₂ O	6,770.68	9.68	78.59
1.A.2	Manufacturing Industries and Construction	CO ₂	5,453.71	7.80	86.39
2.A.1	Cement Production	CO ₂	2,357.06	3.37	89.76
4.B	Manure Management	N ₂ O	2,224.33	3.18	92.94
6.A	Solid Waste Disposal on Land	N ₂ O	1,618.18	2.31	95.25

1.5.2 Disaggregated Key Categories

Ireland uses the Tier 1 methods provided in the IPCC good practice guidance to extend the analysis above to identify key categories that may be treated separately at a more disaggregated level, which gives more information about the individual sources or combination of sources and gases that are of most importance within a Level 2 category. The results of the analysis for Tier 1 level assessment in relation to emissions in both 1990 and 2005 are presented in Table 1.5 and Table 1.6 respectively. Ranking in this way identifies those categories that should be prioritised in the inventory process itself and also the individual components of emissions that could be targeted by specific abatement measures. There is insufficient information available on uncertainties to allow for analysis using the Tier 2 methods. Results for Tier 1 trend assessment for 2005 are shown in Table 1.7. The results of the assessment for 2005 excluding LULUCF categories may be summarised as follows

- (i) level assessment identifies 26 key categories;
- (ii) there are 14 key categories of CO₂ in level assessment, accounting for 66.3 percent of total emissions;
- (iii) there are seven key categories of CH₄ and four key categories of N₂O in level assessment, which account for 18.2 percent and 10.3 percent, respectively, of total emissions;
- (iv) trend assessment identifies the same key categories as level assessment;
- (v) there are 14 key categories of CO₂ in trend assessment, accounting for 72.8 percent of the total trend;
- (vi) there are seven key categories of CH₄ and four key categories of N₂O in trend assessment, which account for 12.5 percent and 8.4 percent, respectively, of the total trend;
- (vii) *Energy* accounts for 14 key categories, *Agriculture* for nine while *Industrial Processes* contributes two and *Waste* contributes one;

The list of key categories given by level assessment in 1990 is very similar to that for 2005 but the higher ranking of the main CO₂ sources in *Energy*, at the expense of CH₄ and N₂O sources in *Agriculture*, is notable in 2005. The top ten key sources contributed

approximately two-thirds of total emissions in both years. The emissions of CO₂ from the combustion of petrol and diesel by road traffic was the single largest source category of greenhouse gas emissions in Ireland in 2005, accounting for 17.8 percent of the total.

1.5.3 Application of Results

The Tier 1 approach to the determination of key source categories is based on the principle that the cumulative uncertainty in their emissions represents 90 percent of the total inventory uncertainty and that 95 percent of total emissions account for this cumulative fraction of uncertainty. This quantitative approach may therefore result in a much larger number of key categories than might be expected using simpler qualitative criteria. In effect, an inventory with only a small number of major emission sources will require the inclusion of many source categories in order to reach the 95 percent emissions threshold.

This is well shown by the results of key source determination for Ireland, based on Tier 1 level assessment. The results excluding LULUCF indicate that half of key categories in 2005 each accounted for less than 3 percent of the total emissions and that only five key categories contributed more than 5 percent each to the total. The Tier 1 analysis adequately identifies the specific sources of emissions that are significant in terms of the overall uncertainty of the inventory but it provides little direction on where to focus priority when the number is large. In these circumstances, information on the uncertainty in the individual source categories and other factors must be taken into account in making decisions regarding the most cost-effective use of inventory capacity related to key source categories.

The results of the Tier 1 key category analysis in Table 1.6 clearly show that the impact of CO₂ emissions from energy consumption on total emissions in Ireland continues to increase. These emissions account for 14 of the key categories listed in Table 1.6 and for two-thirds of total emissions in 2005. While key categories determined by CO₂ emissions from energy consumption have a major bearing on total emissions in Ireland, the remaining potential for significant reduction in the uncertainties associated with these sources is rather limited. The activity data and CO₂ emission factors for *Energy* source categories in general are among the most reliable items of input data in the inventory and there is consequently little scope for improving the accuracy of the emission estimates. The application of a robust Tier 2 methodology for emissions of CH₄ from enteric fermentation in cattle in this submission and the use of verified estimates for CO₂ emissions from cement production means that the contributions from three additional key categories (ranked 3, 10 and 11 in Table 1.6) making up a further 15.1 percent of the total are also known with probably the highest certainty now achievable. The N₂O emissions from *4.D Agricultural Soils*, the CH₄ and N₂O emissions from *4.B Manure Management* and the CH₄ emissions from *6.A Solid Waste Disposal on Land* account for most of the remaining key categories in Table 1.6. The uncertainties in the estimates for these complex sources (Section 1.7) will remain high due to the large number of factors that influence their emissions and the relatively simple methods that must still be used.

1.6 Quality Assurance and Quality Control

In early 2005, the inventory agency in Ireland commissioned a project with UK consultants to establish formal QA/QC procedures in emission inventories that would meet the needs of the UNFCCC reporting requirements. The project developed a QA/QC system including a documented QA/QC plan and procedures along with a QA/QC manual. The manual provides a general overview to the QA/QC system and guidance on the application of the plan and procedures. The QA/QC plan identifies the specific data quality objectives related to the principles of transparency, consistency, completeness, comparability and accuracy required for Ireland's national inventory and provides specific guidance and documentation forms and

templates for the practical implementation of QA/QC procedures. The QA/QC procedures cover such elements as data selection and acquisition, data processing and reporting so that the international requirements under the Kyoto Protocol and Decision 280/2004/EC are met. The manual provides guidance and templates for appropriate quality checking, documentation and traceability, the selection of source data and calculation methodologies and peer review and expert review of inventory data and outlines the annual requirements for continuous improvement for the inventory.

The inventory agency used the 2006 reporting cycle to begin implementation of the basic elements of the new approach to QA/QC and its application has been substantially completed in delivering the 2007 submission. This involved the allocation of responsibilities linked to the national system mentioned in section 1.3.2 and the use of a template spreadsheet system to record the establishment and maintenance of general inventory checking and management activities covering the overall compilation process, as well as the undertaking of specific annual activities and any necessary periodic activities in response to specific events or outcomes in inventory reporting and review. The system facilitates record keeping related to the chain of activities from data capture, through emissions calculations and checking, to archiving and the identification of improvements.

Ireland's calculation spreadsheets in all sectors have been restructured and reorganised to facilitate the QA/QC process and to facilitate more efficient analysis and to ensure ease of transfer of the outputs to the CRF Reporter Tool. This facilitates rapid year-on-year extension of the time-series and efficient updating and recalculation, where appropriate, in the annual reporting cycle. Internal aggregation to various levels corresponding to the CRF tables provides immediate and complete checks on the results.

External reviews of the agriculture sector and of the entire ETS results for 2005 were conducted as important new components of quality assurance at the beginning of 2007. The review for Agriculture was performed by a technical inspector in the Department of Agriculture and Food using the new calculation files with a view to assessing the consistency of the time series which had been subject to considerable improvement and recalculation in the 2006 reporting cycle to account for higher tier methods and advice from the Department on various aspects of input data and calculation parameters. The ETS returns to the Agency's Office of Licence and Guidance (OLG) provided for the complete coverage of CO₂ estimates for categories 1.A.1, 2.A.1, 2.A.2 and 2.A.3 in 2005. When the allocation to these categories from the ETS raw data was completed, the output was returned to the ETS administrator in OLG for final checking against the source data. The participation of Irish inventory experts in the internal review of sector 1.A.1 for the EU and its Member States during 2006 was another useful contribution to external review procedures affecting the present submission.

1.7 Uncertainty Assessment

The Tier 1 method provided by the IPCC good practice guidance has been used to make an assessment of uncertainty in the emissions inventory for 2005 in the same way as for previous years. This method estimates uncertainties for the entire inventory in a particular year and the uncertainty in the trend over time by combining the uncertainties in activity data and emission factors for each source category. The analysis for 2005 is presented in Table 1.8, using emissions on a GWP basis and a level of aggregation that limits the likely dependency and correlation between source categories.

The input values of uncertainty for activity data have been assigned largely on the basis of general information and opinions elicited from the principal data suppliers, such as statistical offices, energy agencies, Government departments and individuals. In the case of country-

specific emission factors for combustion sources, which relate largely to CO₂, expert judgement has been used to assign the uncertainties for the source categories given in Table 1.8 with reasonable confidence, given the well-established properties of the fuels concerned. Uncertainties in the emission factors for other gases released from combustion sources and for other source categories in general are based on information provided in the IPCC Good Practice Guidance and the CORINAIR/EMEP Emission Inventory Guidebook.

The 2003 in-country review report for Ireland concluded that the input values of uncertainty chosen for activity data or emission factors for some sources in the 2001 inventory may not have been entirely appropriate. The uncertainty analysis for subsequent years therefore incorporates changes that have been made following further investigation to determine the most conceptually meaningful values that can be used at the level of source disaggregation being used for the analysis. Sustainable Energy Ireland, the body responsible for compiling the national energy balance, completed a process to expand and improve Irish energy balances in 2006, which provides further insight into uncertainty in the statistical data compiled from annual fuel-use questionnaires. The inventory experts continue to collaborate with SEI in this process to ascertain the views of the energy-use compilers on uncertainty associated with energy quantities disaggregated by sector and by fuel type. New sources of data, such as the ETS returns, are also being investigated in an attempt to substantiate the quantitative estimates of uncertainty in activity data obtained in this way.

In some of the most important emissions sources in *Agriculture* (such as enteric fermentation and agricultural soils) and *Waste* (solid waste disposal, for example) the activity data or emission factors ultimately used are determined by several specific component inputs, which are all individually subject to varying degrees of uncertainty. The uncertainty estimates used for both activity data and emission factor for these sources have been derived by assigning uncertainties to the key component parameters and combining them at the level of activity data or emission factors, as appropriate, for each activity for input to the Tier 1 uncertainty assessment. The footnotes to Table 1.8 show how some of these revised uncertainty inputs were obtained. The application of the Tier 2 method for CH₄ emissions from enteric fermentation in cattle and the use of improved country-specific information related to manure management (Chapter Seven) justifies the adoption of reduced input uncertainties for some of the emission sources in *Agriculture*. Some reduction is also appropriate in the case of CO₂ emissions from cement production, where verified data are now available through implementation of the EU emissions trading scheme (Chapter Three).

The Tier 1 uncertainty analysis for 2005 gives an overall uncertainty of 6.2 percent in total emissions and a trend uncertainty of 3.6 percent for the period 1990 to 2005. The application of improved Tier 2 methods for emissions from enteric fermentation and manure management for cattle has reduced the level of uncertainty to some extent. The overall outcome continues to be determined largely by the uncertainty in the estimate of N₂O emissions from agricultural soils, where an emission factor uncertainty of 100 percent is assumed in order to complete the analysis. Two-thirds of total Irish emissions, i.e. the proportion contributed by CO₂, are estimated to have a level uncertainty of just over one percent. When CH₄ is included, bringing the proportion of total emissions up to 86 percent, the total uncertainty is of the order of two percent, even though there are large uncertainties assigned to the CH₄ emission factors in some source categories. However, it is the influence of N₂O that leads to a substantial uncertainty in total emissions. This influence is not as large in the case of the trend, due to the modest change in emissions of N₂O from 1990 to 2004 and the relatively small share of this gas in total emissions. The impact of HFC, PFC and SF₆ on inventory uncertainty remains negligible because these gases account for only 1 percent of total emissions in Ireland.

1.8 Completeness and Time-Series Consistency

Table 1.9 gives an overview of the level of completeness of the 2005 GHG inventories with respect to the six greenhouse gases covered by the UNFCCC guidelines and the IPCC Level 2 source-category split in operation since 2005. Further detail on source/gas coverage at IPCC Level 3 is provided in the individual chapters describing the inventory methods and data for each Level 1 source-category. The work done for the current reporting cycle serves to maintain a complete and consistent emissions time-series by improving the inventories for the years 1991-2003 to bring them fully into line for those of 1990 and 2004, which were the main focus of the work in 2006. The opportunity has also been taken in this current cycle to improve the first estimates of emissions and removals for all years for LULUCF reported in accordance with the requirements of Decision 13/CP.9. It may be concluded that the principles of completeness and consistency are therefore being observed in so far as can be expected.

Table 1.5 Disaggregated Key Categories 1990

1990 Rank	IPCC Sub-Category	Emission Source/Activity	Gas	1990 Emission exc LULUCF Gg CO ₂ eq	1990 Emission from LULUCF Gg CO ₂ eq	1990 Emission inc LULUCF Gg CO ₂ eq	1990 Level Assessment exc LULUCF %	Cumulative Level exc LULUCF %	1990 Level Assessment inc LULUCF %	Cumulative Level inc LULUCF %
1	1.A.1.	Energy Industries - Solid Fuels	CO ₂	8,009.44		8,009.44	14.46	14.46	13.83	13.83
2	1.A.4.b	Residential- Solid Fuels	CO ₂	5,606.94		5,606.94	10.13	24.59	9.68	23.51
3	4.A.1.	Enteric Fermentation - Non-Dairy Cattle	CH ₄	5,374.69		5,374.69	9.71	34.30	9.28	32.79
4	1.A.3.b.	Road Transportation - Liquid Fuels	CO ₂	4,700.42		4,700.42	8.49	42.78	8.12	40.91
5	4.A.1.	Enteric Fermentation - Dairy Cattle	CH ₄	2,894.77		2,894.77	5.23	48.01	5.00	45.91
6	4.D.1.	Agricultural Soils - Direct Soil Emissions	N ₂ O	2,861.11		2,861.11	5.17	53.18	4.94	50.85
7	4.D.2.	Agricultural Soils - Pasture, Range and Paddock	N ₂ O	2,798.93		2,798.93	5.05	58.23	4.83	55.68
8	1.A.2.	Manufacturing Industries & Construction - Liquid Fuels	CO ₂	2,225.39		2,225.39	4.02	62.25	3.84	59.53
9	1.A.4.a.	Commercial/Institutional - Liquid Fuels	CO ₂	1,976.61		1,976.61	3.57	65.82	3.41	62.94
10	1.A.1.	Energy Industries - Gaseous Fuels	CO ₂	1,880.66		1,880.66	3.40	69.22	3.25	66.19
11	4.D.3.	Agricultural Soils - Indirect Emissions	N ₂ O	1,345.30		1,345.30	2.43	71.65	2.32	68.51
12	6.A.	Waste - Solid Waste Disposal on land	CH ₄	1,332.05		1,332.05	2.41	74.05	2.30	70.81
13	1.A.1.	Energy Industries - Liquid Fuels	CO ₂	1,268.51		1,268.51	2.29	76.34	2.19	73.00
14	4.B.1.	Manure Management - Non-Dairy cattle	CH ₄	1,234.07		1,234.07	2.23	78.57	2.13	75.13
15	1.A.4.b	Residential - Liquid Fuels	CO ₂	1,189.67		1,189.67	2.15	80.72	2.05	77.19
16	5.A.1	LULUCF - Forest land Remaining Forest Land	CO ₂		-1078.90	1,078.90	0.00	80.72	1.86	79.05
17	2.B.	Chemical Industry	N ₂ O	1,035.40		1,035.40	1.87	82.59	1.79	80.84
18	4.A.3.	Enteric Fermentation - Sheep	CH ₄	1,032.48		1,032.48	1.86	84.45	1.78	82.62
19	2.B.	Chemical Industry	CO ₂	989.17		989.17	1.79	86.24	1.71	84.33
20	2.A.1.	Cement Production	CO ₂	884.00		884.00	1.60	87.84	1.53	85.85
21	1.A.2.	Manufacturing Industries & Construction - Gaseous Fuels	CO ₂	873.14		873.14	1.58	89.41	1.51	87.36
22	1.A.2.	Manufacturing Industries & Construction - Solid Fuels	CO ₂	871.24		871.24	1.57	90.99	1.50	88.87
23	1.A.4.c.	Agriculture/Forestry/Fisheries - Liquid Fuels	CO ₂	660.30		660.30	1.19	92.18	1.14	90.01
24	5.C.1	LULUCF - Grassland Remaining Grassland	CO ₂		620.07	620.07	0.00	92.18	1.07	91.08
25	4.B.1.	Manure Management - Dairy Cattle	CH ₄	615.90		615.90	1.11	93.29	1.06	92.14
26	5.A.2.	LULUCF - Land Converted to Forest Land	CO ₂		600.89	600.89	0.00	93.29	1.04	93.18
27	4.B.13.	Manure Management - Solid Storage	N ₂ O	350.78		350.78	0.63	93.93	0.61	93.79
28	4.B.8.	Manure Management - Pigs	CH ₄	327.77		327.77	0.59	94.52	0.57	94.35
29	1.A.1.	Energy Industries - Solid Fuels	N ₂ O	319.01		319.01	0.58	95.09	0.55	94.90
30	1.A.4.b	Residential - Gaseous Fuels	CO ₂	269.73		269.73	0.49	95.58	0.47	95.37

Table 1.6 Disaggregated Key Categories 2005

2005 Rank	IPCC Sub-Category	Emission Source/Activity	Gas	2005 Emission exc LULUCF Gg CO ₂ eq	2005 Emission inc LULUCF Gg CO ₂ eq	2005 Emission inc LULUCF Gg CO ₂ eq	2005 Level Assessment exc LULUCF %	Cumulative Level exc LULUCF %	2005 Level Assessment inc LULUCF %	Cumulative Level inc LULUCF %
1	1.A.3.b.	Road Transport - Liquid Fuels	CO ₂	12,453.90		12,453.90	17.81	17.81	17.28	17.28
2	1.A.1.	Energy Industries - Solid Fuels	CO ₂	8,019.30		8,019.30	11.47	29.27	11.13	28.40
3	4.A.1.	Enteric Fermentation - Non-Dairy Cattle	CH ₄	5,679.05		5,679.05	8.12	37.39	7.88	36.28
4	1.A.1.	Energy Industries - Gaseous Fuels	CO ₂	4,664.00		4,664.00	6.67	44.06	6.47	42.75
5	1.A.2.	Manufacturing Industries and Construction - Liquid Fuels	CO ₂	3,591.34		3,591.34	5.13	49.19	4.98	47.73
6	1.A.4.b	Residential - Liquid Fuels	CO ₂	3,459.38		3,459.38	4.95	54.14	4.80	52.53
7	1.A.1.	Energy Industries - Liquid Fuels	CO ₂	2,973.98		2,973.98	4.25	58.39	4.13	56.66
8	4.D.2.	Agricultural Soils - Pasture, Range & Paddock	N ₂ O	2,772.79		2,772.79	3.96	62.35	3.85	60.50
9	4.D.1.	Agricultural Soils - Direct Soil Emissions	N ₂ O	2,687.67		2,687.67	3.84	66.20	3.73	64.23
10	4.A.1.	Enteric Fermentation - Dairy Cattle	CH ₄	2,515.43		2,515.43	3.60	69.79	3.49	67.72
11	2.A.1.	Cement Production	CO ₂	2,357.06		2,357.06	3.37	73.16	3.27	70.99
12	1.A.4.b	Residential - Solid Fuels	CO ₂	2,159.26		2,159.26	3.09	76.25	3.00	73.99
13	1.A.4.a.	Commercial/Institutional - Liquid Fuels	CO ₂	1,798.00		1,798.00	2.57	78.82	2.49	76.48
14	6.A.	Solid Waste Disposal on land	CH ₄	1,618.18		1,618.18	2.31	81.13	2.24	78.73
15	4.D.3.	Agricultural Soils - Indirect Emissions	N ₂ O	1,310.22		1,310.22	1.87	83.01	1.82	80.54
16	1.A.4.b	Residential - Gaseous Fuels	CO ₂	1,240.51		1,240.51	1.77	84.78	1.72	82.27
17	1.A.2.	Manufacturing Industries and Construction - Gaseous Fuels	CO ₂	1,190.90		1,190.90	1.70	86.48	1.65	83.92
18	4.B.1.	Manure Management - Non-Dairy Cattle	CH ₄	1,164.18		1,164.18	1.66	88.15	1.62	85.53
19	1.A.4.a.	Commercial/Institutional - Gaseous Fuels	CO ₂	958.64		958.64	1.37	89.52	1.33	86.86
20	5.A.1	LULUCF - Forest land Remaining Forest Land	CO ₂		-845.19	845.19	0.00	89.52	1.17	88.04
21	1.A.4.c.	Agriculture/Forestry/Fisheries - Liquid Fuels	CO ₂	828.54		828.54	1.18	90.70	1.15	89.18
22	4.A.3.	Enteric Fermentation - Sheep	CH ₄	809.45		809.45	1.16	91.86	1.12	90.31
23	1.A.2.	Manufacturing Industries and Construction - Solid Fuels	CO ₂	671.47		671.47	0.96	92.82	0.93	91.24
24	4.B.1.	Manure Management - Dairy Cattle	CH ₄	479.33		479.33	0.69	93.51	0.66	91.90
25	5.C.1	LULUCF - Grassland remaining Grassland	CO ₂		477.79	477.79	0.00	93.51	0.66	92.57
26	1.A.3.b.	Road Transport - Liquid Fuels	N ₂ O	443.30		443.30	0.63	94.14	0.61	93.18
27	4.B.8.	Manure Management - Pigs	CH ₄	438.99		438.99	0.63	94.77	0.61	93.79
28	2.F.	Consumption of F Gas and SF ₆	HFC	431.03		431.03	0.62	95.38	0.60	94.39
29	5.C.2.	LULUCF - Land converted to Grassland	CO ₂		-421.22	421.22	0.00	95.38	0.58	94.97
30	4.B.13.	Manure Management - Solid Storage	N ₂ O	353.58		353.58	0.51	95.89	0.49	95.46

Table 1.7 Key Category Trend Assessment 2005 (excluding LULUCF)

Rank	Category	Emission Source	Gas	Emissions in 1990 Gg CO ₂ eq	Emissions in 2005 Gg CO ₂ eq	Level Assessment %	Trend Assessment	Contribution to Trend %	Cumulative Contribution to Trend %
1	1.A.3.b.	Road Transport - Liquid Fuels	CO ₂	4,700.42	12,453.90	17.81	6.8118	18.88	18.88
2	1.A.4.b	Residential - Solid Fuels	CO ₂	5,606.94	2,159.26	3.09	5.6700	15.71	34.59
3	1.A.1.	Energy Industries - Solid Fuels	CO ₂	8,009.44	8,019.30	11.47	2.7375	7.59	42.18
4	1.A.1.	Energy Industries - Gaseous Fuels	CO ₂	1,880.66	4,664.00	6.67	2.3790	6.59	48.77
5	1.A.4.b	Residential - Liquid Fuels	CO ₂	1,189.67	3,459.38	4.95	2.0580	5.70	54.48
6	4.A.1.	Enteric Fermentation - Non-Dairy Cattle	CH ₄	5,374.69	5,679.05	8.12	1.5135	4.19	58.67
7	1.A.1.	Energy Industries - Liquid Fuels	CO ₂	1,268.51	2,973.98	4.25	1.4178	3.93	62.60
8	4.A.1.	Enteric Fermentation - Dairy Cattle	CH ₄	2,894.77	2,515.43	3.60	1.4055	3.90	66.50
9	2.A.1.	Cement Production	CO ₂	884.00	2,357.06	3.37	1.2972	3.60	70.09
10	4.D.1.	Agricultural Soils - Direct Soil Emissions	N ₂ O	2,861.11	2,687.67	3.84	1.1702	3.24	73.34
11	4.D.2.	Agricultural Soils - Pasture, Range & Paddock	N ₂ O	2,798.93	2,772.79	3.96	0.9888	2.74	76.08
12	1.A.4.b	Residential - Gaseous Fuels	CO ₂	269.73	1,240.51	1.77	0.9623	2.67	78.74
13	1.A.4.a.	Commercial/Institutional - Liquid Fuels	CO ₂	1,976.61	1,798.00	2.57	0.8723	2.42	81.16
14	1.A.4.a.	Commercial/Institutional - Gaseous Fuels	CO ₂	223.37	958.64	1.37	0.7223	2.00	83.16
15	1.A.2.	Manufacturing Industries and Construction - Liquid Fuels	CO ₂	2,225.39	3,591.34	5.13	0.7206	2.00	85.16
16	4.A.3.	Enteric Fermentation - Sheep	CH ₄	1,032.48	809.45	1.16	0.5966	1.65	86.81
17	1.A.2.	Manufacturing Industries and Construction - Solid Fuels	CO ₂	871.24	671.47	0.96	0.5160	1.43	88.24
18	4.D.3.	Agricultural Soils - Indirect Emissions	N ₂ O	1,345.30	1,310.22	1.87	0.4997	1.39	89.63
19	4.B.1.	Manure Management - Non-Dairy Cattle	CH ₄	1,234.07	1,164.18	1.66	0.4994	1.38	91.01
20	2.F.	Consumption of F Gas and SF ₆	HFC	0.69	431.03	0.62	0.4673	1.30	92.31
21	1.A.3.b.	Road Transport - Liquid Fuels	N ₂ O	71.30	443.30	0.63	0.3797	1.05	93.36
22	4.B.1.	Manure Management - Dairy Cattle	CH ₄	615.90	479.33	0.69	0.3597	1.00	94.36
23	6.A.	Solid Waste Disposal on land	CH ₄	1,332.05	1,618.18	2.31	0.1462	0.41	94.76
24	1.A.2.	Manufacturing Industries and Construction - Gaseous Fuels	CO ₂	873.14	1,190.90	1.70	0.0457	0.13	94.89
25	1.A.4.c.	Agriculture/Forestry/Fisheries - Liquid Fuels	CO ₂	660.30	828.54	1.18	0.0438	0.12	95.01
26	4.B.8.	Manure Management - Pigs	CH ₄	327.77	438.99	0.63	0.0084	0.02	95.03

Table 1.8 Tier 1 Uncertainty Estimates 2005 (continued on following page)

IPCC Source Category		Gas	Emissions in 1990	Emissions in 2005	Activity Data (AD) Uncertainty	Emission Factor (EF) Uncertainty	Combined Uncertainty	Combined Uncertainty as % of Emissions in 2005	Combined Emissions Uncertainty Squared	Type A Sensitivity	Type B Sensitivity	Uncertainty in Trend in Total Emissions due to AD	Uncertainty in Trend in Total Emissions due to EF	Combined Uncertainty in Trend in Total Emissions	Combined Trend Uncertainty Squared
			Gg CO ₂	Gg CO ₂	%	%	%	%		%	%	%	%	%	%
1A1	Energy-Liquid	CO ₂	1,268.51	2,973.98	1	2.5	2.69	0.11	0.01	0.02	0.05	0.08	0.06	0.10	0.01
1A1	Energy-Solid	CO ₂	8,009.44	8,019.30	1	5	5.10	0.58	0.34	-0.04	0.14	0.20	-0.19	0.28	0.08
1A1	Energy-Gas	CO ₂	1,880.66	4,664.00	1	2.5	2.69	0.18	0.03	0.04	0.08	0.12	0.10	0.16	0.02
1A2	Industry-Liquid	CO ₂	2,010.97	2,566.45	10	2.5	10.31	0.38	0.14	0.00	0.05	0.66	0.00	0.66	0.43
1A2	Industry-Coal	CO ₂	871.24	671.47	2	5	5.39	0.05	0.00	-0.01	0.01	0.03	-0.04	0.05	0.00
1A2	Industry-Pet Coke	CO ₂	214.42	1,024.88	5	10	11.18	0.16	0.03	0.01	0.02	0.13	0.14	0.19	0.04
1A2	Industry-Gas	CO ₂	873.14	1,190.90	2.5	2.5	3.54	0.06	0.00	0.00	0.02	0.08	0.00	0.08	0.01
1A3	Transport-Oil	CO ₂	4,982.98	12,776.92	1	2.5	2.69	0.49	0.24	0.12	0.23	0.33	0.29	0.44	0.19
1A3	Transport-Gas	CO ₂	62.04	165.13	1	2.5	2.69	0.01	0.00	0.00	0.00	0.00	0.00	0.01	0.00
1A4	Comm-Liquid	CO ₂	1,976.61	1,798.00	10	5	11.18	0.29	0.08	-0.01	0.03	0.46	-0.06	0.46	0.21
1A4	Comm-Coal	CO ₂	2.55	103.59	5	10	11.18	0.02	0.00	0.00	0.00	0.01	0.02	0.02	0.00
1A4	Comm-Peat	CO ₂	135.73	1.89	10	20	22.36	0.00	0.00	0.00	0.00	0.00	-0.06	0.06	0.00
1A4	Comm-Gas	CO ₂	223.37	958.64	2.5	2.5	3.54	0.05	0.00	0.01	0.02	0.06	0.03	0.07	0.00
1A4	Res-Liquid	CO ₂	1,101.79	3,311.61	10	5	11.18	0.53	0.28	0.03	0.06	0.85	0.17	0.86	0.75
1A4	Res-Coal	CO ₂	2,483.57	989.59	5	10	11.18	0.16	0.03	-0.04	0.02	0.13	-0.39	0.41	0.17
1A4	Res-Petcoke	CO ₂	87.88	147.76	5	10	11.18	0.02	0.00	0.00	0.00	0.02	0.01	0.02	0.00
1A4	Res-Peat	CO ₂	3,123.37	1,169.67	10	20	22.36	0.37	0.14	-0.05	0.02	0.30	-1.00	1.05	1.09
1A4	Res-Gas	CO ₂	269.73	1,240.51	2.5	2.5	3.54	0.06	0.00	0.02	0.02	0.08	0.04	0.09	0.01
1A4	Agric Liquid	CO ₂	660.30	828.54	10	5	11.18	0.13	0.02	0.00	0.01	0.21	0.00	0.21	0.04
1.B	Fugitive Emissions	CO ₂	138.88	60.20	2.5	10	10.31	0.01	0.00	0.00	0.00	0.00	-0.02	0.02	0.00
2.A.1	Cement Production	CO ₂	884.00	2,357.06	7.5	5	9.01	0.30	0.09	0.02	0.04	0.45	0.11	0.47	0.22
2.A.2	Lime Production	CO ₂	214.08	183.56	5	5	7.07	0.02	0.00	0.00	0.00	0.02	-0.01	0.02	0.00
2.A.3	Limestone and Dolomite Use	CO ₂	7.59	13.14	5	5	7.07	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2.B.1	Ammonia Production	CO ₂	989.17	0.00	1	5	5.10	0.00	0.00	-0.02	0.00	0.00	-0.11	0.11	0.01
3	Solvent and Other Product Use	CO ₂	80.86	75.44	30	5	30.41	0.03	0.00	0.00	0.00	0.06	0.00	0.06	0.00
<i>Total CO₂</i>			<i>32,552.88</i>	<i>47,292.25</i>				<i>1.20</i>	<i>1.45</i>					<i>1.81</i>	<i>3.29</i>
1A	Fuel Comb-All Fuels	CH ₄	156.98	128.63	2	50	50.04	0.09	0.01	0.00	0.00	0.01	-0.06	0.06	0.00
1B	Fugitive Emissions	CH ₄	131.08	56.86	2.5	10	10.31	0.01	0.00	0.00	0.00	0.00	-0.02	0.02	0.00
4A	Ent Ferm. Dairy Cattle	CH ₄	2,894.77	2,515.43	1	15	15.03	0.54	0.29	-0.02	0.05	0.06	-0.31	0.32	0.10
4A	Ent Ferm.Other Cattle	CH ₄	5,374.69	5,679.05	1	15	15.03	1.22	1.49	-0.02	0.10	0.15	-0.30	0.33	0.11
4A	Ent Ferm.Other Livestock	CH ₄	1,068.34	854.62	1	30	30.02	0.37	0.13	-0.01	0.02	0.02	-0.27	0.27	0.07
4B	Man. Manag.Dairy Cattle	CH ₄	615.90	479.33	1	15	15.03	0.10	0.01	-0.01	0.01	0.01	-0.08	0.08	0.01
4B	Man. Manag.Other Cattle	CH ₄	1,234.07	1,164.18	1	15	15.03	0.25	0.06	-0.01	0.02	0.03	-0.11	0.11	0.01
4B	Man. Manag.Other Livestock	CH ₄	463.93	580.82	1	30	30.02	0.25	0.06	0.00	0.01	0.01	0.00	0.02	0.00
6A	Solid Waste	CH ₄	1,332.05	1,618.18	41	47	62.37	1.44	2.08	0.00	0.03	1.69	-0.05	1.70	2.87
6.B	Wastewater Handling	CH ₄	14.70	25.22	10	30	31.62	0.01	0.00	0.00	0.00	0.01	0.00	0.01	0.00
<i>Total CH₄</i>			<i>13,286.51</i>	<i>13,102.32</i>				<i>2.04</i>	<i>4.14</i>					<i>1.78</i>	<i>3.18</i>

Cumulative CO ₂ and CH ₄			45,839.39	60,394.57			2.37		5.59			2.54		6.47	
1A3	Fuel Comb-Road Traffic	N ₂ O	71.30	443.30	1	25	25.02	0.16	0.03	0.01	0.01	0.01	0.16	0.16	0.03
1A	Fuel Comb-Other Sectors	N ₂ O	866.83	1,092.44	2	50	50.04	0.78	0.61	0.00	0.02	0.06	0.00	0.06	0.00
2B	Nitric Acid	N ₂ O	1,035.40	0.00	1	10	10.05	0.00	0.00	-0.02	0.00	0.00	-0.24	0.24	0.06
4B	Liquid System	N ₂ O	55.21	55.76	11.2	100	100.63	0.08	0.01	0.00	0.00	0.02	-0.03	0.03	0.00
4B	Solid Storage and dry lot	N ₂ O	350.78	353.58	11.2	100	100.63	0.51	0.26	0.00	0.01	0.10	-0.16	0.19	0.04
4D	Direct Soil Emissions	N ₂ O	2,861.11	2,687.67	11.2	100	100.63	3.87	14.95	-0.02	0.05	0.77	-1.67	1.84	3.39
4D	Pasture Range and Paddock	N ₂ O	2,798.93	2,772.79	11.2	100	100.63	3.99	15.91	-0.01	0.05	0.79	-1.38	1.59	2.52
4D	Indirect Emissions	N ₂ O	1,345.30	1,310.22	11.2	50	51.24	0.96	0.92	-0.01	0.02	0.37	-0.35	0.51	0.26
6B	Wastewater Handling	N ₂ O	114.00	134.20	10	10	14.14	0.03	0.00	0.00	0.00	0.03	0.00	0.03	0.00
Total N ₂ O			9,498.85	8,849.96			5.72		32.69			2.51		6.30	
Cumulative CO ₂ , CH ₄ , N ₂ O			55,338.24	69,244.54			6.19		38.28			3.57		12.77	
2F	Halocarbons & SF ₆	HFC	0.69	431.03	20	10	22.36	0.14	0.02	0.01	0.01	0.17	0.08	0.19	0.04
2F	Halocarbons & SF ₆	PFC	0.09	173.95	10	2.5	10.31	0.03	0.00	0.00	0.00	0.04	0.01	0.04	0.00
2F	Halocarbons & SF ₆	SF ₆	35.40	95.96	15	5	15.81	0.02	0.00	0.00	0.00	0.03	0.00	0.03	0.00
Total HFC, PFC and SF ₆			36.19	700.94			0.14		0.02			0.20		0.04	
Total all gases			55,374.43	69,945.48					38.30						12.81
			Level Uncertainty in Emissions				6.19				Trend Uncertainty		3.58		

Type A Sensitivity *the change in the difference in overall emissions between the base year and the current year, expressed as a percentage, resulting from a 1 percent increase in emissions of a given source category/gas combination in both the base year and the current year*

Type B Sensitivity *the change in the difference in overall emissions between the base year and the current year, expressed as a percentage, resulting from a 1 percent increase in emissions of a given source category/gas combination in the current year only*

a Based on Equation 6.4 of IPCC GPG with uncertainties of 25%, 25% and 20% for MSW quantity, MSW composition and DOC, respectively

b Based on Equation 6.4 of IPCC GPG with uncertainties of 30%, 30% and 20% for fraction DOC dissimilated, MCF and time of CH₄ release, respectively

c Based on Equation 6.4 of IPCC GPG with uncertainties of 20% and 30% for nitrogen excretion and AWMS proportion, respectively

Table 1.9 Summary of Completeness

IPCC SOURCE AND SINK CATEGORIES	CO₂	CH₄	N₂O	HFC	PFC	SF₆
1. Energy						
A. Fuel Combustion (Sectoral Approach)	All	All	All	NA	NA	NA
1. Energy Industries	All	All	All	NA	NA	NA
2. Manufacturing Industries and Construction	All	All	All	NA	NA	NA
3. Transport	All	All	All	NA	NA	NA
4. Other Sectors	All	All	All	NA	NA	NA
5. Other	NO	NO	NO	NA	NA	NA
B. Fugitive Emissions from Fuels						
1. Solid Fuels	NO	NO	NO	NA	NA	NA
2. Oil and Natural Gas	All	All	Part	NA	NA	NA
2. Industrial Processes						
A. Mineral Products	All	Part	Part	NA	NA	NA
B. Chemical Industry	NO	NO	NO	NO	NO	NO
C. Metal Production	NO	NO	NO	NO	NO	NO
D. Other Production	NE	NA	NA	NA	NA	NA
E. Production of Halocarbons and SF ₆	NA	NA	NA	NO	NO	NO
F. Consumption of Halocarbons and SF ₆	NA	NA	NA	All	All	All
G. Other	NO	NO	NO	NO	NO	NO
3. Solvent and Other Product Use	All	NA	NE	NA	NA	NA
4. Agriculture						
A. Enteric Fermentation	NA	All	NA	NA	NA	NA
B. Manure Management	NA	All	All	NA	NA	NA
C. Rice Cultivation	NA	NO	NA	NA	NA	NA
D. Agricultural Soils	NA	NE	All	NA	NA	NA
E. Prescribed Burning of Savannas	NO	NO	NO	NA	NA	NA
F. Field Burning of Agricultural Residues	NO	NO	NO	NA	NA	NA
G. Other	NO	NO	NO	NA	NA	NA
5. Land-Use Change and Forestry						
A. Forest Land	All	NE	NE	NA	NA	NA
B. Cropland	All	NO	IE	NA	NA	NA
C. Grassland	All	NO	IE	NA	NA	NA
D. Wetlands	All	NE	NE	NA	NA	NA
E. Settlements	Part	NO	NO	NA	NA	NA
F. Other Land	All	NE	NE	NA	NA	NA
G. Other	NO	NO	NO	NA	NA	NA
6. Waste						
A. Solid Waste Disposal on Land	NO	All	NA	NA	NA	NA
B. Wastewater Handling	NA	All	All	NA	NA	NA
C. Waste Incineration	NE	NE	NE	NA	NA	NA
D. Other	NO	NO	NO	NA	NA	NA
7. Other	NO	NO	NO	NA	NA	NA
Memo Items:						
International Bunkers						
Aviation	All	All	All	NA	NA	NA
Marine	All	All	All	NA	NA	NA
Multilateral Operations	NO	NO	NO	NA	NA	NA
CO₂ Emissions from Biomass	All	NA	NA	NA	NA	NA

All : Emissions of the gas are covered for all sources under the source category/memo item

NA : Emissions of the gas not applicable to the source category/memo item

NO : Emissions of the gas does not occur in Ireland for the source category/memo item

NE : Emissions on the gas not estimated for the source category/memo item

Part : Emissions of the gas estimated for some activities in the source category

Chapter Two

Emission Trends

2.1 Trends in Total Emissions

Table 2.1 and Figure 2.1 show the trends in emissions of the six greenhouse gases in Ireland over the period 1990-2005, extracted from the trend tables of the 2005 CRF. The estimates reported here show some changes on those reported in 2006, which reflect recalculations that are fully described in subsequent chapters. The trends in the principal emission components within the six IPCC sectors as CO₂ equivalents are shown on Figure 2.2 through Figure 2.7. Total emissions of the six greenhouse gases in Ireland (excluding net CO₂ from *Land Use Change and Forestry*) increased steadily from 55,374.43 Gg CO₂ equivalent in 1990 to 70,922.59 Gg CO₂ equivalent in 2001 and then decreased slightly to 68,659.31 Gg CO₂ equivalent in 2004. The emissions increased again in 2005 to 69,945.48 Gg CO₂ equivalent, which is 26.3 percent higher than in 1990 and 1.9 percent above the level in 2004.

In 2005, the *Energy* sector accounted for 66.3 percent of total emissions, *Agriculture* contributed 26.4 percent while a further 4.7 percent emanated from *Industrial Processes* and 2.5 percent was due to *Waste*. Emissions of CO₂ accounted for 67.6 percent of the total of 69,945.48 Gg CO₂ equivalent in 2005, with CH₄ and N₂O contributing 18.7 percent and 12.7 percent, respectively. The combined emissions of HFC, PFC and SF₆ accounted for approximately 1 percent of total emissions in 2005. The *Energy* and *Industrial Processes* sectors account for the bulk of the CO₂ emissions, CH₄ emissions are produced mainly in the *Agriculture* and *Waste* sectors and most of the N₂O emissions are generated in *Agriculture*. The trends in the principal drivers of emissions in the key sectors of *Energy* and *Agriculture* are shown in Figure 2.8 and Figure 2.9, respectively.

The large increase in emissions during the period 1990-2001 was clearly driven by the growth in CO₂ emissions from energy use. The increase in CO₂ amounted to 43 percent over these 12 years. The bulk of this increase occurred in the years between 1995 and 2000, during which Ireland experienced a period of unprecedented economic growth and emissions grew by around 3 percent annually. The rate of economic growth slowed down from 2000 to 2004, which together with the closure of ammonia and nitric acid production plants and continued decline in cattle populations and fertilizer use, resulted in some reduction in the emission levels in 2002 and 2003. The increase in 2005 was due largely to increased emissions from road transport and from electricity generation where two new peat-fired stations entered service.

2.2 Trends by Sector and Gas

Fuel combustion in *1.A.3 Transport* and in *1.A.1 Energy Industries* accounted for 28,599.34 Gg CO₂ in 2005 or approximately 41 percent of total greenhouse gas emissions (Figure 2.1). The largest increases in CO₂ emissions have taken place in the transport sector. The CO₂ emissions from transport sources, which are largely accounted for by road traffic in Ireland,

increased by 156 percent between 1990 and 2005, due to sustained growth in the numbers and in the use of passenger cars and goods vehicles. This trend is exaggerated somewhat in latter years by so-called fuel-tourism, whereby a significant proportion of the automotive fuels sold in Ireland is used by vehicles in the UK and other countries. The proportion was estimated to be approximately 12 percent for petrol in 2001-2005 and averaged 25 percent in the case of diesel in the same period. It is worth noting that in 1990 there was significant cross-border movement of automotive fuels into Ireland.

There continues to be heavy reliance on carbon intensive fuels for electricity generation in Ireland and, as electricity demand increased steadily during the 1990s, the associated CO₂ emissions from 1.A.1 Energy Industries (electricity generation and oil refining) increased by 55 percent from 11,158.61 Gg in 1990 to 17,266.56 Gg in 2001. Some gains were achieved from energy efficiency and fuel switching as some new electricity producers entered the market in 2002 and 2003, with the result that CO₂ emissions from energy industries reduced to 15,283.51 Gg in 2004. Emissions subsequently increased to 15,657.29 Gg in 2005, which is approximately 40 percent higher than in 1990, as the levels of peat use for electricity generation returned to former levels with the entry into service of two new stations. Residential fuel combustion (CRF sub-category 1.A.4(b)) is another important source of emissions in the *Energy* sector (Figure 2.1). Although residential energy consumption increased by about 17 percent from 1990 to 2005, the CO₂ emissions in this sub-sector show a decrease of 3 percent due to the decline in the use of carbon-intensive fuels, such as peat and coal, and greater use of oil and natural gas. The emissions of CO₂ from coal and peat use in the residential sector decreased by 61 percent between 1990 and 2005 while those from oil and natural gas more than trebled over this period.

Ireland has only a small number of energy intensive industries and CO₂ emissions from combustion in the industrial sector account for only 8 percent of total emissions but, nevertheless, these emissions increased by approximately 38 percent between 1990 and 2005. The contribution from *Industrial Processes* to total emissions is also relatively small and decreased significantly in 2003 following the closure of Ireland's ammonia and nitric acid plants in June 2002. This reduction is partly offset by sustained increases in emissions of CO₂ from cement manufacture (Figure 2.2). While HFC, PFC and SF₆ emissions account for only 1 percent of the national total, the proportion of these gases in the total for *Industrial Processes* is increasing. The small contribution from the sector *Solvents and Other Product Use* shows a steady decline since 1998.

The component emission trends in *Agriculture* are shown in Figure 2.4 and the main drivers of emissions are shown on Figure 2.8. Large livestock populations produce approximately 500 Gg of CH₄ annually through enteric fermentation and manure management while the sustained application of large amounts of chemical and organic nitrogen to soils results in the emission of approximately 25 Gg N₂O. These emissions from *Agriculture*, equal to approximately 20,000 Gg CO₂ equivalent annually, account for a comparatively larger share of total emissions in Ireland than in most other Annex I Parties. However, according to the estimates given in this NIR, this share decreased from 35 percent in 1990 to approximately 27 percent in 2005 due to the sustained CO₂ increase in *Energy* and a slight downturn in both CH₄ and N₂O emissions from agriculture after 1998, reflecting the decline in the cattle population and fertilizer use.

The full assessment of emissions and removals in the LULUCF sector according to the reporting requirements of Decision 13/CP.9 has given a new understanding of the relative contributions of sub-categories in this sector and it has identified a number of land-use categories that are important in terms of either emissions or removals of CO₂. This sector is a net source of emissions in some years and a net sink of carbon in other years (Table 2.2 and Figure 2.5). This result is determined largely by the balance between 5.A *Forest Land*, which is a major carbon sink, and 5.C *Grassland*, where soil disturbance and liming of

agricultural lands generate relatively large emissions of CO₂. The complex dynamics of land-use changes between categories and the relative contributions from biomass and soils lead to highly fluctuating estimates of emissions and removals over the period 1990-2005.

The *Waste* sector is an important source of CH₄ emissions, the contribution of which is increasing steadily (Figure 2.6) due to the continued dominance of landfill as a means of solid waste disposal in Ireland. The downward shift in the level of emissions after 1996 reflects the effect of landfill gas utilisation at a number of sites since 1996 and a further contribution from gas flaring since 2001.

2.3 Emissions of Indirect Greenhouse Gases

The total emissions of SO₂, NO_x, NMVOC and CO for the years 1990 to 2005 are summarised in Table 2.2. As in the case of CO₂, the emissions of SO₂, NO_x and CO in Ireland are dominated by those emanating from fuel combustion activities while the bulk of VOC emissions are generated by road traffic and solvent use. Substantial decreases have occurred in the emissions of SO₂ and CO. Some reductions have also taken place in NMVOC emissions but emissions of NO_x in 2005 were similar to that in 1990.

Total SO₂ emissions decreased by approximately 61 percent, from 182,839 tonnes in 1990 to 70,402 tonnes in 2005. Power stations remain the principal source of SO₂ emissions, contributing approximately 60 per cent of the total in 2005. Combustion sources in the industrial and residential/commercial sectors largely account for the remainder of emissions, with contributions of 14 percent and 21 percent, respectively in 2005. In 1990, coal combustion accounted for 51 per cent of SO₂ emissions and fuel oil contributed 31 per cent. By 2005, the share of SO₂ emissions from coal had decreased slightly to approximately 50 per cent and that from fuel oil had decreased to 29 per cent.

Table 2.2. Emissions of SO₂, NO_x, VOC and CO 1990-2005 (Tonnes)

	SO ₂	NO _x	NMVOC	CO
1990	182,839	124,228	106,784	410,803
1991	180,017	127,329	107,820	407,101
1992	168,530	135,689	109,042	394,619
1993	159,558	124,256	104,703	362,969
1994	174,095	122,959	103,092	342,835
1995	160,042	124,956	100,949	316,472
1996	148,668	129,173	107,202	326,198
1997	165,350	129,127	107,951	315,948
1998	176,705	134,309	109,825	330,765
1999	157,412	132,106	90,273	301,110
2000	137,162	133,461	81,083	281,872
2001	129,643	134,791	77,678	275,842
2002	99,651	125,671	71,174	254,600
2003	77,949	119,957	67,543	244,953
2004	72,183	118,721	63,939	237,347
2005	70,402	119,088	62,107	225,959

Unlike SO₂, total NO_x emissions show only a small reduction after 2001. Road transport is the principal source of NO_x emissions, contributing approximately 36 per cent of the total in

2005. The power generation sector is the other main source of NO_x emissions, accounting for 27 per cent of emissions in 2005. The reductions in NO_x emissions given by catalytic converters in cars and heavy-duty vehicles have only become apparent in recent years, as the technology has been offset by large increases in vehicle numbers in the past 10 years. This effect is exaggerated in latter years by so-called fuel-tourism, whereby a significant proportion of the automotive fuel sold in Ireland is used by vehicles in the UK and possibly to some extent in other countries but the corresponding emissions are included in the total for Ireland.

The emissions of NMVOC are determined mainly by road traffic and solvent use. These sources typically produce about 70 per cent of the annual total of NMVOC emissions in Ireland. Coal burning in the residential sector is another important source. Technological controls for VOCs in motor vehicles have been more successful than in the case of NO_x, and have given a significant reduction in emissions from road transport over recent years. However, NMVOC emissions from paint application and the domestic use of various solvent-based products are still increasing with the result that overall NMVOC emissions reductions are not large for the period 1990-2005. The emissions of CO continue to decline, driven by major reductions due to catalysts in petrol cars, which is the principal source of CO, and large decrease in the use of solid fuels in residential combustion. Further reductions in the emissions of SO₂, NO_x and NMVOC will occur in the coming years as Ireland implements programmes to comply with the requirements of the National Emission Ceilings Directive (EP and CEU, 2001).

Table 2.1. Greenhouse Gas Emissions 1990-2005 (Gg CO₂ equivalent)

(a) Emissions by Gas

GAS	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
CO ₂ (inc net CO ₂ from LULUCF)	32,673.69	33,672.17	33,607.34	33,586.49	34,666.43	35,662.63	37,447.07	38,851.78	40,551.58	42,137.21	44,992.27	47,361.95	45,776.82	44,847.48	45,552.30	46,635.04
CO ₂ (exc net CO ₂ from LULUCF)	32,552.88	33,399.53	33,286.13	33,430.00	34,683.79	35,480.90	37,139.01	38,622.65	40,688.06	42,288.97	44,883.98	47,342.95	45,902.78	45,146.17	45,746.89	47,292.25
CH ₄	13,286.51	13,474.80	13,555.44	13,667.18	13,667.56	13,722.62	14,031.64	14,102.74	14,331.89	13,919.70	13,438.10	13,260.61	13,268.48	13,881.23	13,338.46	13,102.32
N ₂ O	9,498.85	9,352.58	9,346.12	9,499.28	9,764.81	9,964.97	10,038.59	9,937.06	10,559.77	10,648.13	10,214.20	9,702.05	9,240.36	9,083.32	8,935.99	8,849.96
HFCs	0.69	5.14	5.90	8.90	20.52	44.89	76.23	132.09	189.01	194.82	228.95	251.50	276.55	349.52	384.04	431.03
PFCs	0.09	0.09	0.09	0.09	75.38	75.38	103.09	130.82	61.87	195.93	305.41	295.98	212.40	228.79	186.84	173.95
SF ₆	35.40	36.38	37.36	38.33	81.85	82.83	102.06	132.10	94.24	68.96	55.91	69.49	70.31	118.68	67.09	95.96
Total including LULUCF	55,495.25	56,541.16	56,552.24	56,800.28	58,276.56	59,553.31	61,798.67	63,286.59	65,788.37	67,164.75	69,234.83	70,941.58	68,844.92	68,509.03	68,464.72	69,288.26
Total excluding LULUCF	55,374.43	56,268.52	56,231.04	56,643.80	58,293.92	59,371.58	61,490.61	63,057.46	65,924.84	67,316.51	69,126.54	70,922.59	68,970.89	68,807.72	68,659.31	69,945.48

(b) Emissions by IPCC Category

SOURCE AND SINK CATEGORIES	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
1. Energy	31,603.37	32,588.14	32,538.75	32,739.47	33,802.76	34,676.83	36,324.52	37,520.10	39,808.17	41,528.02	43,683.48	45,944.03	44,634.23	44,913.64	44,814.39	46,384.29
2. Industrial Processes	3,166.43	2,866.26	2,781.45	2,746.12	3,117.73	3,063.04	3,205.31	3,657.35	3,494.73	3,560.02	4,186.67	4,292.79	3,722.18	3,041.57	3,147.14	3,254.70
3. Solvent and Other Product Use	80.86	82.74	82.96	83.42	84.32	86.11	86.79	87.20	88.06	84.75	80.07	79.66	77.37	75.96	75.70	75.44
4. Agriculture	19,063.03	19,235.61	19,286.82	19,488.13	19,652.85	19,857.05	20,247.02	20,362.83	21,024.43	20,594.83	19,535.86	19,128.74	18,889.41	18,983.90	18,830.02	18,453.44
5. LULUCF	120.81	272.64	321.21	156.48	-17.36	181.73	308.06	229.14	-136.47	-151.76	108.29	18.99	-125.96	-298.70	-194.58	-657.22
6. Waste	1,460.75	1,495.77	1,541.04	1,586.65	1,636.26	1,688.55	1,626.97	1,429.97	1,509.45	1,548.90	1,640.45	1,477.36	1,647.69	1,792.65	1,792.06	1,777.61
7. Other	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Total including LULUCF	55,495.25	56,541.16	56,552.24	56,800.28	58,276.56	59,553.31	61,798.67	63,286.59	65,788.37	67,164.75	69,234.83	70,941.58	68,844.92	68,509.03	68,464.72	69,288.26




Figure 2.1 Trend in Emissions from Energy 1990-2005




Figure 2.2 Trend in Emissions from Industrial Processes 1990-2005




Figure 2.3 Trend in Emissions from Solvents and Other Product Use 1990-2005




Figure 2.4 Trend in Emissions from Agriculture 1990-2005




Figure 2.5 Trend in Emissions and Removals from LULUCF 1990-2005




Figure 2.6 Trend in Emissions from Waste 1990-2005




Figure 2.7 Total Primary Energy Requirement 1990-2005




Figure 2.8 Principal Drivers of Emissions from Agriculture 1990-2005

Chapter Three

Energy

3.1 Overview of Energy Sector

The *Energy* source category covers all combustion sources of CO₂, CH₄ and N₂O emissions and the fugitive emissions of these gases associated with the production, transport and distribution of fossil fuels. Table 3.1 presents the CRF Level 3 classification of sources concerned and indicates their degree of coverage in Ireland. Estimates are included for all emission sources that occur in the country and the required level of disaggregation is achieved for sufficiently detailed completion of the CRF tables. The overall approach and methodologies used to estimate emissions in the *Energy* sector for 2005 remain largely as described in the 2006 NIR. However, it was possible to use the CO₂ estimates reported under the ETS for 2005 to achieve complete bottom-up results in respect of some important sub-categories in this sector for the 2005 inventory. This is a significant advance in terms of accuracy as the ETS estimates are verified and they represent a large proportion of the total emissions from *Energy*.

There was particular focus in the 2006 reporting cycle on delivering the best possible estimates of emissions in the base year and on reporting for 2004 in a manner consistent with the base year estimates. Changes in the form and content of the energy balances that were used for 1990 and 2004 in Ireland's 2006 submission were a key part of the work. Similarly improved and revised energy balances are now applied for 2005 and also for the years 1991 through 2003 to give a consistent set of results for all years. Recalculations have therefore been undertaken in this sector for the years 1991-2003 and these are described in section 3.7 below.

Table B.1 of Annex B shows the national energy balance sheets for 2005, published by Sustainable Energy Ireland (SEI). The energy statistics are compiled using a combination of top-down and bottom-up methods and the 2005 example indicates the same form of expanded balance sheet as used for 1990 and 2004. The improved balance sheets reflect revisions made by SEI as part of a programme to harmonise national energy balances in compliance with the needs of the International Energy Agency (IEA) and EUROSTAT and to facilitate their wider use nationally. The Irish energy balances incorporate additional sectoral disaggregation specific to the needs of the greenhouse gas inventory, following close collaboration between SEI and the inventory agency over recent years. A fully consistent set of energy balance sheets for the years 1990-2005 underlies the estimates of emissions for *Energy* in this submission.

Following the methods decision tree of the IPCC good practice guidance for combustion sources, the information in Table B.1 of Annex B allows for the full application of the two available IPCC methods for emission sources in *Energy*, i.e. the Sectoral Approach and the Reference Approach. The Sectoral Approach uses the detailed sectoral breakdown of fuel consumption by all end users as the basis of the calculations for CO₂, CH₄ and N₂O. The relevant activity data are represented by

the disaggregated entries below TPER (Total Primary Energy Requirement) in Table B.1 of Annex B. A combination of top-down and bottom-up methods is used in the sectoral application of the national statistics on fuel consumption to derive the emission estimates in the various sub-categories. The Reference Approach provides an estimate of aggregate CO₂ emissions only, based on the apparent consumption of fuels in the country. This estimate is not used in the compilation of total national emissions but rather for comparison purposes only. The apparent fuel consumption is determined from the energy balance items relating to primary and secondary fuels represented by those above TPER in Table B.1 of Annex B. The application of the Sectoral Approach and the Reference Approach is now described with reference to 2005 data and their results are then compared for CO₂, as required by the UNFCCC reporting guidelines. The Sectoral Approach is described according to the individual sub-categories listed in Table 3.1.

Table 3.1. Level 3 Source Category Coverage for Energy

1 Energy	CO₂	CH₄	N₂O
<i>A. Fuel Combustion</i>			
1. Energy Industries			
a. Public Electricity and Heat Production	All	All	All
b. Petroleum Refining	All	All	All
c. Manufacture of Solid Fuels and Other Energy Industries	All	All	All
2. Manufacturing Industries and Construction			
a. Iron and Steel	All	All	All
b. Non-Ferrous Metals	All	All	All
c. Chemicals	All	All	All
d. Pulp, Paper and Print	All	All	All
e. Food Processing, Beverages and Tobacco	All	All	All
f. Other	NO	NO	NO
3. Transport			
a. Civil Aviation	All	All	All
b. Road Transportation	All	All	All
c. Railways	All	All	All
d. Navigation	All	All	All
e. Other Transportation	All	All	All
4. Other Sectors			
a. Commercial/Institutional	All	All	All
b. Residential	All	All	All
c. Agriculture/Forestry/Fisheries	All	All	All
5. Other	NO	NO	NO
<i>B. Fugitive Emissions from Fuels</i>			
1. Solid Fuels			
a. Coal Mining	NO	NO	NO
b. Solid Fuel Transformation	NO	NO	NO
c. Other	NO	NO	NO
2. Oil and Natural Gas			
a. Oil	NO	NO	NA
b. Natural gas	All	All	NA
c. Venting and Flaring	All	All	NA
d. Other	NO	NO	NO

All : all emission sources covered; NE : emissions not estimated; NO : activity not occurring; NA : not applicable (emissions of the gas do not occur in the source category); IE : emissions included elsewhere

3.2 Sectoral Approach for Emissions from Energy Use

3.2.1 Combustion Sources

The combustion of fossil fuels accounts for the bulk of CO₂ emissions in most countries. In Ireland, fuel combustion contributed almost two-thirds of total emissions in 2005 (Chapter Two). The CO₂ emissions are relatively easy to quantify with reasonable accuracy as the fuel amounts are published annually and information on their carbon contents is well established. The total amount of CO₂ released on combustion can therefore be readily ascertained. Only small amounts of CH₄ and N₂O are associated with fuel combustion activities. The emissions of these gases are generally not quantified with the same reliability as the emissions of CO₂ because the rates of CH₄ and N₂O production depend on several factors, in addition to fuel type, and consequently there is considerable uncertainty in the available emission factors for these gases.

The Irish energy data in the expanded energy balance sheets (Table B.1 of Annex B) are well disaggregated according to fuel and sector for the purposes of calculating emissions in the IPCC Level 3 source categories in a top-down approach. Supplementary sources of information facilitate the use of bottom-up methods in some important sub-categories and they provide greater detail in the overall fuel-sector matrix, making it more compatible with the inventory reporting format required for the Sectoral Approach. The simple calculation spreadsheet given in Table C.1 of Annex C shows how the emissions from combustion sources are computed for the year 2005 using the activity data and emission factors described below. The complete allocation to IPCC Level 1 source categories is readily achieved from this compilation, as shown in Table C.2 of Annex C. The correspondence between the national disaggregation of sources and IPCC combustion source categories is given in Table C.3 of Annex C.

All CO₂ emission factors, except those for petroleum coke and biomass, are country-specific values, determined directly from information on the carbon content and net calorific value of the fuels used in stationary and mobile sources. Where ETS data are used for the 2005 inventory, they provide more precision on CO₂ emission factors on an individual plant basis. The CO₂ emission factor for natural gas now takes into account the increasing contribution of imported gas in the national total given by the energy balance. The import of natural gas from the UK began around 1995 and imported gas accounted for 87 percent of the total in 2005. The CO₂ emission factor appropriate to the split between domestic and imported natural gas, which is more carbon intensive, was first applied for the 2004 inventory. The same approach is again used for 2005 and the emission factors for the years 1995-2003 are produced in the same way for the purposes of recalculations for these years in this submission.

The combustion CO₂ emission factors adopted for use by participants in ETS take account of the fact that a very small fraction of fuel carbon may remain unoxidised and IPCC oxidation factors appropriate to solid, liquid and gaseous fuels are applied to compute the emissions. For other combustion sources, where activity data are in general top-down fuel use from the energy balance, the inventory agency adopts the approach that no specific allowance is needed for unoxidised carbon in the calculation of CO₂ emissions. Default CO₂ emission factors from IPCC are used for petroleum coke and biomass, the latter almost invariably referring to wood wastes. For stationary sources and all mobile sources except road traffic, Ireland has to date relied largely on the default emission factors for CH₄ and N₂O available from the CORINAIR/EMEP Emission Factor Guidebook (McInnes, 1996 and Richardson, 1999).

3.2.1.1 1.A.1 Energy Industries

The returns from ETS participants in respect of their emissions and fuel combustion in 2005 under Directive 2003/87/EC (EP and CEU, 2003) to the EPA's Emission Trading Unit (ETU) were used to compile the complete inventory for category 1.A.1. The data from a total of only 19 individual installations – 16 electricity generating stations in 1.A.1(a), one oil refinery in 1.A.1(b) and two peat briquetting plants under 1.A.1(c) – were sufficient to compute the results in this important category. In each of the three sub-categories, the verified CO₂ estimates reported by the ETS participants were used directly and the corresponding fuel use was used to estimate CH₄ and N₂O emissions using the emission factors as previously applied in these sub-categories.

The bottom-up CO₂ emission estimates received from the ETS participants, along with the emissions of CH₄ and N₂O estimated by the inventory agency, are aggregated on the basis of four main fuel types (peat, coal, oil and natural gas) in the calculation sheets and also by solid, liquid and gaseous fuels for reporting in the CRF. However, the corresponding energy use which appears in the CRF is taken from the energy balance, rather than from the ETS returns, following established practice to always reflect the published national energy data in emission inventories. The differences between ETS energy amounts and those given by the energy balance are very small for 2005 as the inventory agency communicated the ETS data to SEI so that the energy use in respect of category 1.A.1 could be compared and reconciled with that being compiled through SEI's own surveys and procedures.

Figure 3.1 shows the trend in emissions from 1.A.1(a) Public Electricity and Heat Production over the period 1990-2005, which account for 97 percent of the total for category 1.A.1. It may be noted that CO₂ emissions from peat increased in 2005 as peat consumption returned to pre-2004 levels following the construction of two new peat-fired power plants. The emissions from other fuels remained similar to their 2004 levels. Coal is used only at the largest station in the country, which tends to be operated close to capacity giving a corresponding stable CO₂ output. Oil-fired plants are often used to replace downtime at other plants and to meet fluctuating electricity demand with the result that associated emissions are more variable.

One small oil refinery accounts for the emissions reported under 1.A.1(b) Petroleum Refining using ETS data and country-specific emission factors. Emissions for 1.A.1(c) Manufacture of Solid Fuels and Other Energy Industries were reported for the first time in the 2006 submission and refer to the production of peat briquettes from milled peat in two plants. The 2005 values for CO₂ are also taken from ETS returns while CH₄ and N₂O estimates are computed by the inventory agency.

The inventory experts are collaborating with colleagues to fully to consolidate and formalise data gathering for this important sub-set of emissions using prescribed monitoring and verification mechanisms to ensure consistency with reporting under the Convention and Decision 280/2004/EC.




Figure 3.1 Trend in Emissions from 1.A.1 Energy Industries 1990-2005

3.2.1.2 1.A.2 Manufacturing Industries and Construction

The revised and expanded energy balance sheets developed by SEI incorporate a mapping of industrial fuel use in combustion into the CRF sub-categories (a) through (f) under 1.A.2 Manufacturing Industries and Construction. This facilitates the complete disaggregation of emissions in this source category for completion of the CRF Table 1.A(a)s2. In the past, allocation to the lower level was often based on poor information, which resulted in outlier implied emission factors for some of the fuels in sub-categories (a) through (f). Information on emissions or fuel consumption in 2005 was obtained from ETS returns in respect of a small number of energy intensive industries (e.g. alumina production and cement manufacture) allowing their respective estimates to be accounted for on a bottom-up basis. The energy amounts concerned are deducted from the known totals for sub-categories 1.A.2(a) through 1.A.2(f) and the residual fuel quantities is used to compute emissions on a top-down basis using the country-specific emission factors as shown in Table C.1 of Annex C.

3.2.1.3 1.A.3 Transport

The fuel consumption within Ireland associated with sub-category 1.A.3(a) Civil Aviation is calculated from the number of annual Landing and Take-off (LTO) cycles for domestic travel provided by airport authorities, the fuel consumption rates given by the IPCC good practice guidance appropriate to the type of aircraft concerned (Table 2.10, GPG Appendix 2.5A.1) and the length of the flights within Ireland. This approach is used for consistency with other years even though the expanded and updated energy balance sheets record the amount of fuel used in domestic air transport. However, the difference between the two values of fuel amount is less than 10 percent.

Emissions of CO₂ reported under 1.A.3(b) Road Transportation are computed from the amounts of petrol and diesel given under road transport in the energy balance sheet and country-specific emission factors for these fuels as shown in Table C.1 of Annex C. Following the IPCC good practice guidance, the activity data are based on

fuel sales within the State, even though a significant proportion of automotive fuels purchased in Ireland is used in the UK. The CH₄ and N₂O emissions from road traffic are calculated in the COPERT III model (Ahlvik et al, 1997), developed within the CORINAIR programme for estimating a range of emissions from this important source. Road traffic is an important source of N₂O from fuel combustion and the emissions are increasing in line with the increasing share of catalyst-controlled vehicles in the national fleet. The COPERT III model estimates these emissions on the basis of distance travelled using a detailed bottom-up approach (Tier 3) that accounts for such factors as fuel type, fuel consumption, engine capacity, driving speed and a range of applicable technological emission controls that may be applied on the basis of the age of the vehicle. The model is applied annually in Ireland to derive CH₄ and N₂O emissions estimates. The resultant 2005 emission factors have been converted to national average values per fuel type for the purpose of Table C.1 in Annex C.

The CO₂ emissions under 1.A.3(c) Railways and 1.A.3(d) Navigation are calculated from the amounts of oil used by these activities and the country specific emission factors for oil. The emissions reported in sub-category 1.A.3(e) Other Transportation refer to the use of natural gas in pipeline compressor stations. The fuel use is estimated as the difference between the value given for natural gas under own use/losses in the energy balance sheet (Table B.1 of Annex B) and the amount of gas estimated to be lost from the distribution network, as reported under fugitive emissions in sub-category 1.B.2 (b) Natural Gas.

3.2.1.4 1.A.4 Other Sectors

The CRF sub-category 1.A.4 Other Sectors covers combustion sources in the residential, commercial, agriculture and forestry sectors. The residential sector remains the most important source of emissions in this sub-category in Ireland. This is evident from Figure 3.2, which shows the trend in the principal components of emissions in 1.A.4 Other Sectors over the period 1990-2005. While the shift from carbon-intensive fuels, such as coal and peat, to oil and natural gas has been sufficient to maintain emissions relatively constant up to 2005, the benefits from fuel switching are now fully realised and the emissions from oil and gas are increasing in line with higher fuel consumption resulting from greater housing stock and population.

Table B.2 of Annex B shows the calculation of emissions for sub-category 1.A.4 Other Sectors, using the fuel quantities as given by the energy balance (Table B.1 of Annex B). The energy balance provides no indication on the specific end-use of gasoil in the agricultural sector. Consequently, a split based on information from agricultural experts (10 percent stationary sources and 90 percent mobile sources) is used by the inventory agency to distinguish between the use of this fuel in stationary and mobile combustion sources. This split has little bearing on emissions of CO₂, but it is important in relation to CH₄ or N₂O and the indirect greenhouse gases.

3.2.2 Fugitive Emissions

Natural gas has been produced from gas fields off the south coast of Ireland since the 1970s but this source is being rapidly depleted. Substantial reserves of natural gas have recently been discovered off the west coast and they have not yet come into production. Ireland has no coal or oil industries and therefore fugitive emissions of greenhouse gases are limited to those associated with natural gas production and distribution.




Figure 3.2 Trend in Emissions from 1.A.4 Other Sectors 1990-2005

Bord Gais Eireann (BGE), Ireland's gas company has assessed gas losses in the pipeline network in the context of the needs of annual inventory reporting and a long-term programme to replace cast-iron mains with polyethylene pipe in all urban areas served by natural gas. The change to polyethylene pipe is considered to result in negligible losses. The gas company indicated that gas loss in 1995, determined as the difference between system input and metered sales, was 1.92 million therms, which equates to 4,085 tonnes of methane, when the amounts of indigenous and imported gas and their respective properties are taken into account. This value implied a loss of the order of 0.2 percent of total sales. Projections made by BGE for five-year intervals from 2000 show losses decreasing to zero by 2020 on completion of the pipe replacement programme.

The BGE data continue to be used as the best available for this particular fugitive emission source. The rate of loss implied by the 1995 value and the projections is applied to give an emission for all years of the inventory time-series referred to in this report. The gas consumption recorded in the energy balance for the industrial, commercial and residential sectors is used as activity data rather than total sales and the appropriate split between indigenous and imported gas is applied for all years. The inventory agency was informed by BGE in 2004 that natural gas losses from the distribution network were so small that they could not be measured.

Only one company is involved in natural gas production in Ireland. Emissions to the atmosphere from this company's off-shore gas production platforms are reported to the Department of Communications Marine and Natural Resources under the OSPAR Convention. Such reports have been obtained for several years in the 1990-2005 time series and the estimates of CO₂ and CH₄ emissions given therein are used directly for the years concerned. The available data, which relate largely to gas extraction but which also account for a small amount of flaring in some years, indicate a close relationship between emissions and the amount of gas produced.

This relationship has been applied in terms of the indicative emission rates of CO₂ and CH₄ per unit of gas extracted to estimate the emissions for those years for which no reports were received. A report on emissions was supplied for 2005.

3.3 IPCC Reference Approach for CO₂ Emissions from Energy Use

The IPCC Reference Approach is a top-down methodology for CO₂ that estimates emissions by accounting for the overall production of primary fuels, the external trade in primary and secondary fuels, stock changes and for the carbon that may enter long-term storage in non-energy products and feedstocks. It can be used to report national emissions in cases where the detailed activity data required for the Sectoral Approach are not available but it is more usually applied for verification of the results of the latter for those countries that have the information to apply both methods. The Reference Approach is used in Ireland as a verification procedure for CO₂ emissions from fuel combustion activities. The calculation sheet for the Reference Approach (Table 1.A(b) of the 2005 CRF) is reproduced as Table C.4 of Annex C of this report. The apparent consumption of fuels, the basic activity data in this case, is determined as

$$\text{Apparent Consumption} = \text{Production} + \text{Imports} - \text{Exports} - \text{International Bunkers} - \text{Stock Changes}$$

where production applies only to primary fuels. Naphtha was previously the only petroleum product to be considered in relation to non-energy fuel-use, where the carbon is not fully released as in combustion. The IPCC default value of 0.75 is used for the proportion of carbon stored in naphtha. Ireland's only oil refinery is a small hydroskimming refinery where there is no production of other petroleum products normally used for non-energy purposes, such as bitumen, lubricants, plastics and asphalt. The expanded SEI energy balance sheets now record the import of some of these products, thereby allowing improved completeness in the Reference Approach estimation of CO₂ emissions and carbon storage. A significant amount of natural gas feedstock was traditionally used in ammonia production in Ireland but the company closed in 2002 and there is consequently no feedstock use of natural gas since 2002.

3.4 Comparison of CO₂ Estimates from the Sectoral Approach and Reference Approach

The national energy consumption and CO₂ emissions estimates obtained using the Sectoral Approach usually differ to some extent from the corresponding values resulting from the Reference Approach (Table C.5 of Annex C). According to the UNFCCC guidelines, differences greater than 2 percent should be explained and investigated to see whether they indicate systematic underestimation or overestimation of energy consumption by one or other of the methods. Differences of 2.6 percent and –0.6 percent are indicated for total energy and CO₂ emissions, respectively in 2005. The differences are largely due to those for liquid fuels, where they amount to 4.77 percent for energy and –0.78 percent for CO₂ emissions. The small differences are largely explained by the highly aggregated manner in which emission factors for liquid fuels are applied in the Reference Approach and the effect of non-energy fuel use.

3.5 Memo Items

The memo items of the IPCC reporting format refer to activities for which the emissions are excluded from national totals. The use of fuels in international aviation and marine bunkers is the most important of these activities. Some of the associated emissions, particularly CO₂ emissions from international aviation, are increasing very rapidly and it is therefore important that they are closely monitored for comparison with other sources and for the benefit of the international organisations that will have to develop control strategies for them in the future. The emissions of CO₂ from biomass combustion are not included in national totals of greenhouse gases because it is assumed that an equivalent amount of CO₂ is removed from the atmosphere by the growth of the next biomass crop. The estimation of emissions for memo items is described here because they are calculated as part of the general estimation procedures for the Energy sector.

The activity data for biomass appear as a specific item in the Irish energy balance sheets (Table B.1 of Annex B). For the industrial, residential and agricultural sectors, this is known to refer to wood wastes. Default emission factors for CO₂, CH₄ and N₂O for wood burning are used to estimate the emissions from biomass in these sectors using the simple Tier 1 approach. The estimates for all gases appear in the CRF tables covering these sectors, but in the case of CO₂, they do not contribute to the total for Energy or to the national total in the CRF summary tables.

The national energy balance sheets include marine bunkers and international aviation as specific items and the emissions may be calculated directly. The approach used to estimate fuel consumption in domestic civil aviation by the inventory agency is described in section 3.2.1.3 above and gives a result for 2005 close to that in the energy balance. This fuel amount is deducted from the value given in the energy balance sheet for kerosene use in air transport to obtain an estimate of international aviation bunker fuel consumption. In 2005, the amount of fuel allocated to domestic aviation was approximately 4.2 percent of the total recorded under air transport in the energy balance.

3.6 QA/QC in Energy

Extensive QA/QC procedures have been followed for the Energy sector during the present reporting cycle by fully implementing the plan that underpins Ireland's formal national system. This involved the designation of a QA/QC manager within the inventory agency and the application of a system of quality control checks and documentation spreadsheets to the front of all calculation workbooks. These workbooks were restructured to correspond directly to the disaggregation given by the CRF sectoral background data tables and so that calculations may be made on a time-series basis, rather than by individual year. This increases efficiency in the use of the time-series energy data provided by SEI, allows for rapid recalculation and checking across the time-series and facilitates the transfer of the output emission estimates and energy quantities to the CRF Reporter Tool. Additional summary sheets are used for aggregation to various levels to provide full cross-checking with completed CRF tables for any year.

The quality checks at inventory level build on the extensive upgrading and quality control of energy balances now largely completed by SEI. This work, together with further collaboration with inventory experts and thorough evaluation of the SEI role in relation to the national system and QA/QC procedures, has resulted in substantial

improvements that are now taken into account in the emissions for Energy for the years 1990 through 2005 included in the present submission. In recognition of its role as a key data provider, SEI is continuing to develop its own procedures to ensure that energy balances fully harmonised with Eurostat and IEA requirements will be made available in a timely manner to facilitate the annual reporting of greenhouse gas emissions estimates. Arrangements have been established whereby the bottom-up energy data reported to the EPA for individual enterprises in all relevant energy-use sectors covered by the EU emissions trading scheme may be reconciled at an early stage with the corresponding top-down information collected by SEI. Plans are being formulated to supplement traditional approaches to energy accounting by undertaking surveys of greater frequency and increased bottom-up detail.

The formal application of ETS data in the Energy sector for 2005 is a timely and important step towards improved reliability and accuracy of the estimates. Thorough checking of this input was achieved in collaboration with colleagues in the Emissions Trading Unit of the EPA. Having received the raw ETS data from ETU, the inventory experts allocated the CO₂ estimates and corresponding energy amounts to the appropriate sub-categories for CRF reporting and then returned the compilation to the ETU contact for final checking and accounting of any amendments following the ETS verification process.

The internal review of key categories and other issues is part of the QA/QC system used for the EU greenhouse gas inventory. During the 2007 reporting cycle, inventory experts in Ireland participated in the review of category 1.A.1. The exercise was useful as a means of comparing the methods and data used in the Member States and it shows that the inventories have reached an advanced stage of development in all countries and that a high degree of consistency and comparability is being achieved.

3.7 Recalculations in Energy

Recalculations have been undertaken in the Energy sector to account for the following

- Application of the revised and expanded energy balance for years 1991-2003;
- Inclusion of estimates for the sub-category *1.A.1(c) Manufacture of Solid Fuel and Other Energy Industries* for years 1991-2003;
- Reallocation of all heavy fuel oil consumption from the commercial sector under *1.A.4 Other Sectors* to *1.A.2 Manufacturing Industries and Construction*;
- Upward revision of coal consumption in *1.A.2 Manufacturing Industries and Construction* in 2003 and 2004 following comparison of energy balance data and ETS energy data for 2005.

The updated and expanded energy balances were not available to the inventory agency in time to complete the various recalculations for all years in the time-series for reporting in 2006. This work together with some other minor modifications is now reflected in the results of the recalculations as given in Table 3.2. The effects are very small in most years.

Table 3.2. Percentage Change in Emissions from Energy due to Recalculations

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
Sub-category total in 2006 Submission (Gg CO₂ equivalent)															
1.A.1. Energy Industries	11,099.29	11,572.35	12,248.31	12,256.14	12,601.19	13,268.56	13,991.19	14,664.34	15,032.71	15,686.74	15,987.06	17,196.14	16,267.87	15,598.85	15,218.91
1.A.2. Manufacturing Industries and Construction	4,112.24	3,842.26	3,619.81	3,598.98	3,708.70	3,527.72	3,511.76	3,987.60	3,917.48	4,237.75	4,743.34	4,726.29	4,892.17	4,784.71	4,710.24
1.A.3. Transport	5,035.61	5,265.07	5,682.19	5,667.32	5,882.58	6,368.98	7,127.86	7,752.89	8,847.78	9,818.64	10,210.97	11,062.84	11,230.63	11,392.65	12,092.64
1.A.4. Other Sectors	9,997.67	9,639.63	9,388.32	9,049.14	9,548.75	9,316.93	9,091.58	9,420.72	9,973.98	9,902.73	10,364.47	10,413.97	10,296.49	10,262.95	10,594.83
1.B. Fugitive Emissions from Fuels	138.90	142.30	140.75	159.91	162.65	166.80	160.73	141.02	107.85	117.06	70.82	134.00	65.40	58.88	71.30
1. A. Total	30,383.71	30,461.61	31,079.37	30,731.50	31,903.86	32,648.99	33,883.11	35,966.57	37,879.80	39,762.91	41,376.65	43,533.24	42,752.56	42,098.05	42,687.92
Subcategory total in 2007 Submission (Gg CO₂ equivalent)															
1.A.1. Energy Industries	11,158.62	11,617.34	12,279.74	12,297.59	12,634.28	13,317.47	14,031.86	14,692.87	15,080.53	15,732.98	16,050.38	17,266.57	16,345.34	15,643.44	15,283.51
1.A.2. Manufacturing Industries and Construction	3,969.76	4,101.01	3,785.02	3,994.43	4,259.72	4,365.44	4,172.23	4,588.25	4,599.87	4,748.00	5,682.10	5,660.99	5,371.49	5,371.09	5,444.89
1.A.3. Transport	5,045.02	5,245.42	5,694.65	5,653.42	5,891.79	6,123.46	7,162.07	7,504.93	8,847.83	9,810.93	10,535.34	11,041.74	11,237.21	11,419.24	12,101.91
1.A.4. Other Sectors	10,064.90	10,198.47	9,377.33	9,354.88	9,523.50	9,374.17	9,413.81	9,158.46	9,626.52	9,507.14	9,681.31	10,068.53	9,935.22	10,232.99	10,260.41
1.B. Fugitive Emissions from Fuels	138.88	142.34	140.78	159.92	162.67	166.75	160.75	141.04	107.85	117.06	70.82	134.00	65.40	58.88	71.30
1. A. Total	30,377.18	31,304.58	31,277.51	31,460.23	32,471.95	33,347.29	34,940.73	36,085.55	38,262.60	39,916.11	42,019.95	44,171.82	42,954.66	42,725.64	43,162.02
Percentage Change in Emissions from Energy															
1.A.1. Energy Industries	0.53	0.39	0.26	0.34	0.26	0.37	0.29	0.19	0.32	0.29	0.40	0.41	0.48	0.29	0.42
1.A.2. Manufacturing Industries and Construction	-3.46	6.73	4.56	10.99	14.86	23.75	18.81	15.06	17.42	12.04	19.79	19.78	9.80	12.26	15.60
1.A.3. Transport	0.19	-0.37	0.22	-0.25	0.16	-3.85	0.48	-3.20	0.00	-0.08	3.18	-0.19	0.06	0.23	0.08
1.A.4. Other Sectors	0.67	5.80	-0.12	3.38	-0.26	0.61	3.54	-2.78	-3.48	-3.99	-6.59	-3.32	-3.51	-0.29	-3.16
1.B. Fugitive Emissions from Fuels	-0.01	0.02	0.02	0.00	0.01	-0.03	0.01	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1. A. Total	-0.02	2.77	0.64	2.37	1.78	2.14	3.12	0.33	1.01	0.39	1.55	1.47	0.47	1.49	1.11

Chapter Four

Industrial Processes

4.1 Overview of Industrial Processes Sector

The list of activities under *Industrial Processes* in the IPCC reporting format is given in Table 4.1. Some of these activities are well known sources of one particular greenhouse gas, such as cement production for CO₂ or adipic acid production in the case of N₂O, while others may be more important in terms of their indirect greenhouse gas emissions. Major industrial processes within the chemical sector and metal production that are common to many other developed countries have never been an important part of the Irish economy. Consequently, many of the production processes listed in Table 4.1 are not relevant to the inventories of greenhouse gases in Ireland. The four industrial sources that have been covered in the past, mainly due to their emissions of CO₂, are cement and lime production under *2.A Mineral Products* and ammonia and nitric acid production under *2.B Chemical Industry*. The ammonia and nitric acid plants, both operated by Irish Fertilizer Industries, ceased production in June 2002. A small amount of limestone is used to abate SO₂ emissions in peat-fired electricity generating stations and thus *2.A.3 Limestone and Dolomite Use* is a relevant activity in Ireland. The associated CO₂ emissions from this minor source were included in the annual inventories for the first time in the 2006 submission.

The *Industrial Processes* source category is the only IPCC Level 1 category for which emissions of HFC, PFC and SF₆ are reported in annual inventories. Both potential and actual emissions of the 21 individual substances concerned (Table A.1, Annex A) should be reported for source category *2.F Consumption of Halocarbons and SF₆* while actual emissions only are required in other source categories (*2.C Metal Production* and *2.E Production of Halocarbons and SF₆*.) The IPCC methods estimate potential emissions by equating emissions to total consumption while actual emissions are the estimated losses to air of the substances concerned. There is no production of halocarbons or SF₆ in Ireland and therefore source category *2.F Consumption of Halocarbons and SF₆* is the only relevant source category of HFC, PFC and SF₆ emissions in the country. All relevant sub-categories are fully covered (Table 4.1), as described below.

Table 4.2 presents the estimates of greenhouse gas emissions for *Industrial Processes* over the period 1990-2005 for the relevant sources in Ireland. They indicate contributions of 5.7 percent and 4.7 percent to total emissions in 1990 and 2005, respectively. As such the sector is not a particularly important one in the Irish greenhouse gas inventories. Ammonia and nitric acid production were the principal sources of emissions in the sector in 1990, accounting for two-thirds of the total, but the plants ceased operation in 2002, leaving cement production as the dominant process emission source thereafter. The combined contributions of HFC, PFC and SF₆ to the total emissions for *Industrial Processes* remains small and highly variable from year to year. Emissions of HFC show a steady increase up to 2005, largely due to the influence of the air conditioning and refrigeration sub-categories while the emissions of PFC continue to follow the downward trend evident in previous submissions. Emissions of SF₆ remain variable from year to year. The estimates given in Table 4.2 for the period 1990-2005 include recalculated values in several sub-categories for many of these years, which are further described in the following sections.

Table 4.1 Level 3 Source Category Coverage for Industrial Processes

2. Industrial Processes	CO ₂	CH ₄	N ₂ O	HFC	PFC	SF ₆
A. Mineral Products						
1. Cement Production	All	NA	NA	NA	NA	NA
2. Lime Production	All	NA	NA	NA	NA	NA
3. Limestone and Dolomite Use	All	NA	NA	NA	NA	NA
4. Soda Ash Production and Use	NE	NA	NA	NA	NA	NA
5. Asphalt Roofing	NE	NA	NA	NA	NA	NA
6. Road Paving with Asphalt	NE	NA	NA	NA	NA	NA
7. Other	NO	NO	NO	NO	NO	NO
B. Chemical Industry						
1. Ammonia Production*	All	NE	NA	NA	NA	NA
2. Nitric Acid Production*	NA	NA	All	NA	NA	NA
3. Adipic Acid Production	NO	NO	NO	NA	NA	NA
4. Carbide Production	NO	NO	NA	NA	NA	NA
5. Other	NO	NO	NO	NO	NO	NO
C. Metal Production						
1. Iron and Steel Production	NO	NO	NO	NA	NA	NA
2. Ferroalloys Production	NO	NO	NO	NA	NA	NA
3. Aluminium Production	NO	NO	NO	NA	NA	NA
4. SF ₆ Use in Aluminium and Magnesium	NA	NA	NA	NA	NA	NO
Foundries						
5. Other	NO	NO	NO	NO	NO	NO
D. Other Production						
1. Pulp and Paper	NE	NE	NE	NA	NA	NA
2. Food and Drink	NE	NE	NE	NA	NA	NA
E. Production of Halocarbons and SF ₆						
1. By-product Emissions	NA	NA	NA	NO	NO	NO
2. Fugitive Emissions	NA	NA	NA	NO	NO	NO
3. Other	NA	NA	NA	NO	NO	NO
F. Consumption of Halocarbons and SF ₆						
1. Refrigeration and Air Conditioning Equipment	NA	NA	NA	All	All	All
2. Foam Blowing	NA	NA	NA	All	All	All
3. Fire Extinguishers	NA	NA	NA	All	All	All
4. Aerosols/ Metered Dose Inhalers	NA	NA	NA	All	All	All
5. Solvents	NA	NA	NA	All	All	All
6. Semiconductor Manufacture	NA	NA	NA	All	All	All
7. Electrical Equipment	NA	NA	NA	All	All	All
8. Other	NA	NA	NA	All	All	All
G. Other	NA	NA	NA	NO	NO	NO

All : all emission sources covered; NE : emissions not estimated; NO : activity not occurring; NA : not applicable (emissions of the gas do not occur in the source category); IE : emissions included elsewhere

* ammonia and nitric acid plants closed down in June 2002

4.2 Emissions from 2.A Mineral Products

The IPCC Level 3 emission source categories relevant under *2.A Mineral Products* in 2005 are limited to *2.A.1 Cement Production*, *2.A.2 Lime Production* and *2.A.3 Limestone and Dolomite Use*. Total CO₂ emissions from these activities amounted to 2,533.76 Gg, of which cement production accounted for 92 percent.

4.2.1 2.A.1 Cement Production

During the cement manufacturing process, calcium carbonate in the cement kiln feed (typically CaCO_3 in limestone) undergoes calcination at high temperature to produce lime (CaO) and CO_2 . The activated lime that results from this process combines with silica in the kiln feed to form cement clinker. The emissions of CO_2 are usually calculated from the amount of clinker produced and the stoichiometric ratio of CO_2 to CaO . A small amount of raw material may be converted into cement kiln dust (CKD) due to incomplete calcination. If the CKD is not recycled as part of subsequent kiln input, the CO_2 emissions based on clinker production must be corrected to account for the carbonate fraction lost in CKD.

An emission factor of 0.5 tonne CO_2 per tonne of cement clinker was used to estimate CO_2 emissions from cement production for all years up to 2002 using the Tier 1 method. Information on CO_2 emissions from cement production supplied to the EPA during 2004 for the development of Ireland's National Allocation Plan under Directive 2003/87/EC (EP and CEU, 2003) on emissions trading provided a basis for re-examination of both the historical activity data and the emission factor used to estimate CO_2 from cement production for the years 1990-2002. The new information was obtained from a number of additional cement producers who had entered the Irish market in 2000, in addition to the single larger original manufacturer.

As the emission estimates from individual cement plants are subject to verification under Directive 2003/87/EC, their validity is being fully established in the context of the companies' documented methods and data and the associated guidance on emissions estimation methods provided by Decision 2004/156/EC (CEC, 2004). Such verification was achieved in respect of the four cement plants in operation in 2005, allowing for more accurate accounting of combustion emissions and process emissions separately. The process CO_2 emissions from these plants were calculated using the Tier 2 method given in Decision 2004/156/EC, which is fully consistent with the Tier 2 method in the IPCC good practice guidance, based on reliable data on clinker production, corrected as appropriate for CKD, and CaO content of the clinker. The emission factors in 2004 ranged from 0.533 t CO_2 /t clinker to 0.540 t CO_2 /t clinker with a weighted average of 0.536 t CO_2 /t for all clinker production. The procedure was repeated for 2005, giving process emissions of 2,357.76 Gg CO_2 and an implied emission factor of 0.536 t CO_2 /t clinker.

4.2.2 2.A.2 Lime Production

Statistical data on lime production in Ireland are obtained annually from the lime manufacturers (three companies up to 1998 and two companies thereafter). The CORINAIR default value for CO_2 emissions from lime production (0.75 t CO_2 /t lime) was used consistently to estimate process emissions from this source using the Tier 1 method for all inventory years up to 2003. This default value is also given for high-calcium lime in the IPCC good practice guidance and it was considered appropriate for Ireland as high-grade limestone is the standard raw material available for high calcium quicklime manufacture (at least 95 percent CaO content).

As in the case of cement production, lime producers have provided their own estimates of CO_2 emissions from lime manufacture calculated in accordance with the methods described in Decision 2004/156/EC, thus enabling the inventory agency to review and revise the previously submitted estimates for another important source of CO_2 emissions in *Industrial Processes*. The CO_2 estimates have been obtained for all years up to 2005, as given in Table 4.2. They indicate implied emission factors in the range 0.75 to 0.88 t CO_2 /t lime produced.

Table 4.2. Emissions from Industrial Processes 1990-2005

	Gas	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
INDUSTRIAL PROCESSES	All	Gg CO ₂ eq	3,166.43	2,866.26	2,781.45	2,746.12	3,117.73	3,063.04	3,205.31	3,657.35	3,494.73	3,560.02	4186.671	4,292.79	3,722.18	3,041.57	3,147.14	3,254.70
2.A. Mineral Products																		
Cement Production	CO ₂	Gg	884	782	753	729	859	879	983	1,145.00	1,059.00	1,166.00	1,700.90	1,851.20	1,859.80	2,127.00	2,295.10	2,357.10
Lime Production	CO ₂	Gg	214.1	192.2	162.4	204.9	205.4	187.5	198.2	221.9	211.7	170.1	190.4	189.4	190.3	206.2	201.5	183.6
Limestone and Dolomite Use	CO ₂	Gg	7.594	7.394	7.294	6.994	7.294	7.994	7.794	8.794	8.644	8.944	9.342	13.479	10.954	11.368	12.55	13.143
2.B. Chemical Industry																		
Ammonia Production*	CO ₂	Gg	989.2	1030.6	1003	945.5	1055.8	973.0	922.5	1074.2	1058.1	943.1	883.3	1037.4	809.7	NO	NO	NO
Nitric Acid Production*	N ₂ O	Gg CO ₂ eq	1035.4	812.2	812.2	812.2	812.2	812.2	812.2	812.2	812.2	812.2	812.2	585.9	291.4	NO	NO	NO
2.F. Consumption of Halocarbons and SF₆																		
Emissions of HFC	HFC	Gg CO ₂ eq	0.693	5.133	5.891	8.887	20.525	44.889	76.226	132.088	189.01	194.825	228.945	251.504	276.547	349.517	384.041	431.026
Emissions of PFC	PFC	Gg CO ₂ eq	0.093	0.093	0.093	0.093	75.382	75.382	103.085	130.823	61.87	195.933	305.406	295.984	212.403	228.795	186.84	173.954
Emissions of SF ₆	SF ₆	Gg CO ₂ eq	35.405	36.381	37.357	38.334	81.852	82.827	102.062	132.1	94.245	68.956	55.907	69.485	70.308	118.682	67.093	95.962
Implied Emission Factor for CO ₂ in Cement Production		t CO ₂ /t clinker	0.549	0.549	0.549	0.549	0.549	0.549	0.549	0.549	0.549	0.549	0.547	0.547	0.545	0.536	0.536	0.536
Implied Emission Factor for CO ₂ in Lime Production		t CO ₂ /t lime	0.839	0.83	0.753	0.843	0.832	0.839	0.826	0.877	0.826	0.823	0.767	0.778	0.785	0.861	0.819	0.814

* Ammonia and nitric acid plants ceased production in 2002

4.2.3 2.A.3 Limestone and Dolomite Use

Up to 2006 Ireland had not previously reported emissions arising from this activity. Information became available in 2005 to allow for the inclusion of CO₂ emissions associated with the use of carbonates in the manufacture of building bricks and ceramics and from limestone use for the absorption of SO₂ emitted in one new peat-burning power station. The emissions from these sources are rather small, amounting to 0.4 percent of the total for *Industrial Processes* in 2005 but nevertheless their inclusion adds further to the level of inventory completeness.

Information on the raw materials used in brick manufacture (clay, carbonates and shale) has been supplied for the years 1990-2005 by three companies who are participants in the EU emissions trading scheme. Carbon dioxide emissions estimates from the three individual companies are used in inventory calculations, similar to the approach undertaken in *2.A.1 Cement Production*.

Limestone has been used to capture sulphur emitted from peat burning in one new electricity generating station since 2001. The CO₂ emissions from this use of limestone are estimated on the basis of limestone quantity reported by the company and an emission factor of 0.44 t CO₂/t limestone, which is the stoichiometric ratio of CO₂ to CaCO₃. The reported emissions for *2.A.3 Limestone and Dolomite Use* refer to the manufacture of bricks and ceramics only up to the year 2000 and thereafter also include the emissions from limestone use in the new peat-fired power plant.

4.3 Emissions of HFC, PFC and SF₆ from *Industrial Processes*

4.3.1 Special Studies

The compilation of emissions estimates for fluorinated gases presents major new challenges for inventory agencies because they emanate from diverse sources that are entirely different to those traditionally covered by atmospheric emissions inventories and the uses of many of the substances concerned are continuing to change very rapidly in the marketplace. Issues of confidentiality are common among many of the source activities concerned and this also hinders the inventory process and the transparency of reporting in relation to fluorinated gases. The first attempts to quantify emissions of HFC, PFC and SF₆ in Ireland were made for the year 1995 for inclusion in Ireland's Second National Communication published in 1997 (DOE, 1997). Little was known at that time about the sources of these emissions and the methodologies to quantify them were not well established. The results for 1995 were therefore regarded as tentative and incomplete. However, the indications were that, in common with emissions from industrial processes in general in this country, those of HFC, PFC and SF₆ were likely to be rather small.

In 2000, the EPA commissioned special studies on HFC, PFC and SF₆ emissions, led by the Clean Technology Centre (CTC) at Cork Institute of Technology that were designed to identify the important sources in Ireland and to quantify the emissions in 1998 on the basis of separate bottom-up and top-down methodologies. The reports on these studies (O'Doherty and McCulloch, 2002 and O'Leary *et al*, 2002) describe a very comprehensive investigation into the emissions of fluorinated gases in Ireland and the bottom-up method provided a readily applicable approach that could be used for developing inventories of these gases for other years. The bottom-up approach took full account of the available IPCC methodologies and IPCC good practice guidance in developing the 1998 emissions estimates for HFC, PFC and SF₆. Tier 2 methods were used for estimating the emissions from the majority of sources that have non-zero emissions. The actual and potential emissions in 1998 were

compiled in the CRF tables, with table modifications where necessary to facilitate transparent reporting of the country-specific data.

The methodological approach adopted in the special study for 1998 was subsequently used in early 2002, again under contract with CTC (O'Leary, 2002), to compile emissions estimates for HFC, PFC and SF₆ for the time-series 1995 through 2000, which were incorporated in the recalculated inventories submitted in 2002. Estimates were also compiled to the extent possible at that time for 1990, but data were difficult to obtain and it was clear that the use of many of the substances had not become established in the country by then. The focus in this particular follow-up study was on the years from 1995 to 2000, in the knowledge that 1995 could be selected as the base year for emissions of fluorinated gases. The inventory agency subsequently continued reporting for the years up to 2003, based broadly on the CTC approach used for the 1995-2000 time-series.

4.3.2 HFC, PFC and SF₆ Time-Series 1990-2005

As part of the work on the 2004 inventory and the general round of improvements conducted for the 2006 submission, the inventory agency decided that it would be useful to again examine, on a contract basis, the known sources of HFC, PFC and SF₆ emissions over an extended time period. The contract was undertaken jointly by CTC and UK consultants NETCEN, the latter having considerable experience in developing emission inventories for the UK. The work and results are fully described in a supplementary document (Adams *et al*, 2005). The intention was to re-assess the use and application of the various substances in the Irish market as a whole, initially to compile the best possible estimates of emissions in 2004, and to make revisions as appropriate for earlier years based on better information, particularly for 1995 (the base year adopted by Ireland with respect to HFC, PFC and SF₆) and for those years (2001-2003) for which the estimates had been produced by the inventory agency. A second objective of the study was to extend the F-gas emissions time-series back to 1990 so that Ireland could make available information that had been lacking for the years 1990-2004, requested under Decision 280/2004/EC, to enable the European Union to complete the inventories at the European level for all years. In performing this update of the previous emission inventories for fluorinated gases, a number of users and distributors of the fluids were contacted and any data obtained were used for estimating emissions of the various gases for the period 2001-2004. Where data has allowed, emission estimates have been calculated following the sectoral guidance provided by the Intergovernmental Panel on Climate Change (IPCC) publications. Where changes in methodology have occurred in any sector, or where new or corrected information for previous years has become available, emissions have been recalculated for the entire time-series 1995-2001.

Emissions from the consumption of HFC, PFC and SF₆ i.e. through the production, use and disposal of equipment containing these fluids (e.g. refrigerators, electrical switchgear etc) remain the only relevant emissions in Ireland. The following paragraphs describe the overall approach and assumptions made to achieve the estimates of both actual and potential emissions with reference to the relevant source categories, as identified by the earlier 2002 studies. Reference should be made to the special study report for further clarification on those sources actually covered and those that have non-zero emissions.

In the consumption of HFC, PFC and SF₆, a distinction can be made between uses as aerosols and fire-fighting equipment that release fluid at the near-instantaneous time it is used and applications such as mobile air conditioning and foams where the release happens slowly over its lifetime and on disposal. The IPCC guidance recommends that, where information is available, the whole product life cycle be considered when estimating emissions, including imported and exported goods. For example, production-related

emissions are typically estimated from the amount of fluid charged into the equipment during manufacture multiplied by a percentage loss rate. In this study, where this data is available, the production levels and loss factors are generally based on the information received from manufacturers, and IPCC default values are used in other cases.

For many applications, the major form of emissions is gradual loss from a bank of fluid contained in a large number of similar products. This may occur through leakage or through equipment failure. The applications subject to loss in this way include refrigeration and air conditioning equipment, closed-cell foams, fire extinguisher systems and electrical switch-gear. The annual emissions from these types of equipment can be calculated as the sum of the manufacturing, usage and disposal emissions for all the different types of equipment within a specific sector.

The IPCC Good Practice Guidance (IPCC, 2000) proposes a Tier 3b 'top down' approach and a Tier 3a 'bottom up' approach. It is clear however, as is the case for many other countries including the U.K., that data availability in Ireland means that it is not possible to use Tier 3b for all sources. In order to apply Tier 3b it is necessary to obtain complete, accurate and precisely defined data. This is most likely to be possible when there are a small number of operators with well-established data collection systems in a given sector (e.g. in electrical switchgear use), but in contrast for sectors such as the refrigeration and air conditioning sector, there are large numbers of small and medium-sized service companies in addition to the main distributors and equipment manufacturers. Hence, the likely accuracy and completeness of the data are not sufficient to apply Tier 3b to all relevant sectors in Ireland. Instead a pragmatic approach has been used in sectors where lack of data precludes the use of a full Tier 3b methodology. Instead, the use of an alternative Tier 1 methodology for estimating emissions as described in the Good Practice Guidance is used in several sectors.

In the 2005 contract study (Adams et al, 2005), emissions of HFC, PFC and SF₆ have been estimated from the consumption of halocarbons and SF₆ for the following activities:

- 2.F.1 Stationary Refrigeration and Air Conditioning;
- 2.F.2 Foams;
- 2.F.3 Fire-extinguishers;
- 2.F.4 Aerosols and Metered Dose Inhalers;
- 2.F.7 Semiconductor manufacture;
- 2.F.8 Electrical equipment;
- 2.F.9 Other, which includes emissions from window sound-proofing, medical applications, sporting goods and as a gas-air tracer in research and leak detection.

The category *2.F.9 Other* includes estimates for SF₆ emissions from its use in medical applications and sporting goods. These two sources were noted as potential emission sources in the 2002 inventory study but the actual emissions were not quantified. The inclusion of emission estimates for these uses, although both relatively minor in terms of the level of their respective emissions, therefore helps to improve the completeness of Ireland's inventory.

The results of the inventory update for F-gases are shown in Table 4.3. They clearly indicate that for most sectors emissions of HFC, PFC and SF₆ have generally increased year on year. This trend reflects the increasing use of these fluorinated species across a range of applications (e.g. often as replacements in applications where the use of CFC and HCFCs is no longer permitted under the Montreal Protocol) and hence the presence of larger fluid banks from which operational leakage potentially occurs. One exception to this trend is in the manufacture of semiconductors, where the reported emissions received from manufacturing companies in Ireland show annual fluctuations reflecting changing manufacturing activity in

response to the global trends in this market. For the years 1999-2005, this sector produced much higher emissions of HFC, PFC and SF₆ than any other (in terms of CO₂ equivalent emissions), accounting for half of the annual emissions in these years.

4.4 Recalculations for *Industrial Processes*

The recalculations for *Industrial Processes* being submitted in 2007 may be summarised as set out below. Most of the changes relate to the methods and data used for the individual source categories of HFC, PFC and SF₆. Specifically there was a inconsistency between the UK population data which is used in some cases to derive emission estimates for Ireland on the basis that Ireland and the UK have for example similar purchasing trends as is the case in recalculations in 2.F.4 Aerosols and 2.F.9 Other-sporting goods. Table 4.4 lists the previously reported estimates and recalculated values of emissions for Industrial Processes. Plant-specific data is used to calculate process CO₂ emissions from 2.A.1 Cement Production for the years 1990-2005. As a result of a review as part of Irelands QA/QC procedures under the National Inventory System, data for 2003 and 2004 has been revised. Irish HFC emissions from aerosols are derived from the BAMA estimate for the UK, using population as the scaling factor, however the UK population estimates used to calculate this scaling factor were inconsistent with population estimates used to calculate scaling factors in Chapter 5 Solvent and Other Product Use. To maintain consistency one single estimate of UK population is now in use over the time series 1990-2005 and has been verified against published statistics by the UK Office for National Statistics. Similarly, to maintain consistency in the use of scaling up factors on the basis of circumstances in the UK, there are recalculated emission estimates in respect of medical applications and sporting goods under 2.F.9 Other.

4.6 Improvements in *Industrial Processes*

The inventory agency will continue to use verified CO₂ emissions estimates for cement and lime production that become available through the EU emissions trading scheme as the most reliable data for these two sources. The agency also plans to continue the practice of outsourcing contracts on a periodic basis to re-examine and extend the inventory time-series for F-gases. This approach has been found to be an efficient way of compiling the estimates for sources and gases that the inventory experts in the agency have not worked on in detail in the past, as shown by the complete time-series of HFC, PFC and SF₆ emissions included in the 2007 submission.

Table 4.3. Emissions of HFC, PFC and SF₆ from Industrial Processes 1990-2005 (Gg CO₂ eq)

IPCC Source Category	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
2.F.1 Refrigeration and Air-Conditioning	NO	3.322	3.512	3.690	3.984	4.481	7.145	20.056	40.616	47.660	51.213	47.527	62.879	110.613	133.089	150.972
2.F.1 Mobile Air Conditioning	NO	NO	NO	1.222	5.068	10.141	16.357	25.077	38.229	54.119	70.461	83.364	96.376	110.148	126.120	145.631
2.F.2 Foams	NO	0.000	0.016	0.063	0.303	0.644	1.107	1.779	3.619	5.222	6.275	9.211	11.499	13.705	15.865	18.038
2.F.3 Fire-extinguishers	0.219	0.700	1.179	1.677	2.235	2.839	3.496	4.210	4.988	5.834	6.755	7.806	9.026	10.385	11.915	13.636
2.F.4 Aerosols	0.006	0.648	0.721	1.784	7.109	24.959	45.142	76.202	97.501	72.336	78.879	86.369	77.891	83.810	77.286	79.271
2.F.4 Metered Dose Inhalers	0.000	0.000	0.000	0.000	0.000	0.000	6.100	6.700	7.500	8.300	9.800	14.126	16.992	18.211	18.409	21.738
2.F.7 Semiconductor manufacture	0.468	0.468	0.468	0.468	1.825	1.825	2.937	4.633	3.877	9.418	12.379	3.101	1.884	2.644	1.357	1.739
TOTAL HFC	0.693	5.138	5.897	8.903	20.525	44.889	82.284	138.657	196.330	202.889	2358.762	251.504	276.547	349.517	384.040	431.026
2.F.7 Semiconductor manufacture	0.093	0.093	0.093	0.093	75.382	75.382	103.085	130.823	61.870	195.933	305.406	295.984	212.403	228.795	186.840	173.954
TOTAL PFC	0.093	0.093	0.093	0.093	75.382	75.382	103.085	130.823	61.870	195.933	305.406	295.984	212.403	228.795	186.840	173.954
2.F.7 Semiconductor manufacture	0.478	0.478	0.478	0.478	43.020	43.020	62.140	81.260	52.580	16.730	31.070	20.435	28.584	59.917	35.635	65.554
2.F.8 Electrical equipment	21.510	22.466	23.422	24.378	25.334	26.290	26.386	37.284	25.238	34.990	7.787	32.050	22.786	38.446	21.553	23.518
2.F.9 Other - window soundproofing	0.431	0.451	0.472	0.492	0.512	0.532	0.551	0.570	0.590	0.465	0.333	0.195	0.193	0.191	0.189	0.187
2.F.9 Other - medical applications	0.797	0.797	0.797	0.797	0.797	0.797	0.797	0.797	0.797	0.797	0.797	0.797	0.797	0.797	0.797	0.797
2.F.9 Other - sporting goods	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	2.851	3.786	3.731	3.821	5.759	7.142	5.813	5.903
2.F.9 Other - gas-air tracers	12.189	12.189	12.189	12.189	12.189	12.189	12.189	12.189	12.189	12.189	12.189	12.189	12.189	12.189	6.095	0.000
TOTAL SF₆	35.405	36.381	37.357	38.334	81.852	82.827	102.062	132.100	94.245	68.956	55.907	69.486	70.308	118.682	67.093	95.958
TOTAL HFC, PFC and SF₆	36.191	41.612	43.347	47.330	177.758	203.098	281.373	395.011	345.125	459.713	590.259	616.974	559.258	696.993	637.974	700.941

Table 4.4. Recalculated Emission Estimates for Industrial Processes 1990-2004 (Gg CO₂ eq)

IPCC Source Category	Gas	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
Previously Reported Values																
2.A.1 Cement Production	CO ₂	884.000	782.000	753.000	729.000	859.000	879.000	983.000	1145.000	1059.000	1166.000	1700.900	1851.200	1859.800	2128.000	2290.000
2.F.1 Mobile Air Conditioning	HFC	NO	NO	NO	1.222	5.068	10.141	16.357	25.077	38.229	54.119	70.461	83.364	96.376	110.148	124.290
2.F.2 Foams	HFC	NO	0.000	0.016	0.063	0.303	0.644	1.107	1.779	3.619	5.222	6.275	9.211	11.499	13.705	15.812
2.F.4 Aerosols	HFC	0.006	0.643	0.715	1.767	7.036	24.668	38.499	68.539	90.186	64.279	72.045	87.955	90.163	92.205	94.257
2.F.7 Semiconductor manufacture	HFC	0.468	0.468	0.468	0.468	1.825	1.825	2.937	4.633	3.877	9.418	12.379	3.101	1.884	2.644	1.498
2.F.7 Semiconductor manufacture	PFC	0.093	0.093	0.093	0.093	75.382	75.382	103.085	130.823	61.870	195.933	305.406	295.984	212.403	228.795	196.370
2.F.9 Other - sporting goods	SF ₆	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	2.851	3.786	3.731	3.765	5.675	7.051	5.740
Recalculated Values																
2.A.1 Cement Production	CO ₂	884.000	782.000	753.000	729.000	859.000	879.000	983.000	1145.000	1059.000	1166.000	1700.900	1851.200	1859.800	2127.000	2295.100
2.F.1 Mobile Air Conditioning	HFC	NO	NO	NO	1.222	5.068	10.141	16.357	25.077	38.229	54.119	70.461	83.364	96.376	110.148	126.120
2.F.2 Foams	HFC	NO	0.000	0.016	0.063	0.303	0.644	1.107	1.779	3.619	5.222	6.275	9.211	11.499	13.705	15.865
2.F.4 Aerosols	HFC	0.006	0.648	0.721	1.784	7.109	24.959	45.412	76.202	97.501	72.336	78.879	86.369	77.891	83.810	77.286
2.F.7 Semiconductor manufacture	HFC	0.468	0.468	0.468	0.468	1.825	1.825	2.937	4.633	3.877	9.418	12.379	3.101	1.884	2.644	1.357
2.F.7 Semiconductor manufacture	PFC	0.093	0.093	0.093	0.093	75.382	75.382	103.085	130.823	61.870	195.933	305.406	295.984	212.403	228.795	186.840
2.F.9 Other - sporting goods	SF ₆	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	2.851	3.786	3.731	3.821	5.759	7.142	5.813
% Change in Emission																
2.A.1 Cement Production	CO ₂	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	-0.05	0.22
2.F.1 Mobile Air Conditioning	HFC	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	1.47
2.F.2 Foams	HFC	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	0.34
2.F.4 Aerosols	HFC	NO	0.78	0.84	0.96	1.04	1.18	17.96	11.18	8.11	12.53	9.49	-1.80	-13.61	-9.10	-18.01
2.F.7 Semiconductor manufacture	HFC	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	-9.41
2.F.7 Semiconductor manufacture	PFC	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	-4.85
2.F.9 Other - sporting goods	SF ₆	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	1.49	1.48	1.29	1.27

Chapter Five

Solvent and Other Product Use

5.1 Overview of *Solvent and Other Product Use* Sector

This IPCC source category is considered separately because of its importance in relation to the emissions of NMVOC (non-methane volatile organic compounds), one of the indirect greenhouse gases, which result from the use of solvents and various other volatile compounds. However, some minor direct uses of N₂O (such as anaesthesia) are covered in this source category and the IPCC reporting format also explicitly provides for the inclusion of CO₂ emissions that result from the oxidation of the carbon in VOC emissions. This is consistent with the overall approach adopted for estimating CO₂ from the combustion of fuels using the sectoral approach (Section 3.2), where the CO₂ emissions are based on the full carbon content of the fuel even though some of the carbon is usually emitted as NMVOC or CO. The Irish inventories include an estimate of CO₂ emissions in this way but emissions associated with the direct use of N₂O are not estimated.

The activity data used for computing estimates of CO₂ emissions in *Solvent and Other Product Use* are the mass emissions of NMVOC computed for the relevant source categories (3.A *Paint Application*, 3.B *Degreasing and Dry Cleaning*, 3.C *Chemical Products* and 3.D *Other Solvent Uses*). The Irish data used for this purpose are the VOC emissions compiled according to the CORINAIR methodology for reporting to UNECE under the Convention on Long Range Transboundary Air Pollution (CLRTAP). As part of the work on recalculations for the 2002 submission, Ireland produced a revised and consistent time-series of such NMVOC emissions estimates based on the results of detailed analysis and investigations for 1998 (Finn et al, 2001). The CO₂ emissions were derived by assuming that 85 percent of the mass emissions of NMVOC in the four categories is converted to CO₂.

5.2 NMVOC and CO₂ Inventory Time Series

Emission control strategies are being implemented in Ireland to comply with a limit of 65 kt for total emissions of NMVOC in 2010 under the National Emissions Ceilings Directive. The levels of solvent use and the emissions from solvents are changing substantially in response to product replacement and reformulation and emission controls being implemented under Integrated Pollution Control (IPC) and the Solvents Directive. In these circumstances, the inventories of VOC emissions from solvent use over recent years were reassessed as part of the general improvements conducted for Irish emission inventories during 2005. The inventory agency commissioned a project to carry out in-depth analysis of the specified NMVOC source categories (CTC, 2005) in order to compile the best possible estimates of emissions in 2004 as a follow-up to the earlier commissioned work and to revise the inventories for the years 1998-2003 as necessary in the light of new information. The revised estimates for these target years indicated lower NMVOC emissions than had been previously reported and used as the basis for CO₂ in the sector *Solvent and Other Product Use*.

A bottom-up approach was possible for activities subject to IPC licensing in the four source categories. Relevant data on emissions and solvent use were extracted from their electronic or paper Annual Environmental Reports (AERs) or Pollution Emissions Registers (PERs). Where such information was not available, European PERs were assessed. Top-down methods were used for activities not covered by the IPC licensing system. These included the use of paints and the use of domestic solvents, the two principal source categories. Input, usage and emissions data for each individual activity was collated into IPC and non-IPC spreadsheets and emissions were estimated by applying EMEP/CORINAIR methods, default emission factors and general guidance as appropriate. Scaling up to national level was applied where necessary.

The estimates of CO₂ emissions from Solvent and Other Product Use for 1990-2005 are presented in Table 5.1. The largest contributor to overall emissions is domestic solvent use. It is also to be noted that emissions from this sector have increased while those from the majority of sectors are decreasing. The main drivers are considered to be the increased number of vehicles, growth in the number of individual households, and higher per-capita consumption of non-aerosol automotive products, cosmetics, toiletries, and household products. It should be noted that UK emission factors together with Irish statistics for number of vehicles, persons and households were used in the absence of any other data. One of the only two other significant sectors for which emissions are increasing is industrial application of paint in the wood products sector. This is as a result of an expansion in activity in the sector as well as the continued use of conventional high solvent content coatings. The vast majority of these companies are small operations outside the remit of IPC.

Emissions from architectural paint use are decreasing (even while paint sales are increasing) as a result of an increased market share for water-based paints and a reduction in the VOC content of water based paints (VOC content of solvent based paints remains more or less static). From discussions with industry, one of the key drivers for the decrease in solvent use in architectural paint has been as a result of pressure from some of the larger retailers. The decrease in VOC emissions from architectural painting should be set to continue with the advent of the deco-paints Directive (2004/42/EC) and can only benefit from continued and expanded retailer/consumer pressure. There have been significant drops in both printing and wood impregnation. The decrease in printing is principally due to the installation of abatement equipment in the plant, which is the largest user of solvents. The decrease in the use of wood preservatives can be attributed to several site closures and to the switch from solvent-borne to water-borne wood preservatives.

Other industrial paint application and other manufacturing taken together show a decrease in emissions between 1998 and 2005. The diversity within these sectors is very large in terms of the type of process, the products made, and the scale involved. There have been closures, particularly of a few of the large emitters, which have decreased emissions, but there has also been some new processes licensed. In addition there is a large degree of uncertainty associated with the non-IPC element of the emissions estimates for these sources. However, the study found that there are specific instances of IPC licensed sites reducing VOC emissions through prevention at source or through abatement.

5.3 Recalculations for Solvents and Other Product Use

In the compilation of emission estimates for 2005, it was found that underlying UK population statistics used as scaling factors for the period 1990 to 2004 inclusive underlying the calculation of emissions from sectors 3A (Paint Application) and 3D (Other Use of Solvents) were not consistent within the inventory and with those use in other emission inventories. This anomaly has been rectified. A comparison of these recalculated totals with those in the 2006 submission is presented in Table 5.2.

Table 5.1 Estimates of NMVOC and CO₂ Emissions from Solvent and Other Product Use 1990-2005

	3A Paint Application	3B Degreasing, dry cleaning, electronics	3C Chemical Products Manufacturing & Processing	3D Other Use of Solvents	Total NMVOC Emissions	Estimated CO ₂ emissions from NMVOC
2007 Submission	Mg NMVOC					Gg
1990	6,878	2,304	3,534	13,228	25,944	80.86
1991	7,036	2,304	3,534	13,674	26,548	82.74
1992	7,231	2,304	3,534	13,550	26,619	82.96
1993	7,485	2,304	3,534	13,444	26,767	83.42
1994	7,782	2,291	3,534	13,446	27,053	84.32
1995	8,082	2,496	3,534	13,517	27,629	86.11
1996	8,406	2,423	3,534	13,485	27,848	86.79
1997	8,687	2,311	3,534	13,447	27,979	87.20
1998	9,053	2,158	3,534	13,508	28,253	88.06
1999	8,761	2,043	3,442	12,947	27,193	84.75
2000	8,071	1,883	3,350	12,386	25,690	80.07
2001	8,375	1,660	3,168	13,356	25,560	79.66
2002	8,649	1,067	2,986	12,123	24,826	77.37
2003	8,625	1,141	2,804	11,801	24,371	75.96
2004	8,700	1,155	2,622	11,811	24,288	75.70
2005	8,785	1,079	2,440	11,901	24,206	75.44

Table 5.2 Recalculations for Solvent and Other Product Use 1990-2004

	3A Paint Application	3B Degreasing, dry cleaning, electronics	3C Chemical Products Manufacturing & Processing	3D Other Use of Solvents	Total NMVOC Emissions	Estimated CO ₂ from NMVOC
	Mg NMVOC					Gg
2006 Submission						
1990	6878	2304	3534	13253	25969	80.86
1991	7036	2304	3534	13699	26573	82.74
1992	7231	2304	3534	13576	26645	82.96
1993	7485	2304	3534	13470	26792	83.42
1994	7782	2291	3534	13472	27079	84.32
1995	8082	2496	3534	13543	27655	86.11
1996	8406	2423	3534	13511	27874	86.79
1997	8687	2311	3534	13474	28005	87.20
1998	9060	2158	3534	13588	28340	88.06
1999	8767	2043	3433	13032	27275	84.75
2000	8079	1883	3332	12476	25770	80.07
2001	8375	1660	3132	12355	25523	79.66
2002	8649	1067	2932	12121	24770	77.37
2003	8625	1141	2732	11794	24293	75.96
2004	8700	1155	2532	11508	23895	75.70
2007 Submission						
1990	6878	2304	3534	13228	25944	80.94
1991	7036	2304	3534	13674	26548	82.82
1992	7231	2304	3534	13550	26619	83.04
1993	7485	2304	3534	13444	26767	83.50
1994	7782	2291	3534	13446	27053	84.40
1995	8082	2496	3534	13517	27629	86.19
1996	8406	2423	3534	13485	27848	86.87
1997	8687	2311	3534	13447	27979	87.28
1998	9053	2158	3534	13508	28253	88.33
1999	8761	2043	3442	12947	27193	85.04
2000	8071	1883	3350	12386	25690	80.37
2001	8375	1660	3168	12356	25560	79.66
2002	8649	1067	2986	12123	24826	77.37
2003	8625	1141	2804	11801	24371	75.97
2004	8700	1155	2622	11811	24288	75.70
% Change in Emission						
1990	NO	NO	NO	-0.19	-0.10	0.10
1991	NO	NO	NO	-0.18	-0.09	0.10
1992	NO	NO	NO	-0.19	-0.10	0.10
1993	NO	NO	NO	-0.19	-0.09	0.10
1994	NO	NO	NO	-0.19	-0.10	0.10
1995	NO	NO	NO	-0.19	-0.09	0.09
1996	NO	NO	NO	-0.19	-0.09	0.09
1997	NO	NO	NO	-0.20	-0.09	0.09
1998	-0.08	NO	NO	-0.59	-0.31	0.31
1999	-0.07	NO	0.26	-0.65	-0.30	0.34
2000	-0.10	NO	0.54	-0.72	-0.31	0.38
2001	NO	NO	1.15	0.01	0.14	0.00
2002	NO	NO	1.84	0.02	0.23	-0.01
2003	NO	NO	2.64	0.06	0.32	0.02
2004	NO	NO	3.55	2.63	1.64	0.00

Chapter Six

Agriculture

6.1 Overview of Agriculture Sector

Table 6.1 lists the IPCC Level 3 source categories in *Agriculture*, where CH₄ and N₂O are the key greenhouse gases. The agricultural activities of particular importance in Ireland are those under *4.A Enteric Fermentation*, *4.B Manure Management* and *4.D Agricultural Soils* only, some of which are identified as being among the largest greenhouse gas emission sources in the country (Chapter Three). The inventory time-series for the years 1990-2005 contains emission estimates for all relevant sources and gases in these three important source categories. The availability of better up-to-date data and the completion of major national research in agriculture has facilitated major improvements in methodologies and in the manner of data application for many of the sources concerned. Source categories *4.C Rice Cultivation*, *4.E Prescribed Burning of Savannas* and *4.F Field Burning of Agricultural Residues* are not relevant to Ireland and the notation key NO is used in relation to all associated emissions in the CRF. Although the practice of field burning of agricultural residues did exist on a small scale in the past, the emissions are considered negligible, and it has been discontinued since the mid 1990s.

The methods provided by the IPCC good practice guidance are now being applied as completely as possible for agricultural emission sources under Irish circumstances. The IPCC methods require considerable information detail on activity data, emission factors and other input parameters needed for the emission calculations. There were major changes in the inventories for *Agriculture* in the 2006 submission with the adoption of Tier 2 methods for CH₄ emissions from enteric fermentation in cattle and robust improvement in estimates of emissions from manure management based on the results of major research and an extensive farm facilities survey conducted in recent years. This research, together with other relevant work related to the development of an elaborate new NH₃ inventory for agriculture in 2005 and guidelines on implementation of the EU Nitrates Directive (CEC, 1991) has facilitated the application of a large amount of country-specific information underlying the various estimates of emissions. The same method is used for the purposes of this submission but further development and minor updating of the underlying activity data remains part of the ongoing work and assessment in relation to agricultural emissions.

Because of the importance of agriculture in the country, Ireland has very extensive and up-to-date statistical data on all aspects of the sector, compiled and published by the Central Statistics Office. This is the official source of the basic data for inventory purposes, except for synthetic fertilizer use and poultry population statistics, for which annual data are obtained from the Department of Agriculture and Food. The time-series of key agricultural statistics as used for the various activity data (livestock populations and fertilizer use) is given in Annex D. It may be noted that in the case of cattle, the populations related to housing (Table D.1(a)) are different to those for pasture (Table D.1(b)) to take full account of the respective production systems. The manner in which the populations are applied is explained in the documentation boxes of the CRF tables.

Table 6.1. Level 3 Source Category Coverage for Agriculture

Agriculture	CO ₂	CH ₄	N ₂ O
A. Enteric Fermentation			
1. Cattle	NA	All	NA
Dairy Cattle	NA	All	NA
Non-Dairy Cattle	NA	All	NA
2. Buffalo	NA	NO	NA
3. Sheep	NA	All	NA
4. Goats	NA	All	NA
5. Camels and Llamas	NA	NO	NA
6. Horses	NA	All	NA
7. Mules and Asses	NA	All	NA
8. Swine	NA	All	NA
9. Poultry	NA	NE	NA
10. Other	NA	NO	NA
B. Manure Management			
1. Cattle	NA	All	All
Dairy Cattle	NA	All	All
Non-Dairy Cattle	NA	All	All
2. Buffalo	NA	NO	NO
3. Sheep	NA	All	All
4. Goats	NA	All	All
5. Camels and Llamas	NA	NO	NO
6. Horses	NA	All	All
7. Mules and Asses	NA	All	All
8. Swine	NA	All	All
9. Poultry	NA	All	All
10. Anaerobic Lagoons	NA	NA	NA
11. Liquid Systems	NA	All	All
12. Solid Storage and Dry Lot	NA	All	All
13. Other	NA	NO	NO
C. Rice Cultivation	NO	NO	NO
D. Agricultural Soils	NO	NO	NO
1. Direct Soil Emissions	IE*	NE	All
2. Pasture Range and Paddock Manure	NA	NO	All
3. Indirect Emissions	NA	NO	All
4. Other	NO	NO	NO
E. Prescribed Burning of Savannas	NO	NO	NO
F. Field Burning of Agricultural Residues	NO	NO	NO
G. Other	NO	NO	NO

All : all emission sources covered; NE : emissions not estimated; NO : activity not occurring; NA : not applicable (activity exists but no emissions of the gas occurs); IE : emissions included elsewhere

* CO₂ emissions from Liming of Agricultural Lands included in category 5.B of LULUCF (Chapter Seven)

6.2 CH₄ Emissions from Enteric Fermentation

6.2.1 Overall Approach

Ireland used Tier 1 methods to estimate CH₄ emissions from enteric fermentation for all inventory submissions up to 2006. This basic approach showed that enteric fermentation in dairy and non-dairy cattle produced 8.5 percent and 3.6 percent respectively, of total GHG emissions in 2003. In 1990, enteric fermentation in non-dairy cattle was the single largest individual source of emissions in Ireland, accounting for 9.6 percent of the national total. Implementation of the IPCC good practice guidance for GHG inventories requires that Parties use Tier 2 (detailed country-specific) methods for key sources of greenhouse gases, as in the case of CH₄ from enteric fermentation from cattle in Ireland. The recommendation to use Tier 2 methods has also been made in several reports on the review of Ireland's inventory submissions to UNFCCC. This recommendation was first implemented in the 2006 submission.

The Irish cattle herd is now characterised by 11 principal animal categories as given in Table 6.1 for which annual census data are published by CSO. In-depth analysis of production

systems and the associated animal feed and energy requirements was conducted for all categories to determine CH₄ production. Substantial further subdivision was incorporated for dairy and beef cattle to adequately describe the wide range of cattle rearing and finishing systems applicable in Ireland. Dairy cows were covered by 12 systems and 18 were analysed for suckler cows, while up to 30 systems were used for both male and female beef cattle. The exercise to develop Tier 2 emission factors for the 11 animal categories was initially carried out for the 2003 herd and then repeated for 1990 and 1994. The following paragraphs outline the approach and a detailed description of the comprehensive study and analysis underlying the new emission factors is available (O'Mara, 2006).

Table 6.2 Animal Classification for Cattle Population

Cattle Type	Classification		
Dairy cattle	Dairy cows	Suckler (Beef) cows	
Beef cattle	Male < 1 year Female < 1 year	Male 1 – 2 years Female 1 – 2 years	Male > 2 years Female > 2 years
Other cattle	Breeding bulls	Dairy in-calf heifers	Beef in-calf heifers

6.2.2 Enteric Fermentation in Dairy Cattle

For both dairy cows and suckler cows, the country is divided into three regions: (1) south and east, (2) west and midlands, and (3) north west, coinciding with regions used for implementing the Nitrates Regulations (SI No 788 of 2005) based on slurry storage requirements of local planning authorities. This facilitates in-depth analysis for separate regions with different lengths of winter housing and takes account of different animal feeding practices. The number of cows in each category given by CSO statistics is allocated to the regions identified above using the Cattle Movement Monitoring System (CMMS) reports published by the Department of Agriculture and Food (DAF, 2006). The cattle production system in each region was defined in terms of calving date, dates of winter housing and spring turn-out to grass, milk yield and composition, forage and concentrate feeding level, cow live-weight and live-weight change and lactation period.

In the approach outlined by O'Mara (2006), the daily energy requirement of cows in each region was calculated by month or part thereof based on maintenance requirements, milk yield and composition, requirements for foetal growth, and gain or loss of bodyweight (INRA, 1989). In this system, net energy requirements are defined in terms of units of feed for lactation (UFL), where 1 UFL is the net energy value of 1 kg of barley at 86 percent dry matter and is equal to 7.11 MJ net energy for lactation (NE_l). This international energy system, well established and used locally in Ireland, was considered more appropriate to the local conditions than the system and equations used by the IPCC guidelines and IPCC good practice guidance.

The important equations are:

Maintenance NE_l requirements (MJ) = 9.96 + (0.6 × LW/100), where LW is liveweight.

A 10 percent activity allowance was added for the housed period and a 20 percent allowance was added for the grazing period as outlined by INRA (1989);

NE_l (MJ) required per kg milk = 0.376 × fat content + 0.209 × protein content + 0.948;

Pregnancy: mean of 12.1 MJ NE_l /day for the last 3 months of pregnancy;

Liveweight change: each kg liveweight lost contributed 24.9 MJ NE_i to energy requirements, while each kg of liveweight gained required 32 MJ NE_i.

The composition of the diet of cows in each region was described by month or part thereof and daily intake was calculated by reference to the daily energy requirement. The concentrate allowance was fixed while forage intake varied according to energy requirements. Daily methane emissions (MJ/day) were calculated from digestible energy intake using the equation of Yan et al. (2000).

$$\text{CH}_4 = \text{DEI} * [0.096 + (0.035 \times \text{S}_{\text{DMI}}/\text{T}_{\text{DMI}})] - 2.298 * (\text{FL} - 1)$$

where DEI is digestible energy intake (MJ/day), S_{DMI} and T_{DMI} are silage and total dry matter intakes (kg/day), respectively, and FL is feeding level (multiples of the maintenance energy requirement).

A constant methane conversion rate of 0.065 of gross energy intake was applied when the diet consisted of grazed grass and 3 kg or less of concentrate supplement per day. This was based on a large New Zealand database of measurements for grazing animals on similar production systems to those in Ireland. A methane output of 21.6 g/kg DM is used for pasture diets with a grass GE content of 18.45 MJ/kg, which is equivalent to 6.5 percent of GE. (Harry Clark, Personal Communication). The daily CH₄ emissions were summed to give annual emissions for cows in each region, and a weighted national average was then calculated.

6.2.3 Enteric Fermentation in Beef Cattle

Emission factors for the beef cattle categories given in Table 6.3 were determined by calculating lifetime emissions for the animal and by partitioning between the first, second and third years of the animal's life. This approach allows the published CSO animal populations for June to be used directly as the activity data most representative of the inventory year for enteric fermentation while taking into account the movement of cattle from one category to another, as enumerated by the June census, up to two times in their lifetime.

Analysis was undertaken for a total of 11 separate production systems covering the three groups of male and female beef cattle given in Table 6.3 after the proportion of the herd in each category was calculated. Important parameters such as housing dates (expert opinion), turnout dates (expert opinion) and live-weight gains (expert opinion reconciled with actual national carcass weights) during winter housing periods and grazing seasons were defined for each system. The most important parameter is liveweight gain as it directly affects the energy requirement and thus feed intake. There is little statistical information on the liveweight gain of the different types of cattle in the Irish cattle herd, but the weight of carcasses of all slaughtered cattle is recorded by the Department of Agriculture and Food. Using data for the average carcass weight of male and female cattle, appropriate liveweight gains were applied to the various life stages of each animal category, such that when all categories were combined, that data were consistent with the national statistics for carcass weight (plus or minus 10 kg difference).

Given these data for liveweight and liveweight gain, energy requirements of animals were estimated during the winter housing periods and grazing seasons of the animals lifetime using the INRAtion computer programme, version 3.0. This programme is devised by the French research organisation INRA, and is based on the net energy system for cattle. In version 3 of INRAtion, some adaptation for Irish conditions was made to the equations for estimating the energy requirements of growing and finishing animals (O'Mara, 1997, Crowley, 2001 and Crowley *et al*, 2002). Net energy requirements of growing beef cattle are

defined in terms of UFL, as in the case of dairy cattle, while for finishing cattle, net energy requirements are defined in terms of UFV (from the French *unite forrige viande*) and 1 UFV is equal to 7.61 MJ NE_{mg}.

The composition of the diet in each system was described by grazing season and winter housing period and daily intake was calculated by reference to the daily energy requirement. The concentrate allowance was fixed while forage intake varied according to energy requirements. The Irish modifications to the INRA programme were predominantly for animals at weanling and finishing stages (i.e. at times that concentrates were likely to be fed). No modifications were made for 'heavy' growing animals, (typically animals in their second grazing season or later that were not being finished). For animals in these stages, intakes were adjusted as appropriate by expert opinion. Daily methane emissions were calculated using the equation of Yan et al. (2000), however a constant of 0.065 of gross energy intake was applied when the diet was grazed grass plus 3 kg or less of concentrate supplement/day. The daily emissions were aggregated to give annual emissions per system and a weighted national average was then calculated.

6.2.4 Enteric Fermentation in Other Cattle

Bulls for breeding and in-calf heifers account for approximately 6 percent of the national cattle herd. Separate production systems are not defined for these categories because of lack of published data on their feed intake and the small number of animals involved. Bulls for breeding are mostly of continental breeds, and their emission factors are based on those for late maturing male beef cattle of suckler origin in their second year. The value of 66.5 kg/year for animals in this category is determined by an applicable period of 310 days in their second year, which is adjusted to 81.5 kg/year using the full period of 365 days for breeding bulls.

In-calf heifers were assigned the same emission factors as female beef cattle (50.16 and 53.58 kg/year for dairy and beef animals, respectively) in their second year (i.e. corresponding to category 1–2 years old). In-calf heifers only need emissions associated with the period March – December of their second year to be accounted for, as they are subsequently enumerated as dairy or suckler cows in the CSO animal census thereafter. Female beef cattle in the category 1-2 years old are assumed to be slaughtered on 3rd February of their third year. Adjustment for the slightly longer period was not made in respect of in-calf heifers, as they are carrying a calf in addition to normal growth.

6.2.5 Summary of Tier 2 Emission Factors for Cattle

The Tier 2 emission factors developed by the detailed analysis outlined above for the years 2005, 2004 and 1990 are summarised in Table 6.3 for the 11 principal categories chosen to characterise the Irish cattle herd. Emission factors for the full time series 1990–2005 in respect of the 11 principal categories are presented in Table D.2 of Annex D. The emission factor for dairy cows in 1990 is very close to the IPCC default emission factor of 100 kg CH₄/head/year for highly productive dairy cattle in Western Europe, which Ireland has used to date. The corresponding value for 2004 indicates an increase of about 8 percent from 1990 in line with increased milk yield, which is not captured by the Tier 1 approach previously used. As such, annual milk yield may be used as a convenient basis for deriving aggregate weighted emission factors for dairy cattle in other years. The emission factors for beef cattle indicate an overall weighted average of approximately 40 kg/head, compared to the value of 50 kg/head previously used. Little change is indicated between 1990 and 2004, except in the case of male cattle in the category of animals greater than two years old. This is explained by the earlier finishing time for male beef cattle since the BSE crisis that affected agriculture during the 1990s.

Table 6.3 Tier 2 CH₄ Emission Factors for 1990, 2004 and 2005

	Enteric Fermentation (kg/head/year)			Manure Management (kg/head/year)		
	2005	2004	1990	2005	2004	1990
Dairy cows	107.22	108.50	101.38	20.43	20.53	21.57
Suckler cows	74.12	74.41	74.03	13.86	13.90	14.02
Male cattle < 1 year	29.72	29.70	30.46	8.60	8.58	9.73
Male cattle 1 - 2 years	58.96	59.26	62.22	13.82	13.75	16.68
Male cattle > 2 years	37.67	35.23	55.08	1.91	1.60	4.57
Female cattle < 1 year	27.86	27.89	27.05	8.29	8.30	8.79
Female cattle 1 - 2 years	45.64	44.50	53.54	9.63	9.11	14.74
Female cattle > 2 years	22.46	22.46	21.65	0.34	0.34	0.33
Bulls for breeding	80.87	81.21	86.38	18.21	18.58	23.79
Dairy in-calf heifers	49.90	50.03	51.82	10.55	10.74	13.40
Beef in-calf heifers	53.30	53.44	55.42	12.45	12.66	15.61

6.2.6 Enteric Fermentation in Other Livestock

The type of information used to derive the Tier 2 emission factors for cattle is not available for other important livestock categories, such as sheep and swine. Therefore, the inventory agency continues to use the Tier 1 approach for enteric fermentation for all livestock categories other than cattle. The emission factors used are generally those for Western Europe given in Table 4.3 of the IPCC Guidelines. However, in order to fully utilize the detailed CSO breakdown in respect of sheep and swine populations, the base emission factors from IPCC are adjusted in each case on the basis of animal weight, as shown in Table D.2 of Annex D. As a result, the implied emission factors produced by the CRF related to sheep and swine are lower than the default values.

6.3 CH₄ Emissions from Manure Management

The decomposition of the organic material in animal manures may be a significant source of CH₄ emissions if anaerobic conditions prevail in the animal waste management systems being used. The estimation of such emissions requires information on the quantity of manure production for the animal groups concerned, the type of waste management systems employed and the CH₄ production potential of the wastes. New information obtained from farm facilities surveys that were undertaken as part of the research underlying an elaborate new NH₃ inventory for Ireland and the work on emission factors for enteric fermentation in cattle described above is the basis for revised CH₄ emission factors for manure management. The results of the farm facilities surveys provide a much improved representation of animal waste allocation among the relevant waste management systems in the country while the excretion of organic matter by cattle is fully characterised as part of the analysis of their feed and energy requirements relating to enteric fermentation. The NIRs prior to the 2006 submission have stated that sheep remain outdoors all year round and that there is no management of sheep manures in Ireland. The farm facilities surveys show that lowland sheep are housed for some time during the year thus allowing for the inclusion of

sheep manures in the estimation of emissions from manure management. The manures from horses, mules and asses are also included on the assumption that they are distributed equally between solid storage and pasture systems.

The analysis of the feeding regime for cattle included a full evaluation of the organic matter content of the feeds applicable to the 11 categories that characterise the national herd, which facilitated the estimation of their respective levels of organic matter excretion. The emission factors for manure management were derived using the quantified organic matter excretion as volatile solids (VS), a BO (the methane production potential of animal waste) value of 0.24 m³ CH₄/kg VS, the allocation to animal waste management system based on the farm facilities survey and the corresponding values of MCF (methane conversion factor) given for the cool climate zone in Table 4.10 of the IPCC good practice guidance. The emission factors for cattle are given in Table 6.3. They are higher than those used prior to the 2006 submission mainly because a much higher proportion of waste is allocated to liquid systems for which the applicable updated MCF value is 0.39.

The calculation of CH₄ emissions from domestic livestock includes the derivation of the emission factors for manure management for sheep, swine, horses and poultry. The allocations to animal waste management system are again based on the farm facilities survey and appropriate values of BO and VS are taken from the IPCC Guidelines while MCF is again as given in Table 4.10 of the IPCC good practice guidance. The emission factors for swine are substantially higher than previously used, as all wastes are allocated to liquid systems, which have a relatively high MCF of 0.39. The application of the manure management emission factors for sheep, horses and poultry means that all CH₄ emissions from livestock are included in current estimates. The CH₄ emissions from manure management in 2005 amounted to 24.6 percent of those from enteric fermentation.

6.4 N₂O Emissions from Manure Management

Nitrogen excretion rates have been adopted in Ireland for all animal categories for which annual census data are published by the CSO. These rates of nitrogen excretion are endorsed by the Department of Agriculture and Food and by TEAGASC for national use and guidance for farmers in relation to implementation of the Nitrates Regulations. In the case of cattle, the excretion rates are consistent with the nitrogen content of cattle feeds and the quantities excreted by the animal, as analysed in conjunction with the determination of Tier 2 CH₄ emission factors for cattle. The published nitrogen excretion rates are used by the inventory agency, along with the information on the allocation of animal manures to each applicable animal waste management system as the basis of CRF Table 4.B(b).

Approximately two-thirds of animal waste nitrogen is excreted at pasture annually, reflecting the relatively short period that cattle are housed in Ireland and a significant contribution from the large sheep population. Animal wastes excreted at pasture are unmanaged and the associated emissions are accounted for under agricultural soils. The bulk of animal wastes in housing are managed in liquid storage systems (93 percent and 70 percent for dairy cattle and other cattle, respectively and 100 percent for swine) for eventual spreading on agricultural lands. The remainder of animal wastes produced in-house is treated in solid systems. The emission factors given by the IPCC good practice guidance indicate that 1 kg of nitrogen per tonne of nitrogen handled in liquid systems is lost as N₂O while the corresponding loss is 20 kg per tonne for nitrogen in solid storage systems. These default emission factors, for which uncertainty ranges of up to 100 percent are assigned in the IPCC good practice guidance, are used to estimate N₂O emissions from manure management in Ireland. The N₂O emissions from manures managed in liquid and solid storage systems in 2005 amounted to 1.32 Gg.

6.5 N₂O Emissions from Agricultural Soils

Agricultural soils are the principal source of N₂O emissions in many countries. The IPCC methodologies for the source categories concerned are essentially an accounting of all inputs of nitrogen to agricultural soils and the subsequent application of default rates of nitrogen for losses to the atmosphere as N₂O. The primary nitrogen inputs are subject to complex processes and partitioning between various nitrogen compounds within soils and the emissions are highly dependent on soil properties and meteorology. The methodologies are therefore simplified and they are based on a consideration of separate direct and indirect contributions to national emissions. Ireland uses the IPCC good practice guidance methodology completely to estimate N₂O emissions from agricultural soils and the procedure may be followed from the description below. Values for each of the terms used in the calculation of direct and indirect soil emissions for the full time series 1990-2005 are presented in Table D.3 of Annex D.

6.5.1 4.D.1 Direct Soil Emissions

According to the IPCC good practice guidance the direct emissions of N₂O to be reported in CRF sub-category 4.D.1 Direct Soil Emissions may be calculated in a Tier 1 approach from

$$N_2O_{\text{direct}} = (F_{\text{SN}} + F_{\text{AM}} + F_{\text{S}} + F_{\text{BN}} + F_{\text{CR}}) * EF_1 + F_{\text{OS}} * EF_2$$

where

N_2O_{direct} = the direct emissions of N₂O

F_{SN} = amount of synthetic fertilizer nitrogen applied to soils, adjusted for the amount that volatilizes as NH₃ and NO_x

F_{AM} = amount of animal manure nitrogen applied directly to soils, adjusted for the amount that volatilizes as NH₃ and NO_x

F_{S} = amount of organic nitrogen in sludges applied to agricultural soils

F_{BN} = amount of nitrogen fixed by nitrogen-fixing crops

F_{CR} = amount of nitrogen in crop residues returned to soils

F_{OS} = the area of cultivation of organic soils

EF_1 = N₂O emission factor for emissions from direct nitrogen inputs (kg N₂O-N/kg N)

EF_2 = N₂O emission factor for emissions from cultivation of organic soils (kg N₂O-N/kg N)

The estimates of direct N₂O emissions from agricultural soils for the years 1990-2005 take into account the nitrogen inputs from all these sources, except that due to the cultivation of organic soils. Tillage farming in Ireland is concentrated in the south-east of the country while the bulk of organic soils occur in the midlands and west. Consequently, nitrogen inputs due to the cultivation of organic soils can be taken as negligible. The equation for estimating N₂O emissions in Ireland reported in sub-category 4.D.1 *Direct Soil Emissions* therefore becomes

$$N_2O_{\text{direct}} = (F_{\text{SN}} + F_{\text{AM}} + F_{\text{S}} + F_{\text{BN}} + F_{\text{CR}}) * EF_1$$

Where

$$F_{\text{SN}} = N_{\text{fert}} * (1 - \text{Frac}_{\text{GASF}})$$

$$F_{\text{AM}} = [N_{\text{ex}} * (1 - \text{Frac}_{\text{GRAZ}}) * (1 - \text{Frac}_{\text{GASM1}})] - N_2O\text{-}N_{\text{hs}}$$

$$F_{\text{S}} = \text{SS}_i * \text{NSSF}$$

$$F_{\text{BN}} = \sum_i \text{Crop}_i * (1 + \text{Res}_i/\text{Crop}_i) * \text{DMF}_i * \text{NCRF}_i$$

$$F_{CR} = \sum_j \text{Crop}_j * \text{Res}_j / \text{Crop}_j * \text{DMF}_j * \text{NCRF}_j$$

and

N_{fert} = total amount of synthetic fertilizer nitrogen applied to soils (kg N)

$\text{Frac}_{\text{GASF}}$ = fraction of synthetic fertilizer nitrogen that volatilizes as NH_3 (0.016 in 2005)

N_{ex} = total amount of animal manure nitrogen excreted by livestock (kg N)

$\text{Frac}_{\text{GRAZ}}$ = fraction of N_{ex} that is excreted by livestock during grazing

$\text{Frac}_{\text{GASM1}}$ = fraction of animal manure nitrogen that volatilizes as NH_3 during housing, manure storage and landspreading (0.493 in 2005)

$\text{N}_2\text{O-N}_{\text{hs}}$ = amount of animal manure nitrogen emitted as N_2O in housing and storage (kg $\text{N}_2\text{O-N}$)

SS_i = quantity of sewage sludge spread on agricultural lands (kT)

NSSF = nitrogen fraction of sewage sludge (3 percent of dry solids)

Crop_i = production of nitrogen-fixing crop i (kT)

$\text{Res}_i / \text{Crop}_i$ = residue to crop product mass ratio of nitrogen-fixing crop i

DMF_i = dry matter fraction of nitrogen-fixing crop i

NCRF_i = nitrogen fraction of nitrogen-fixing crop i

Crop_j = production of crop j (including nitrogen-fixing crops) (kT)

$\text{Res}_j / \text{Crop}_j$ = residue to crop product mass ratio of crop j (including nitrogen-fixing crops)

DMF_j = dry matter fraction of crop j (including nitrogen-fixing crops)

NCRF_j = nitrogen fraction of crop j (including nitrogen-fixing crops)

The annual statistics on nitrogen fertilizer use (N_{fert}) are obtained from the Department of Agriculture and Food while the organic nitrogen inputs (N_{ex}) are known from the analysis in the previous section in relation to manure management. Significant proportions of the nitrogen applied to soils in synthetic fertilizers and animal manures are normally volatilized as NH_3 with some additional conversion to NO_x . These proportions, $\text{Frac}_{\text{GASF}}$ and $\text{Frac}_{\text{GASM}}$ respectively in the IPCC guidelines, must be taken into account in order to determine the amount of nitrogen available for direct N_2O production. The IPCC good practice guidance gives the default proportions of chemical fertilizer and animal manure nitrogen lost in this way as 10 percent and 20 percent, respectively. The volatilization rates for Ireland are however determined from an elaborate NH_3 inventory for agriculture and it is assumed that nitrogen lost as NO_x is negligible in comparison to NH_3 . In addition, $\text{Frac}_{\text{GASM}}$ is split into $\text{Frac}_{\text{GASM1}}$ and $\text{Frac}_{\text{GASM2}}$ with $\text{Frac}_{\text{GASM1}}$ referring to $\text{NH}_3\text{-N}$ losses from animal manures in housing, storage and landspreading and $\text{Frac}_{\text{GASM2}}$ being the proportion of nitrogen excreted at pasture that is volatilised as NH_3 . The 2005 values of $\text{Frac}_{\text{GASM1}}$ and $\text{Frac}_{\text{GASM2}}$ are 0.493 and 0.038, respectively indicating an overall volatilisation rate of 0.195 for animal manure nitrogen.

The expression for F_{AM} given above is used to estimate the amount of animal manure nitrogen ultimately available for direct application to agricultural soils. It is more precise than that given in the IPCC good practice guidance, as the nitrogen in animal manures emitted as N_2O and as NH_3 during animal housing and storage of manures is deducted from total nitrogen excreted in housing. Accordingly, the fraction $\text{Frac}_{\text{GASM1}}$ used here refers to the loss of nitrogen by volatilization as NH_3 during housing and storage together with that from landspreading. These modifications have been made to achieve more accurate accounting of nitrogen and to maintain consistency with Ireland's Tier 2 inventory of NH_3 . The fractions $\text{Frac}_{\text{GASF}}$ and $\text{Frac}_{\text{GASM1}}$ are estimated at 0.016 and 0.493, respectively in 2005 from the NH_3 inventory. Published estimates of sludge production (Smith et al, 2004) and the proportion applied on agricultural lands are used to estimate F_s on the basis of 3 percent nitrogen content in sewage sludges with typical dry solids content of 25 percent (Fehily Timoney, 1985). The estimate of F_s is included in $\text{N}_2\text{O}_{\text{direct}}$ without deduction for volatilisation and the value is added to F_{AM} for reporting purposes in CRF Table 4.D. Although the amount of

sludge spreading on land is increasing, it contributed less than 1 percent of the organic nitrogen input to agricultural soils in 2005.

The Tier 1b method given by the IPCC good practice guidance is used to estimate the nitrogen contributions from nitrogen-fixing crops (F_{BN}) and from crop residues (F_{CR}) returned to the soil. Annual crop production statistics and the default values of nitrogen content and other input parameters given by the IPCC good practice guidance are the basis for these estimates. The IPCC default value of 0.0125 kg N_2O -N/kg N is currently used for EF_1 to estimate direct emissions of N_2O from the inputs calculated from the above equations. The direct emissions of N_2O in 2005 for category 4.D.1 *Direct Soil Emissions* amounted to 8.67 Gg, of which synthetic fertilizers accounted for 6.80 Gg, 1.5 Gg was due to land spreading of animal manures and crops produced 0.34 Gg.

6.5.2 4.D.2 Pasture Range and Paddock Manure

The direct N_2O emissions associated with nitrogen excretion by animals during grazing is not allocated to sub-category 4.D.1 *Direct Soil Emissions* but is reported instead in the CRF under 4.D.2 *Animal Production*. The amount of organic nitrogen input concerned ($N_{ex} * \text{Frac}_{\text{GRAZ}}$) from the equations above, is large in Ireland due to the relatively short period that cattle remain in housing and the contribution from large sheep populations, the majority of which are not housed. The fraction $\text{Frac}_{\text{GASM2}}$, estimated at 0.037 for 2005, is used to adjust nitrogen inputs from grazing animals due to NH_3 volatilisation. The direct N_2O emission factor (EF_3) for this unmanaged nitrogen input is 0.02 kg N_2O -N/kg N and the estimate of emissions in 2005 was 8.94 Gg.

6.5.3 4.D.3 Indirect Emissions

The IPCC methodology for indirect emissions reported in CRF sub-category 4.D.3 *Indirect Emissions* is based on a simple approach that allocates emissions of N_2O due to nitrogen deposition resulting from NH_3 and NO_x emissions in agriculture and from nitrogen leaching to the country that generated the source nitrogen. The contributions from NH_3 and NO_x emission sources in other sectors, such as transport and stationary combustion, are excluded and the import of nitrogen from other countries through atmospheric transport and runoff is not considered. Accordingly, the total nitrogen volatilized as NH_3 , deducted from total nitrogen inputs in synthetic fertilizers and animal manures for estimating the amount contributing to direct N_2O emissions as described in the previous section, becomes the input value of nitrogen used to calculate indirect emissions due to deposition, as follows

$$N_2O_{\text{indirect-dep}} = [N_{\text{fert}} * \text{Frac}_{\text{GASF}} + (N_{\text{ex}} * (1 - \text{Frac}_{\text{GRAZ}}) * \text{Frac}_{\text{GASM1}}) + N_{\text{ex}} * \text{Frac}_{\text{GRAZ}} * \text{Frac}_{\text{GASM2}}] * EF_4$$

$$N_2O_{\text{indirect-leach}} = [N_{\text{fert}} + F_{\text{AW}} + N_{\text{ex}} * \text{Frac}_{\text{GRAZ}}] * \text{Frac}_{\text{LEACH}} * EF_5$$

where

$N_2O_{\text{indirect-dep}}$ = the indirect emissions of N_2O due to atmospheric nitrogen deposition

$N_2O_{\text{indirect-leach}}$ = the indirect emissions of N_2O due to nitrogen leaching

$\text{Frac}_{\text{GASM2}}$ = fraction of animal manure nitrogen that volatilizes as NH_3 during grazing (0.037 in 2005)

$\text{Frac}_{\text{LEACH}}$ = fraction of synthetic fertilizer nitrogen and animal manure nitrogen that leaches from agricultural soils (0.1 in 2005)

EF_4 = N_2O emission factor for nitrogen inputs from atmospheric deposition

EF_5 = N_2O emission factor for nitrogen leaching

The expressions for $N_2O_{\text{indirect-dep}}$ and $N_2O_{\text{indirect-leach}}$ are slightly modified to be consistent with those for estimating direct emissions above and to account for the two separate volatilisation fractions $\text{Frac}_{\text{GASM1}}$ and $\text{Frac}_{\text{GASM2}}$. There is no contribution to $N_2O_{\text{indirect-dep}}$ from F_S , the nitrogen input from sludge spreading, but F_S increases $N_2O_{\text{indirect-leach}}$ through its inclusion in F_{AW} . The default value for $\text{Frac}_{\text{LEACH}}$, the fraction of nitrogen lost through leaching, in the IPCC Guidelines is 30 percent. Estimates of the nitrogen loads in Irish rivers reported under the OSPAR Convention (NEUT, 1999) suggest that approximately 10 percent of all applied nitrogen in Irish agriculture is lost through leaching. This level of leaching is also indicated by farm budget studies where the nitrogen runoff equivalent to 60 kg N/ha has been measured in streams adjoining farmland receiving 200 kg N/ha from chemical fertilizer and 100 kg N/ha from animal manures per year. The value of 0.1 is considered to be a more realistic estimate of $\text{Frac}_{\text{LEACH}}$ than the default value of 0.3 and it is used for 2005, as it was for previous years.

The IPCC default values of the emission factors EF_4 and EF_5 (0.01 kg N_2O -N/kg NH_3 -N emitted for synthetic fertilizer and animal waste nitrogen and 0.025 kg N_2O -N/kg N leached) are used to estimate indirect N_2O emissions. Total indirect emissions in 2005 amounted to 4.23 Gg N_2O , or approximately 49 percent of direct emissions from soils (sub-category 4.D.1).

6.6 QA/QC in Agriculture

The inventory agency has discontinued the use of the IPCC software in the compilation of the CH_4 and N_2O emissions in Agriculture. Instead, a new spreadsheet system developed for the 2006 submission is used to calculate these emissions in a more efficient and transparent manner, which takes into account the strong links to Ireland's Tier II inventory of NH_3 in *Agriculture* and other factors relevant to a more complete country-specific application of the IPCC good practice guidance. The general QA/QC procedures set down in Ireland's new QA/QC plan (section 1.6) have been undertaken in this compilation and inventory management system, from which the time-series outputs may be readily imported to the CRF Reporter. The spreadsheets incorporate transparent linking between input data and calculations as well as internal checks on the calculations and the outputs are directly compatible with the CRF Reporter Tool. The entire compilation for 2005 and all previous years was reviewed externally by a technical person from the Department of Agriculture and Food as an important element of quality assurance for the 2007 submission.

The intensive collaboration between inventory experts and researchers involved in developing the improved inventory methodologies for both CH_4 and NH_3 together with assessment and endorsement of the outcomes by other experts in TEAGASC and the DEHLG adds significantly to the quality and reliability of the emissions estimates produced in this system according to the IPCC good practice guidance.

6.7 Recalculations in Agriculture

Recalculated estimates of emissions in the Agriculture sector are due to the following inventory revisions carried out in the 2007 reporting cycle for the years 1990-2004.

- a) A calculation error was found in the estimates of the total NH_3 -N lost, which is used in the calculation of F_{AW} (N input from manure after adjustment for volatilization) for the years 1990-2004 inclusive.
- b) The emission estimates for CH_4 from manure management for laying hens in the 2006 submission used the emission factor for broilers.

The recalculations undertaken as a result of items (a) and (b) on emission estimates from the agricultural sector are presented in Table 6.4. The quantitative effect of the recalculation of total $\text{NH}_3\text{-N}$ lost used in the calculation of $\text{Frac}_{\text{GASM}}$ is an average decrease in N_2O emissions from 4.D.1 *Direct Soil Emissions* and 4.D.3 *Indirect Soil Emissions* of 0.65 Gg N_2O and 0.14 Gg, respectively for the years 1990-2004 inclusive (Table 6.4). The quantitative effect of item (b) above is a net increase in CH_4 emissions from 4.B *Manure Management* of approximately 4 Gg CH_4 for the years 1990-2004 inclusive. When expressed in terms of CO_2 equivalent, net emissions from *Agriculture* are reduced by approximately 1 percent over the period 1990-2004 inclusive.

6.8 Improvements in Agriculture

Clearly, it is important that high priority is given to emissions of CH_4 and N_2O from agricultural sources in Ireland so that they may be quantified as reliably as possible, given their large overall contributions to the national total (Chapter Three). A large number of input variables determine the emissions in the case of both gases and the final results are very sensitive to changes in many of these variables. Assumptions relating to some parameters have an important bearing on the outcome. While the IPCC methodologies for the agricultural emission sources that are relevant in Ireland are now very comprehensive, they remain generalised and necessarily simplified considering the complex systems and processes that produce the CH_4 and N_2O emissions. The key to developing better estimates and reducing uncertainty is to take full account of national circumstances of climate, soil types, livestock and crop production practices, management systems and other influencing factors in a robust and justifiable manner when applying these methodologies.

The inventory agency made substantial improvement in the overall inventory compilation for *Agriculture* during 2005, particularly with respect to CH_4 emissions, and now achieves closer compliance with the IPCC good practice guidance by the use of Tier 2 methods for CH_4 emissions from enteric fermentation and manure management in cattle. The agency has also been intensively engaged with researchers working on the DNDC model (Li et al, 1994, 1996) that quantifies N_2O emissions from soils with a view to adopting a methodology for such emissions that systematically accounts for the influences of soil type, fertilizer type and application rates, temperature and rainfall, which are not captured by the current IPCC methodology. The results of both field and lysimeter studies conducted in Ireland in recent years suggest that N_2O emission rates from agricultural soils may be substantially higher than the value of 1.25 percent given by the current IPCC default emission factor. The DNDC model has been applied at both regional and national scale and the outcomes were assessed at a workshop in September 2005 attended by the developer of the model and by other eminent agricultural experts in Ireland and the UK. Soil properties, particularly bulk density and organic carbon content, were found to be the key model variables controlling N_2O and it was concluded that a lack of reliable data on these two parameters precludes routine application of the DNDC model at this stage for the purposes of annual emissions reporting. Nevertheless, the model is a very useful supplementary method for assessing emissions, especially in the analysis of the relative importance of all input variables and their effects on total N_2O emissions. Field studies on N_2O fluxes are ongoing and the use of DNDC is under review.

Table 6.4 Percentage Change in Emissions from Agriculture due to Recalculations

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
2006 Submission - Total CH ₄ (Gg CH ₄)	550.668	558.194	560.647	564.587	563.208	564.393	580.958	594.528	602.02	581.495	554.109	552.672	546.506	541.712	542.76
2006 Submission - Total N ₂ O (Gg N ₂ O)	24.762	24.851	24.797	25.266	25.901	26.551	26.63	26.119	27.629	27.665	26.054	24.806	24.459	25.037	24.464
2007 Submission - Total CH ₄ (Gg CO ₂)	554.843	562.191	565.599	568.613	567.015	567.404	584.709	598.044	605.479	584.869	557.557	556.349	550.057	545.904	546.93
2007 Submission - Total N ₂ O (Gg CO ₂)	23.908	23.966	23.901	24.346	24.986	25.618	25.704	25.174	26.804	26.815	25.249	24.017	23.672	24.258	23.692
Manure Management - CH ₄	4.175	3.998	4.952	4.026	3.806	3.011	3.751	3.516	3.459	3.373	3.448	3.677	3.551	4.192	4.17
Manure Management - N ₂ O	0.001	-0.002	-0.001	-0.008	-0.007	-0.007	-0.005	-0.002	0.002	-0.002	-0.002	-0.002	-0.001	-0.002	-0.004
Animal Production N ₂ O	-0.118	-0.125	-0.129	-0.127	-0.124	-0.12	-0.117	-0.118	NO	NO	NO	NO	NO	NO	NO
Agricultural Soils - Direct N ₂ O	-0.602	-0.618	-0.625	-0.641	-0.64	-0.659	-0.658	-0.676	-0.689	-0.707	-0.669	-0.656	-0.655	-0.648	-0.64
Agricultural Soils - Indirect N ₂ O	-0.135	-0.139	-0.141	-0.144	-0.144	-0.147	-0.146	-0.15	-0.138	-0.141	-0.134	-0.131	-0.131	-0.13	-0.128
TOTAL CH ₄ (Gg CH ₄)	4.175	3.997	4.952	4.026	3.807	3.011	3.751	3.516	3.459	3.374	3.448	3.677	3.551	4.192	4.17
TOTAL N ₂ O (Gg N ₂ O)	-0.854	-0.885	-0.896	-0.92	-0.915	-0.933	-0.926	-0.945	-0.825	-0.85	-0.805	-0.789	-0.787	-0.779	-0.772
Total % Change in Emission from Agriculture	-0.921	-0.979	-0.894	-1.019	-1.026	-1.125	-1.019	-1.065	-0.862	-0.927	-0.898	-0.867	-0.889	-0.803	-0.799

Chapter Seven

Land-Use, Land-Use Change and Forestry

7.1 Introduction

Following the publication of the IPCC Special Report on Land Use, Land-Use Change and Forestry and adoption of the IPCC good practice guidance on LULUCF, the source category classification for reporting on the LULUCF sector was revised by Decision 13/CP.9 to that given in Table 7.1. The six top-level categories are used to represent managed land areas and they are broadly defined to accommodate all land areas in most countries, taking into account possible differences in national classification systems. Each category is split into two sub-categories, which may be further sub-divided to reflect national circumstances and the level of detail considered most appropriate for the estimation of relevant emissions and removals. The conversion sub-categories allow for the tracking of land to the principal fixed categories by assuming that a unit of land subject to a change of use remains in the conversion sub-category for 20 years before it is reported in the top-level category to which it has been converted. The revised area-based approach is intended to make the best use of the various types of data likely to be available for the given categories of land and reduce possible overlaps and omissions in reporting for national total land areas.

The net emissions of CO₂ to, or removals of CO₂ from the atmosphere are to be reported with respect to overall carbon gain or loss for up to four relevant carbon pools for the defined land categories. These pools are above-ground biomass, below-ground biomass, dead organic matter (litter and dead wood) and soils. The good practice guidance on LULUCF provides basic methodologies for calculating changes in carbon pools where land areas form the basic activity data and carbon stock change is determined from a number of other parameters. Various levels of land sub-division may be used to capture differences due to climate, management system, vegetation type or other factors influencing carbon exchange. As for other sectors of the inventory, the guidance also provides higher tier methods for estimating emissions and removals, which may be used if the necessary data are available. The liming of agricultural lands, which produces CO₂ emissions, is another important source included in the LULUCF sector. Emissions of N₂O in the LULUCF sector are reported for such activities as nitrogen fertilization of forest land, soil disturbance associated with land-use conversion to cropland and optionally for land drainage while taking into account potential overlap with the *Agriculture* sector in some cases. Emissions of N₂O and CH₄ are also to be reported for biomass burning.

7.2 Overview of LULUCF

7.2.1 Sector Coverage

The 2006 inventory submission included the results of Ireland's first attempts to comply with the reporting requirements of Decision 13/CP.9 for the LULUCF sector. Following the same approach, complete coverage of the relevant gases has been achieved for the years 1990-2005 in all IPCC land categories, as indicated by Table 7.1, whereas in previous submissions (pre 2006) Ireland reported CO₂ estimates only in respect of carbon stock change in forests and CO₂ emissions from the liming of agricultural soils. The reporting of estimates for all

land-use categories in LULUCF represented a major improvement in terms of inventory completeness for Ireland. This chapter presents a broad description of data treatment and the use of default methods to estimate emissions and removals for the relevant land categories in the extended time-series 1990-2005. A more detailed report on the work undertaken to report on the LULUCF sector in 2006 is available (O'Brien, 2007).

The 2007-inventory submission follows the same methodology as that used in the 2006 submission, with some revisions and the use of updated datasets where appropriate. The inventory estimates for 2005 presented here are provisional, pending the availability of final quality assured agricultural statistics for 2005. The provisional estimates for 2005 are based on linear extrapolation of trends in agricultural land-use change in 2003 and 2004. These revisions and others undertaken for this submission are outlined in section 7.10.

The estimates of emissions and removals from LULUCF over the period 1990-2005 are presented in Table 7.2 for all land-use categories. Potential emissions of N₂O and CH₄ are not reported by Ireland according to the following rationale

- The amount of nitrogen fertilizer used for forests is negligible compared to that in agriculture and therefore all N₂O emissions from fertilization are reported under in the Agriculture sector and the notation IE is used in CRF Table 5(I);
- The N₂O emissions from drainage of forest lands and wetlands is an optional reporting category for which Ireland has elected not to include at this time and hence the notation NA appears in CRF Table 5(II);
- No information is available on biomass burning and this activity is assumed not to occur and therefore the notation NO applies in respect of CH₄ and N₂O emissions from biomass burning in CRF Table 5(V).

The data in Table 7.2 show that the LULUCF sector was a significant net source of emissions in the first half of the time-series and was a net sink of carbon in most years thereafter. This outcome is determined mainly by the balance between the removals in category 5.A *Forest Land* and the emissions from 5.C *Grassland* and from lime applications. The most important individual emission categories over the time-series are the carbon release from soils in 5.A.2 *Land Converted to Forest Land* and the CO₂ emissions from agricultural lime application on Grassland. The increase in carbon stocks in living biomass in the category 5.A.1 *Forest Land remaining Forest Land* is the dominant removal that offsets CO₂ emissions. The Wetland, Settlements and Other Land categories are comparatively unimportant in terms of emissions or removals but Cropland constitutes a significant net source towards the end of the time series. The results contained in the 2007 submission for the years 1990-2005 for the LULUCF sector according to the requirements of Decision 13/CP.9 are not directly comparable with those provided in respect of land use change and forestry in submissions prior to 2006.

7.2.2 Land Use Definitions and Land Use Change Matrices

Table 7.3 summarises the definitions and coverage of the IPCC land-use categories in the LULUCF sector as they relate to Ireland along with the data sources that have been used for estimating the respective areas remaining in the category and areas converted to the category and their associated greenhouse gas emissions and removals. The IPCC *Wetlands* category has been split into natural unexploited wetlands, and peatlands, which are wetlands drained for the purpose of commercial and domestic harvesting of peat for combustion or horticultural use.

Table 7.1. Level 3 Category Coverage for Land Use, Land-Use Change and Forestry

5 Land Use Land-Use Change and Forestry	Carbon Stock Change Emissions of CO ₂			CH ₄	N ₂ O
	Biomass	DOM	Soils		
A. Forest Land					
1. Forest Land remaining Forest Land	All	All	NO*	NA	IE
2. Land converted to Forest Land	All	All	All	NA	IE
B. Cropland					
1. Cropland remaining Cropland	NO	NO	NO*	NA	NE
2. Land converted to Cropland	All	NO	All	NA	NE
C. Grassland					
1. Grassland remaining Grassland	NO	NO	NO*	NO	NE
2. Land converted to Grassland	All	NO	All	NO	NE
D. Wetlands					
1. Wetlands remaining Wetlands	All	NO	All	NO	NE
2. Land converted to Wetlands	NO	NO	NO	NO	NE
E. Settlements					
1. Settlements remaining Settlements	NO	NO	NA	NO	NE
2. Land converted to Settlements	All	NO	All	NO	NE
F. Other Land					
1. Other Land remaining Other Land	NO	NO	NO*	NO	NO
2. Land converted to Other Land	NO	NO	All	NO	NO
G. Other					
Agricultural Lime Application	NA	NA	All	NA	NA

DOM : dead organic matter

All : all emission sources covered; NE : emissions not estimated; NO : activity not occurring; NA : not applicable (no emissions of the gas occur in the pool/source category); IE : emissions included elsewhere.

** Under the Tier 1 method, there is no carbon stock change in soil for these land categories*

Table 7.5 records the land-use changes among the various categories over the period 1990-2005 in the form of land-use change matrices for the individual years. The matrices of land use are intended to show the dynamism of changes in Irish circumstances and to identify the conversions that are most significant in terms of their potential as contributors to either emissions or removals of greenhouse gases over the inventory time-series. The annual totals for individual years in the matrices do not necessarily correspond with the areas that appear as activity data in the various sectoral background data tables in the CRF tables because the latter account for the rolling 20-year transition period that began in 1970. In addition, the area relevant to the biomass pool is not the same as that for the soils pool for *5.A.2 Land Converted to Forest Land* due to the combination of the three national forest area categories and because different lengths of transition period apply to organic and mineral soils. *Grassland* is the dominant land-use category in all years, accounting for 57 percent of total area in 1990. The *Other Land* category was the next largest at almost 11 percent, followed by *Forest Land* which was 7 percent of the total. The major land-use change since 1990 has been the conversion of grassland and peatland to forest land. The area of forests increased by 44 percent from 1990 to 2005, but the proportion of *Forest Land* to total land in the country is still only 10 percent, which is low compared to many Annex I Parties.

7.2.3 Soil Type and Soil Organic Carbon

Soil organic carbon (SOC) is the basic parameter in the IPCC estimation methods for determining carbon stock changes in soils, which is the dominant source of carbon emissions in land conversion categories in LULUCF. The organic carbon status of Irish soils under native vegetation is established from the soil type and the default reference soil organic carbon stocks (SOC_{ref}) for cold, temperate moist regions (GPG Tables 3.2.4, 3.3.3 and 3.4.4). The General Soil Map of Ireland (Gardiner and Radford, 1980) is the basic data

source for soil type. Mineral soils as identified from the general soil map were allocated to the HAC (high activity clay), LAC (low activity clay), sandy and humic soil classes used by IPCC while peats were allocated to the IPCC wetlands class as shown on Table 7.4, based on detailed national assessment of soil carbon stocks in Ireland (Tomlinson, 2005). The values of SOC_{ref} appropriate to each soil association may then be assigned using the correspondence to IPCC classes given in Table 7.4. The distribution of CORINE Land Use over IPCC soil classes was established in the same way to facilitate complete correspondence between land use, soil and SOC_{ref}.

Table 7.4. Soil Class Coverage and Soil Organic Carbon

General Soil Map Soil Association	IPCC Soil Class					Proportion of Soil Association in Area of Ireland
	HAC	LAC	Peaty/ Humic	Sandy Soil	Wetlands Soil	
basin peat					0.34	0.06
brown earth		0.19				0.13
brown podzolic		0.21				0.15
gley		0.30			0.02	0.22
grey brown podzolic		0.30				0.21
lithosol			0.22	1.00		0.04
lowland blanket peat					0.31	0.05
podzol			0.78			0.08
Renzinas	1.00					0.01
upland blanket peat					0.33	0.06
Proportion of IPCC Soil Class in Area of Ireland	0.01	0.71	0.10	0.01	0.17	
SOC _{ref} (t C/ha)	95	85	115	71	87	

7.2.4 Estimation of Emissions from Soils

Mineral Soils

The annual change in SOC in mineral soils over the appropriate transition period determines the carbon emissions or removals for the various land-use conversion categories as follows

$$\Delta C = A * (SOC_0 - SOC_{0-T}) / T \quad (7.1)$$

$$SOC = SOC_{ref} * F_{LU} * F_{MG} * F_I$$

where

- ΔC = annual change in carbon stocks
- A = area of land converted from a former land use
- SOC_0 = soil organic carbon stock for current land use
- SOC_{0-T} = soil organic carbon stock for former land use
- SOC_{ref} = reference soil organic carbon under native vegetation for a given soil type in area A
- T = transition period
- F_{LU} = stock change factor for land use or land-use change type
- F_{MG} = stock change factor for management regime
- F_I = stock change factor for organic matter input

Table 7.2. Emissions^a and Removals^a from Land Use Land-Use Change and Forestry 1990-2005 (Gg CO₂ eq)

IPCC Source Category	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
A. Forest Land	-478.02	-391.91	-201.01	-275.98	-207.58	-245.02	-216.91	-349.44	-505.17	-591.85	-474.15	-624.88	-738.42	-1,060.00	-665.77	-811.38
1. Forest Land remaining Forest Land	-1,078.90	-1,163.38	-884.40	-876.45	-587.48	-437.51	-358.95	-238.61	-521.46	-579.24	-116.18	-37.62	-358.01	-1,258.08	-724.25	-845.19
2. Land converted to Forest Land	600.89	771.47	683.39	600.46	379.90	192.50	142.05	-110.82	16.29	-12.61	-357.96	-587.26	-380.41	198.08	58.48	33.81
B. Cropland	46.83	46.34	42.30	37.46	-32.34	-7.85	75.02	51.66	32.55	8.39	48.39	129.77	120.66	165.77	126.57	148.09
1. Cropland remaining Cropland	46.83	46.34	34.60	-33.48	-60.91	-36.42	-0.27	5.46	-13.66	-37.82	2.19	-5.00	-10.85	-11.88	-2.23	19.29
2. Land converted to Cropland	NE,NO	NE,NO	7.70	70.95	28.58	28.58	75.30	46.20	46.20	46.20	46.20	134.77	131.52	177.65	128.80	128.80
Agricultural Lime Application ^b	38.28	32.72	26.25	36.05	27.69	50.39	49.50	44.55	31.84	39.48	37.32	39.21	29.17	42.04	27.15	26.71
C. Grassland	491.91	559.29	433.02	347.37	176.32	383.12	417.93	500.31	313.87	402.46	473.67	458.89	463.29	559.32	340.65	56.57
1. Grassland remaining Grassland	620.07	582.80	529.72	619.22	522.83	718.00	708.21	648.16	542.97	612.98	598.29	615.30	513.35	612.72	479.50	477.79
2. Land converted to Grassland	-128.16	-23.51	-96.71	-271.84	-346.51	-334.88	-290.28	-147.86	-229.09	-210.51	-124.63	-156.41	-50.05	-53.39	-138.85	-421.22
Agricultural Lime Application ^b	316.76	282.43	229.35	321.25	241.95	444.21	434.53	378.94	273.74	343.75	329.07	346.07	244.72	344.72	213.64	214.09
D. Wetlands	46.61	46.00	45.39	44.79	42.37	39.92	37.48	35.03	32.58	30.13	38.13	34.02	29.91	25.80	21.69	21.27
1. Wetlands remaining Wetlands	46.61	46.00	45.39	44.79	42.37	39.92	37.48	35.03	32.58	30.13	38.13	34.02	29.91	25.80	21.69	21.27
2. Land converted to Wetlands	NE, NO	NE, NO	NE, NO	NE, NO	NE, NO	NE, NO	NE, NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO
E. Settlements	13.69	12.91	13.81	15.24	17.80	18.28	22.81	25.91	29.04	31.18	33.04	44.88	38.29	46.39	52.99	59.38
1. Settlements remaining Settlements	NE, NO	NE,NO	NE,NO	NE, NO	NE, NO	NE, NO	NE, NO	NE, NO	NE, NO	NE, NO	NE, NO	NE, NO	NE, NO	NE, NO	NE, NO	NE, NO
2. Land converted to Settlements	13.69	12.91	13.81	15.24	17.80	18.28	22.81	25.91	29.04	31.18	33.04	44.88	38.29	46.39	52.99	59.38
F. Other Land	-0.21	NE, NO	-12.31	-12.40	-13.93	-6.73	-28.27	-34.33	-39.34	-32.07	-10.79	-23.69	-39.70	-35.98	-70.70	-131.16
1. Other Land remaining Other Land	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
2. Land converted to Other Land	-0.21	NO	-12.31	-12.40	-13.93	-6.73	-28.27	-34.33	-39.34	-32.07	-10.79	-23.69	-39.70	-35.98	-70.70	-131.16
G. Other	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
TOTAL LULUCF	120.81	272.64	321.21	156.48	-17.36	181.73	308.06	229.14	-136.47	-151.76	108.29	18.99	-125.96	-298.70	-194.58	-657.22

a positive values indicate emissions and negative values indicate removals

b the emissions from lime application to grassland and cropland are reported in CRF Table 5(IV) rather than under Grassland in CRF Tables 5.B and 5.C, respectively. These emissions are not included in the totals for 5.C Grassland and 5.B Cropland

Table 7.3. Land Use Categories

Land Use Category	Definition and Coverage	Area 1990 (ha)	Area 2005 (ha)	Sources of Information	Principal Conversions	
					To	From
Forest Land	All public and private plantation forests. Forest land is an area of land where tree crown cover is greater than 20% of the total area occupied or 50% of conventional stocking and includes recently clearfelled areas. It has a minimum width of 20m and a minimum area of 0.1ha and includes all trees with a potential to reach 5m in height. Trees grown for fruit or flowers are excluded, as are woody species such as furze and rhododendron	370,123	536,3149	FIPS (Forest Inventory Planning System) 1995 COILLTE database Forest Service Premiums database LPIS (Land Parcels information System) CORINE Land Cover General Soil Map		Grassland
Cropland	Permanent crops and tillage areas (including setaside) recorded by the Central Statistics Office (CSO)	404,563	368,300	CSO, CORINE Land Cover LPIS (Land Parcels information System)		Grassland
Grassland	Areas of improved grassland (pasture and areas used for the harvesting of hay and silage) and unimproved grassland (rough grazing) in use as recorded by CSO annual statistics	4,147,597	3,935,264	CSO, CORINE Land Cover LPIS (Land Parcels information System) CORINE Land Cover General Soil Map		Other Land
Wetlands	Natural unexploited wetlands	1,227,810	1,150,038	CORINE Land Cover General Soil Map	Peatlands	
Peatland	Wetland areas commercially exploited for public and private extraction of peat and areas used for domestic harvesting of peat	73,980	61,112	Bord na Mona (BNM) area statistics; Expert opinion		
Settlements	Urban areas, roads, airports and the footprint of industrial, commercial/institutional and residential buildings	98,579	114,014	CORINE Land Cover; National Roads Authority (NRA) road construction statistics; CSO housing stock, house completions and other construction floor area statistics; General Soil Map		Grassland
Other Land	Natural grasslands not in use for agricultural purposes, water bodies, bare rock	796,748	995,045	Natural grasslands not in use for agricultural purposes, water bodies, bare rock	Grassland	
Total Land	National territorial area (including inland water bodies and salt marshes and intertidal zones)	7,111,785	7,111,785	CORINE Land Cover		

Table 7.5 Land Use Matrices 1990-1997 (ha)

	Forest Land	Cropland	Grassland	Peatland	Wetland	Settlements	Other Land	Total
1990	370,123	404,563	4,140,385	73,980	1,227,810	98,178	796,747	7,111,785
Forest Land	370,089					35		370,123
Cropland	10,062		4,130,025			297		4,140,385
Grassland	435	401,694	2,408			27		404,563
Peatland	320		61	73,380	219			73,980
Wetland	8,330				1,219,479			1,227,810
Settlements						98,178		98,178
Other Land			17,759			19	778,969	796,747
1991	389,236	401,694	4,150,253	73,380	1,219,698	98,556	778,969	7,111,785
Forest Land	389,199					37		389,236
Cropland	8,680	1,307	4,119,655			317	20,294	4,150,253
Grassland	356	401,309				29		401,694
Peatland	320		61	72,780	219			73,380
Wetland	7,343				1,212,355			1,219,698
Settlements						98,556		98,556
Other Land						21	778,948	778,969
1992	405,898	402,616	4,119,716	72,780	1,212,574	98,959	799,242	7,111,785
Forest Land	405,857					41		405,898
Cropland	8,372	11,556	4,099,263			350	175	4,119,716
Grassland	355	402,229				31		402,616
Peatland	320		61	72,181	219			72,780
Wetland	6,951				1,205,623			1,212,574
Settlements						98,959		98,959
Other Land						23	799,219	799,242
1993	421,855	413,785	4,099,324	72,181	1,205,842	99,404	799,394	7,111,785
Forest Land	421,808					48		421,855
Cropland	10,408		4,085,952			409	2,554	4,099,324
Grassland	486	410,301	2,961			37		413,785
Peatland	320		61	71,581	219			72,181
Wetland	8,245				1,197,597			1,205,842
Settlements						99,404		99,404
Other Land						27	799,367	799,394
1994	441,267	410,301	4,088,974	71,581	1,197,815	99,925	801,922	7,111,785
Forest Land	441,218					49		441,267
Cropland	12,910		4,075,642			422		4,088,974
Grassland	647	409,277	340			38		410,301
Peatland	140		30	70,721	690			71,581
Wetland	10,013				1,187,802			1,197,815
Settlements						99,925		99,925
Other Land			11,835			28	790,059	801,922
1995	464,928	409,277	4,087,847	70,721	1,188,492	100,461	790,059	7,111,785
Forest Land	464,866					62		464,928
Cropland	11,586	7,935	4,032,277			530	35,520	4,087,847
Grassland	611	408,618				48		409,277
Peatland	140		30	69,861	690			70,721
Wetland	8,644				1,179,848			1,188,492
Settlements						100,461		100,461
Other Land						35	790,025	790,059
1996	485,847	416,552	4,032,307	69,861	1,180,538	101,135	825,545	7,111,785
Forest Land	485,777					70		485,847
Cropland	6,523		4,015,156			606	10,022	4,032,307
Grassland	383	414,100	2,014			55		416,552
Peatland	140		30	69,001	690			69,861
Wetland	4,388				1,176,151			1,180,538
Settlements						101,135		101,135
Other Land						40	825,505	825,545
1997	497,211	414,100	4,017,200	69,001	1,176,841	101,906	835,527	7,111,785

Table 7.5 (continued) Land Use Matrices 1997-2004 (ha)

	Forest Land	Cropland	Grassland	Peatland	Wetland	Settlements	Other Land	Total
1997	497,211	414,100	4,017,200	69,001	1,176,841	101,906	835,527	7,111,785
Forest Land	497,131					80		497,211
Cropland	7,112		4,001,102			685	8,301	4,017,200
Grassland	370	408,000	5,668			62		414,100
Peatland	140		30	68,141	690			69,001
Wetland	5,305				1,171,535			1,176,841
Settlements						101,906		101,906
Other Land						45	835,482	835,527
1998	510,059	408,000	4,006,800	68,141	1,172,225	102,777	843,783	7,111,785
Forest Land	509,974					86		510,059
Cropland	7,234		3,998,827			739		4,006,800
Grassland	426	400,900	6,607			67		408,000
Peatland	140		30	67,281	690			68,141
Wetland	4,868				1,167,358			1,172,225
Settlements						102,777		102,777
Other Land			11,936			48	831,798	843,783
1999	522,642	400,900	4,017,400	67,281	1,168,048	103,717	831,798	7,111,785
Forest Land	522,550					91		522,642
Cropland	8,914		4,007,698			787		4,017,400
Grassland	517	400,100	212			71		400,900
Peatland	327		75	65,335	1,544			67,281
Wetland	5,937				1,162,111			1,168,048
Settlements						103,717		103,717
Other Land			33,914			51	797,832	831,798
2000	538,245	400,100	4,041,900	66,421	1,163,655	104,718	796,746	7,111,785
Forest Land	538,120					125		538,245
Grassland	8,934	15,042	3,994,425			1,077	22,422	4,041,900
Cropland	545	399,458				97		400,100
Peatland	327		75	64,477	1,542			66,421
Wetland	5,658				1,157,998			1,163,655
Settlements						104,718		104,718
Other Land						70	796,676	796,746
2001	553,584	414,500	3,994,500	65,580	1,159,540	106,087	817,995	7,111,785
Forest Land	553,477					108		553,584
Grassland	8,696	8,814	3,948,525			926	27,539	3,994,500
Cropland	531	413,886				83		414,500
Peatland	327		75	63,636	1,542			65,580
Wetland	5,501				1,154,039			1,159,540
Settlements						106,087		106,087
Other Land						60	817,935	817,995
2002	568,531	422,700	3,948,600	64,739	1,155,581	107,264	844,371	7,111,785
Forest Land	568,400					131		568,531
Cropland	5,263	13,324	3,928,884			1,129		3,948,600
Grassland	323	422,276				102		422,700
Peatland	327		75	62,795	1,542			64,739
Wetland	3,184				1,152,397			1,155,581
Settlements						107,264		107,264
Other Land			4,941			74	839,356	844,371
2003	577,497	435,600	3,933,900	63,897	1,153,939	108,699	838,253	7,111,785
Forest Land	577,347					150		577,497
Cropland	5,637		3,868,587			1,290	58,386	3,933,900
Grassland	346	422,800	12,338			116		435,600
Peatland	327		75	61,953	1,542			63,897
Wetland	3,429				1,150,510			1,153,939
Settlements						108,699		108,699
Other Land						84	838,169	838,253
2004	587,086	422,800	3,881,000	61,953	1,152,052	110,339	896,555	7,111,785

Table 7.5 (continued) Land Use Matrices 2005 (ha)

	<i>Forest Land</i>	<i>Cropland</i>	<i>Grassland</i>	<i>Peatland</i>	<i>Wetland</i>	<i>Settlements</i>	<i>Other Land</i>	<i>Total</i>
2004	587,086	422,800	3,881,000	63,056	1,152,052	110,339	895,452	7,111,785
Forest Land	586,918					168		587,086
Cropland	5,852		3,774,016			1,445	99,687	3,881,000
Grassland	361	368,300	54,009			130		422,800
Peatland	327		75	61,112	1,542			63,056
Wetland	3,556				1,148,496			1,152,052
Settlements						110,339		110,339
Other Land						94	895,358	895,452
2005	597,014	368,300	3,828,100	61,112	1,150,038	112,176	995,045	7,111,785

The factors F_{LU} , F_{MG} and F_I account for changes in SOC due to management practices that impact on soil carbon. Table 7.6 shows the adjustment factors derived from the product of F_{LU} , F_{MG} and F_I taken from GPG Table 3.3.4 for the land uses defined for Ireland in Table 7.3. Equation 7.1 is the basic Tier I method used for estimating emissions from mineral soils for all land-use categories as described in the following sections. The default transition period of 20 years is applied for all mineral soils. The estimation procedure is performed following a simple approach that provides reasonable first estimates of emissions from soils for the defined land uses in accordance with the IPCC good practice guidance and the available information for the country. It involves the identification and quantification of the land areas subject to a change of use, the application of the data in Table 7.4 to assign SOC_{ref} for the soil types in those land areas and the calculation of carbon stock change on the basis of the factors given in Table 7.6.

Organic Soils

The basic methodology for estimating emissions from organic soils is to assign a direct annual carbon loss rate that accounts for the oxidation of organic matter due to drainage, tillage or disturbance of the land area concerned. The default emission factors of 0.25 t C/ha per year for managed grassland soils and 1 t C/ha per year for cultivated cropland soils in cold temperate climatic regions given in the IPCC good practice guidance are adopted for Ireland. Some information is available to suggest that a transition period shorter than the default duration of 20 is appropriate for some land-use conversions on organic soil, which is taken into account in the analysis described below.

Table 7.6. Adjustment Factors for SOC

Land Use	F_{LU}	F_{MG}	F_I	Adjustment factor, AF
Cropland	0.71	1.09	1.11	0.86
Improved grassland	1.0	1.0	1.14	1.14
Unimproved grassland	1.0	1.0	NA	1.0
Rough grazing	1.0	0.95	NA	0.95
Other agricultural land (Native grassland)	1.0	1.0	NA	1.0

7.3 5.A Forest Land

7.3.1 Carbon Stock Change in Living Biomass

Previous NIRs have described a well-established country-specific Tier 2 methodology used to estimate the annual increase in forest carbon stocks in Ireland's expanding forests. A detailed account of the model used (CARBWARE) and how it was applied for the period 1990-2004 is available (Gallagher et al, 2004). The output from the model has been updated to include 2005 forestry data. The model has been used to calculate the total standing carbon content of forests year-on-year using Irish forest yield models and appropriate values of biomass expansion factor, wood density and carbon content for the various tree species. Wood harvest is determined separately from national statistics and converted to carbon using the same values of biomass expansion factor and carbon content. In the submissions up to 2005, the value of carbon removals reported for a particular year in LUCF Table 5.A of the former CRF is the difference between standing carbon stock at the end of that year and carbon stock at the end of the previous year. This value represents the total for the above-ground biomass and below-ground biomass pools in both *5.A.1 Forest Land Remaining Forest Land* and *5.A.2 Land Converted to Forest Land* under the present reporting regime.

Given that it fully quantifies annual change in forest biomass, the CARBWARE model is retained as the basic methodology for estimating carbon stock increment in LULUCF categories 5.A.1 and 5.A.2 by making the appropriate split between their respective contributing areas on the basis of the age of forests. The model as used to date accounts for total forest area in the following classes

- (i) Areas of young forest from 7 to 25 years of age;
- (ii) Areas of mature forests greater than 25 years old and
- (iii) Cleared and unclassified areas, which are assumed not to store carbon. This area class represents total identified forest area by the Forest Service less covered forest as located by remote sensing and classified in FIPS.

The area representing category *5.A.2 Land Converted to Forest Land* may be readily determined from the area of young forests in class (i) above and the area for category *5.A.1 Land Remaining Forest Land* is then the total productive area less that for category 5.A.2. The allocation of carbon uptake to above-ground biomass and below-ground biomass is achieved by partitioning between these two carbon pools in the ratio 0.8:0.2. Carbon storage in this approach is attributed only to forests old enough to have biomass (i.e. more than six years old), rather than to all planted areas. The CARBWARE model does not cover deforestation and therefore cannot provide information on forest lands converted to other land categories. The following paragraphs summarise the carbon accounting methodology applied for carbon stock change in living biomass in the CARBWARE model. The activity data (forest area, afforestation, harvest, etc) and the estimated carbon stock changes in biomass for *5.A Forest Land* are compiled in Table E.1 of Annex E for the years 1990-2005.

Forest Area and Species

A time series of forest strata by area and age was constructed for the years 1990-2005 using information from the Forest Inventory and Planning System (FIPS) base year of 1995 and the total forest area as given by the Forest Service. The FIPS survey data comprise recorded and interpreted information on areas and species for identified state and private forests. The young crop (7 to 25 years of age inclusive) and mature crop (greater than 25 years) categories in FIPS were broken down by species to provide nine individual strata. A third broad category of cleared/unclassified areas (age up to 7 years) was included so that

the total Forest Service area was accounted for in all years. This includes felled areas in which forest cover had not been re-established, recent plantings less than 7 years old, which are assumed to have no measurable biomass, and other productive un-forested areas.

Having established the basic area-species matrix for 1995, the corresponding data for the years 1996 to 2005 were obtained by growing the forest estate forward in time, using annual data on planting and clear felling rates (Annex E), while taking into account the progression of forested areas between the cleared, young and mature categories on the basis of age. The process was worked in reverse for the years 1994 to 1990 to obtain consistent time-series data for this period, as shown in Annex E.

Volume

The FIPS survey results do not contain wood volume or increment data. Therefore, the volume of stemwood was determined from Irish yield models (Hamilton, 1975 and Forest and Wildlife Service) and is based on periodic current annual increment. The Coillte average weighted yield class (wood production model) was applied to all public and private sector forests for each of the FIPS categories. Main crop volume *after* thinning was used for conifers. The ages assumed for young and mature conifers were 15 and 35 years, respectively. Young broadleaved crops were allocated a nominal standing volume of 10 m³/ha. The volume in mature broadleaved forests was determined from the total timber plus firewood volume recorded in the inventory of private woodlands, divided by area. Mixed mature forest volume was based on an average for the mature other conifers and broadleaves strata. The standing volume is reduced by 15 percent to allow for forest roads and rides. The reduced volumes are multiplied by biomass expansion factors (BEF) of 2 for young forests and 1.4 for mature forests (which give a weighted BEF of 1.64 for all forests) and by dry density in the range 0.35-0.55, depending on species and age, to obtain whole-tree wood volume (m³/ha).

Harvest

Coillte records are the main source of data for wood harvesting. These data (Annex E) are compiled through the company's timber sales reporting system. The annual wood harvest volumes for the main species (broadleaves, spruce, pine and other conifers) are converted to carbon using the average carbon content of 0.5 and weighted biomass expansion factor of 1.64, as in the case of volume increment. Harvest volumes include firewood, which is estimated to be in the region of 30,000 m³/year.

Carbon Stock Increment

The carbon uptake of each FIPS category is calculated by multiplying whole-tree volume by a carbon content of 0.5 and by area. In the original version of the CRF, increment values were used to determine annual increments in carbon stocks and from these the harvest was subtracted to find the net changes in carbon stocks. In the current approach, reduced actual standing volumes (standing volumes less thinnings) on a *net areas basis* are used to estimate standing volume. Annual increment in the latest year is then calculated by subtracting the carbon stock in the previous year from the estimated carbon stock in the latest year. This is the increment less the harvest, as the thinning volumes have already been deducted in the data used and the areas are net of clear-felled volumes.

1990-2005 Time Series Data

The carbon stock change estimates for living biomass in forests are slightly revised on those given in the 2006 submission due to the effect of some updated information for afforestation in 2004 provided by COFORD.

7.3.2 Carbon Stock Change in Dead Organic Matter

Dead organic matter consists of the dead wood and litter pools. For dead wood the Tier 1 approach is used, which assumes that input is equal to output and therefore the net carbon stock change is zero (Section 3.2.1.2 of the good practice guidance for LULUCF). In the case of litter, the default net litter accumulation values for wet temperate forests are adopted, as outlined in Table 3.2.1 of the good practice guidance for LULUCF and the stock change is estimated with reference to young and mature forests separately. The values from Table 3.2.1 are 0.3 t C/ha/yr for broadleaves and 0.5 t C/ha/yr for conifers in young forests (less than 20 years) and 0.8 t C/ha/yr for broadleaves and 1.3 t C/ha/yr for conifers in mature forests (greater than 20 years).

It is assumed that afforested and reforested areas less than 7 years old do not contribute to the dead organic pool, which is consistent with the approach above for estimating the carbon stock change in living biomass. Young forest areas, computed for rolling 20-year periods (e.g. 1971-1990, 1972-1991 etc), are split as 7 percent broadleaf and 93 percent conifer based on the species distribution in the 1995 forest inventory baseline year. The area of mature forests is calculated as total forest area less young forest area and reforested area for the preceding seven years. This represents the litter producing area assuming there is no litter input in the first 7 years following reforestation. It is also assumed that the reforested area equals the harvested area. The broad leaf to conifer split for old forests is 31 percent and 69 percent, respectively, again on the basis of the 1995 baseline year distribution.

For comparison it may be noted that experimental data for Sitka spruce stands up to 20 years old show a net accumulation (net decomposition) of 0.8 to 3 t C/ha/year and about 2 t C/ha/year for older stands, based on litter inputs of 1 to 4 t C/ha/year. The decomposition was based on a mean residence time (total litter/annual input rate) for litter on wet mineral gley soils for Sitka spruce of about 5 to 7 years (i.e. 14-20 percent of the litter is decomposed annually). These values are slightly higher than the defaults but the default values are adopted because we do not have sufficient country specific data.

7.3.3 Net Carbon Stock Change in Soils

5.A.1 Forest land Remaining Forest Land

Under the Tier 1 approach it is assumed that the carbon stock in soil organic matter for category *5.A.1 Forest Land remaining Forest Land* remains constant, regardless of changes in forest management, forest type and disturbance. The notation key NO is therefore used under this item in CRF Table 5.A.

5.A.2. Land Converted to Forest Land

There has been an annual increase in the national forest area since 1970. Initially, the lands converted to forestry were of relatively poor quality, with marginal potential for economic returns under other agricultural practices. In more recent years, and especially with the increase in private afforestation, land of higher quality has been converted, reflecting improved grant-aid under the afforestation programme, the decline in economic returns for conventional farming practices and a preference for less labour-intensive land usage. In order to maintain consistency, the activity areas for estimating carbon stock changes in soil are those used by COFORD in the estimation of carbon stock changes in forest biomass and in dead organic matter. The same split of the total for *5.A.2.Land Converted to Forest Land* is applied for all years (Table E.1 of Annex E) with *5.A.2.3 Wetlands Converted to Forest Land* accounting for 57 percent of the area.

The GIS analysis provided the means by which to allocate the land areas in the conversion categories 5.A.2.1 through 5.A.2.5 to soil classes using Table 7.5. The agricultural lands converted to forest land were determined from the LPIS (Land Parcel Information Systems) database, supplied by the Forest Service, which records the conversions as spatially defined areas. The Forest Service GIS database is a comprehensive description of all existing holdings and activities back to 1920. This database system provides detailed information on individual land conversion areas and plantation date from 1990 for private afforestation under grant-aided schemes. Prior to 1990, total annual afforestation area was used. It was assumed that planting practice was consistent with the practices in the early 1990's, and therefore forest areas were allocated to the various soil types in the same proportions as prevailed in the early 1990's.

The afforested areas were superimposed on the general soil map and the CORINE 1990 Land Cover Map of Ireland (level 6). This overlay combination delineated the individual areas and underlying soil type of afforested lands. It also revealed the plantation date and gave an indication of the previous land use. The previous land use given by CORINE was used as a general guidance. Where the previous land use was clearly anomalous, for example where it was indicated by CORINE that the afforested area was a water body, it was assumed that the trees were actually planted on a sub-area of unimproved grassland, which is included in the category 5.F Other Land. Although there is evidence that afforestation on mineral soils has little or no impact on the carbon stock within mineral soils under Irish conditions, there is not sufficient published data to apply a country-specific or region-specific emissions scheme. Therefore the Tier 1 IPCC defaults in the good practice guidelines were used. Accordingly, afforestation on mineral soils has been assigned the default transition period of 20 years, requiring evaluation of new forests on mineral soils from 1970 onwards. Carbon stock changes for afforested areas on mineral soils were estimated using Equation 7.1.

Afforestation occurs on mineral and organic soils at a ratio of approximately 60:40. Recent forest research in the UK (Hargreaves et al, 2003) under climatic and organic soil conditions similar to those in Ireland suggests that following plantation, organic soils emit carbon at an elevated rate of approximately 16 t C/ha over a typical period of 4-5 years. This implies an emission rate of 4 t C/ha annually over a transition period of 4 years. Thereafter the emission from afforested organic soils reduces to zero, or indeed the soil may become a modest sink of carbon. While the emission rate is large compared to the default rate of 0.68 t C/ha/year for organic soils in cold wet temperate conditions, the transition period is much shorter than the 20-year default period. The accumulated default emission of 13.6 t C/ha over 20 years is only 15 percent less than total emissions according to the UK findings (Hargreaves et al, 2003). A country specific transition period of four years is therefore considered appropriate to afforested areas on organic soils.

7.3.4 Emissions of Non-CO₂ Gases

Ireland does not report emissions of non-CO₂ greenhouse gases for *5.A Forest Land*. The amount of synthetic fertilizer used in forests is negligible compared to that used in agriculture and therefore all N₂O emissions from fertilizer applications are reported under agriculture. The notation IE is therefore used in CRF Table 5(I). No estimates have been made of N₂O emissions associated with the drainage of forest soils, an optional reporting category in the LULUCF sector, and the notation key NA is used in CRF Table 5(II). Biomass burning does not occur in Ireland with the result that the notation key NO is used in CRF Table 5(V) for the gases concerned.

7.4 5.B Cropland

7.4.1 Cropland Areas

Cropland areas are based on CSO annual statistics for tillage crops, revised by the inventory agency to account for inconsistencies due to the impact of changes in total farmed area reported in 1997, as described in section 7.10. At the time of writing this NIR, the CSO had not released a complete set of figures for crop areas for 2005. The missing data have been estimated based on an extrapolation of the 2003 and 2004 figures.

Croplands are assumed to revert to natural grassland status during set-aside (the temporary exclusion of tillage areas from production), but stay within the category *5.B Croplands Remaining Croplands*, as a land parcel that is given over to set-aside in one year will probably be tilled in subsequent years. The CSO data include set-aside areas within the Other Crops class. This area of Other Crops is used as the upper limit to give a conservative estimate of set-aside area. In order for the net change in cropland to correspond to that indicated by the CSO statistics, the cropland areas lost to *5.A Forest Land* and *5.E Settlements* must be offset by new lands converted from *5.C Grassland*. This is achieved by adding those areas of cropland in transition to forest lands and settlements to the area of land in transition to cropland, and deducting an equal amount from the area under *5.B.1 Croplands Remaining Croplands*. The relevant emissions and removals are determined by net carbon stock changes in living biomass and soils for *5.B.2 Lands Converted to Cropland*.

7.4.2 Carbon Stock Change in Biomass

The stock change relates only to above-ground biomass and the estimation is based on the difference between initial and final carbon content of biomass for the lands converted. In the conversion of land to cropland, it is assumed under the Tier I approach that the dominant vegetation from the initial land use is removed entirely. The carbon stock change is then quantified as the net sum of carbon lost on conversion and the carbon added by the first year's growth of crops. Grassland is the only relevant land-use type undergoing conversion to cropland in Ireland. The dry matter content of grassland is taken as 13.6 tonnes/ha and the carbon content of dry matter is 0.5 percent. The default value of 5 t C/ha from GPG Table 3.3.8 is adopted for the carbon stock in crop biomass after one year. The carbon stock change in biomass on the area A converted to cropland is then calculated from Equation 3.3.8 of the IPCC good practice guidance as follows

$$\Delta C = A * [(C_{\text{after}} - C_{\text{before}}) + \Delta C_{\text{growth}}] \quad (7.2)$$

$$\Delta C = A * [(13.6 * 0.5 - 0.0) + 5.0]$$

7.4.3 Carbon Stock Change in Soils

The spatial distribution of cropland areas over IPCC soil class is derived from GIS analysis of LPIS 2004 data, superimposed on the general soil association map. The GIS analysis shows that a very high proportion (98 percent) of croplands are located on Low Activity Clay (LAC) soils. It is assumed that only grasslands on LAC soils are suitable for direct conversion to croplands, which is consistent with the requirement for cropland productivity. It is therefore reasonable to assume that all grassland areas converted to croplands are also on LAC soils and that no other land categories are converted to croplands.

Carbon stock changes in mineral soils are estimated using Equation 7.1. Farm management and input practices are assumed to have been constant over the inventory period for established croplands. Therefore the SOC will not have changed for mineral soils, with the exception of those lands going to set-aside for short periods within the transition period of 20

years. Organic soils continue to lose carbon until such time as they are no longer organic. It is assumed that none of the soils classified as organic managed to change this status during the inventory period. Therefore all croplands that were on organic soils in 1990 continue to be assigned to organic soils.

7.4.4 N₂O Emissions in Cropland

Soil disturbance associated with land-use conversions to cropland result in minor emissions of N₂O. Such emissions are estimated from the change in soil organic carbon over the 20 year transition period, obtained using Equation 7.1 for the land-use and soil type converted to cropland, and the soil C:N ratio as follows

$$N_2O = (\Delta C / R_{C:N}) * 44/28 \quad (7.3)$$

where ΔC is the annual change in carbon stocks given by Equation 7.1 and $R_{C:N}$ is the C:N mass ratio in soil organic matter for which a default value of 15 is given in the IPCC good practice guidance.

7.5 5.C Grassland

7.5.1 Grassland Areas

Grassland is the dominant land-use category in Ireland. Areas are based principally on CSO annual statistics on improved grassland (pastures and areas harvested for silage and hay) and unimproved grassland, which refers to rough grazing. As with croplands, the CSO figures for the years 1990-1996 have been adjusted, as outlined in section 7.10. At the time of writing this NIR, the CSO had not released a complete set of figures for grassland areas for 2005. The missing data has been estimated based on an extrapolation of the 2003 and 2004 figures.

It is important to note that both improved and unimproved grassland areas are estimates of grasslands *in use* for agricultural purposes. Rough grazing areas *in use* are native grasslands that are unmanaged with regard to drainage or other inputs, such as fertilizer, but which may be quite intensively grazed by cattle or sheep. The CSO annual statistics for rough grazing exclude other areas of grassland not reported to be in use for agricultural purposes. These grasslands are assumed to be unmanaged natural grasslands, in a carbon stable state, with no associated emission or sink activity. However, they do represent a reserve of lands available for conversion to other land uses. Given the uncertainty of the area of un-used grassland, it was decided to include this type of grassland in the category *5.F Other Land*. When there is a demand for new grassland for use as rough grazing, it is met by a conversion from *5.F Other Land* to unimproved grassland. Overall, the area of improved pasture has been increasing slightly and the area of rough grazing, or unimproved grassland has been decreasing. This is probably in response to sheep farming policy, which in recent years has sought to decrease over grazing on vulnerable commonage and mountain areas. The grazing of unimproved grasslands leads to degradation of the soil, with consequent emission of carbon.

From the data available, it is difficult to estimate the flux of area within the category *5.C.1 Grassland Remaining Grassland*. The annual CSO figures refer to the areas of land which farmers have declared to be “in use” under the specified types of use. Given the economic investment required to maintain “improved” grassland, it is probable that the declared “in use” areas are a good indicator of the actual extent of well-maintained managed grasslands.

Therefore, significant changes in the improved grassland areas do represent changes in land use, with lands either being neglected, or actively managed, depending on the potential for good economic return. The neglect of improved grasslands will cause the land to revert to the nominally managed or native grassland state over time. The transition to rough grazing causes a degradation of the soil, leading to an emission of carbon. However, it is assumed that the average biomass remains constant. This is an underestimate of the effect of grazing, but insufficient data exists to quantify the impact.

There is a strong dynamic of lands moving between grassland and cropland (with a knock effect on the area assigned to other land). This is because of the nature of the CSO figures, which record only the area given over to each activity in a particular year. Under Irish conditions, conversion of grassland to cropland leads to a net loss of carbon from the soil, and also a loss of living biomass when the Tier I default methods are applied. There is little data on conversion of forest land to grassland. For the purposes of the 2007 submission it is assumed that the amount of deforestation is negligible, especially as there has been an annual increase in forest over recent decades. However, deforestation is an issue to be addressed in future.

7.5.2 Carbon Stock Changes in Grassland

The relevant carbon stock changes are for living biomass under *5.C.2 Land Converted to Grassland* and for soils under both *5.C.1 Grassland Remaining Grassland* and *5.C.2 Land Converted to Grassland*.

Carbon Stock Changes in Living Biomass

The Tier I methodology assumes that grassland remaining grassland has zero biomass carbon stock change under static management practices. This approach is adopted here and the notation NO is entered in CRF Table 5.C. The category *5.C.2.5 Other Land Converted to Grassland* is the most important conversion category in most years while some conversions from cropland and exhausted peatlands also occur. Carbon stock changes are estimated at Tier I in the same way as for land converted to cropland using Equation 7.2 above. The biomass value of cropland converted to grassland is taken to be 10 t/ha and the carbon stock increase due to growth in grasslands (ΔC growth) in the first year is 6 t C/ha from GPG Tables 3.4.2 and 3.4.3. In the case of peatlands there is no initial biomass at the time of conversion to grassland and therefore the carbon stock change is due only to the first year's growth at 6.0 t C/ha. The category *5.C.2.5 Other Land converted to Grassland* is in effect the transition of unmanaged native grassland to improved or unimproved pasture, as indicated in section 7.5.1 above. There is a change in carbon stock associated with conversion to improved grassland, as the land will invariably be subject to ploughing and reseeded. This is accounted for through Equation 7.2 as a loss of 6 t C/ha for standing biomass followed by a gain of 6.0 t C/ha through growth in the first year, using the default values¹.

Carbon Stock Changes in Soils

The distribution of grassland areas converted from other land uses over the IPCC soil classes is determined from GIS analysis of CORINE 1990 land cover data superimposed on the general soil association map. Mineral soils as identified from the general soil map were allocated to the five IPCC soil groups and their organic carbon status is established from the soil type and the default reference soil organic carbon stocks (Table 7.5). Table 7.6 shows the adjustment factors applied to the default SOC_{ref} to correct for land use and farming practice. The principal conversion affecting carbon stock change in soils is that from native

¹ There appears to be some inconsistency between default biomass carbon stocks given in GPG Table 3.4.9 and those derived from GPG Tables 3.4.2 and 3.4.3. The inventory agency believes that the value of 13.6 tonnes DM/ha for the cold wet temperate climate zone should be 12 tonnes DM/ha.

grassland to rough grazing, which causes a decrease in soil carbon. Conversely, it can be seen from Table 7.6 that conversion from cropland to improved grassland implies an increase in the soil carbon. A significant secondary source of carbon emission is the use of wetland soil types as pasture. It is assumed here that the wetlands soils under pasture is to some extent artificially drained, and so encourages the emission of carbon from this organic soil type. The default emission rate of 0.25 t C/ha for drained organic soils under grassland have been applied.

7.5.3 Agricultural Lime Application

Much of the total emission of carbon for productive agricultural land derives from the use of lime applied to control soil acidity. As outlined in section 7.10, agricultural lime application has been revised to allow a disaggregation of associated CO₂ emissions between grassland and cropland. Data on the annual amounts of lime applied to land are currently obtained from the Irish Business and Employers Federation. Limestone is the standard form of the application. The CO₂ emissions are calculated using the default emission factor of 120 kg C/tonne lime. The estimates are reported in CRF Table 5(IV) rather than in CRF Tables 5.B and 5.C, the carbon stock change tables for cropland and grassland, respectively.

7.6 5.D Wetlands

7.6.1 Wetland Areas

The national wetland area was split into two types (Table 7.3). Wetlands as applied to Ireland refer to natural unexploited wetlands while peatlands are those wetland areas drained for the purpose of commercial exploitation and harvesting of peat. This split was necessary to account for the conversion of wetlands to peatland, which would be an internal change under the IPCC definition of wetlands. The activity data areas that appear under category *5.D.1 Wetlands Remaining Wetlands* in CRF Table 5.C therefore refer to peatlands in the Irish context and the conversion to wetland is not applicable.

The commercial exploitation of wetlands as peatlands by Bord na Mona (the Irish Peat Board) according to the land-use definition in Table 7.3 proceeds in three separate stages, all of which may lead to changes in carbon stocks. Drainage is the first management activity, followed after several years by removal of the top layers of plant growth in the first season of peat extraction and then by the industrial extraction and harvesting of a layer of 10 to 15 cm of peat annually. The average working life of commercially developed Irish peatland is of the order of 30-50 years. Conversion to grasslands or forest land has been the historically favoured use of cutaway peatland. However, in recent years wetlands reclamation has been investigated, and achieved with some success. The areas reported under category *5.D.1 Wetlands Remaining Wetlands* refer to all lands drained, whether the peat remains covered by vegetation or is exposed. The company manages its peat reserves to meet present demand and is therefore progressing to extract peat from new sites only when an older field is exhausted. It is assumed that the decrease in reserves of peatland indicate the new extraction areas, and therefore they are an estimate of the area from which biomass has been removed. Until recently, Bord na Mona held a small area of undrained wetlands in reserved. However, these lands have been transferred to the National Parks and Wildlife for conservation.

Bord na Mona supplied the area estimates for the company's commercial activities and for private industrial and domestic harvesting of peat. The data for Bord na Mona commercial peat extraction areas were given as totals for consecutive five-year periods for a variety of

peatland categories (Table 7.7). The average value obtained from this total is used for each of the five years to obtain the full time series. Private industrial exploitation of peatlands is of the order of one-eighth that of the commercial activity of Bord na Mona. As similar harvesting methods are used, the areas have been extrapolated from the Bord na Mona values for individual years. Domestic harvesting of peat bogs by private landowners for their own household use is a strong tradition in many parts of Ireland, and although well documented in a social and cultural context, the amount of such peat extraction is poorly quantified. Current estimates are that approximately 1Mt of peat are extracted each year by hand cutting. This represents a bog area of about 400 hectares per year.

Table 7.7 Area Statistics for Peatlands (ha)

Peatland Category	1985-1990	1991-1995	1996-2000	2001-2005	Vegetation Cover
Active Production Bog	49,715	48,961	46,319	43,761	None
Production Reserve (Drained)	16,250	14,100	12,772	5,930	Heather
Fringe Bog (Undrained)	8,300	8,300	8,300	8,300	Heather dominated Bog Vegetation
Partially Drained	3,090	3,090	3,090	3,090	Typical Bog vegetation
Undrained Intact Bog	4,150	2,508	-	-	Intact Bog vegetation
Cutaway Areas					
Forestry (Plantation)	2,500	4,000	4,000	4,200	Conifers
Forestry (Natural)	-	100	800	2,235	Birch / Willow
Wetland (Acidic)	483	483	2,703	9,044	Eriophorum, Carex, Sphagnum
Wetland (Alkaline)	250	1,250	2,150	3,200	Typha, Phragmites, Open water
Lands Sold/Transferred	2,541	1,946	2,658	374	
Total owned (at end of period)	84,738	82,792	80,134	79,760	

7.6.2 Carbon Stock Changes in Wetland

Biomass

Carbon stock changes in biomass are determined by the balance between carbon loss due to the removal of vegetation on preparation for peat harvesting and gain on areas of peatland restoration. These changes have been estimated on the basis that the entire cover of vegetation is removed to prepare for peat harvesting and that an equivalent amount of biomass is returned on restoration of cutaway areas. In NIR 2006, it was assumed the restoration of biomass occurred in the year of conversion. As outlined in section 7.2.1, discussion with experts from Bord na Mona suggest a more appropriate biomass transition period of 5 years.

The area from which vegetation is removed is given by the amount of reserve that enters production annually and the restoration area is taken as the annual increase in cutaway wetland from Table 7.7. The vegetation is typically heather-dominated bog or heathland cover for which a biomass carbon content of 3 t C/ha is adopted (Cruickshank et al, 2000).

Soils

The CO₂ emissions associated with the combustion of peat are accounted for in the *Energy* sector. An additional loss of carbon is associated with drainage and the exposure of the new peat surface annually after harvesting takes place. The annual activity data are the active production areas of Bord na Mona bog together with the areas of peatland in use by private commercial enterprises and by domestic users. All such peatlands are nutrient poor raised bogs or rain-fed blanket bogs for which the appropriate carbon emission factor is 0.2 t C/ha given for boreal and temperate climatic regions in the IPCC good practice guidance.

7.7 5.E Settlements

7.7.1 Areas of Settlements

The area of settlements in 1990 is that given by CORINE 1990. Land converted to settlements is the area taken up by new road building, available from the National Roads Authority, and the area covered by new residential, commercial and industrial construction based on CSO annual statistics, which are extracted from floor area records for permitted development. An incomplete time series of housing types (for the years 1995-2000) was used to estimate the residential building footprint from floor area. It was assumed that approximately 50 percent of the planning permits granted for construction were for green-field sites previously not part of the urban fabric. Section 7.10 outlines a significant revision of the area of land taken by dwellings, which has been incorporated into this submission.

The identification of the land use from which settlement areas are converted is based on an analysis of the distribution of land use classes given by CORINE 1990, with the exclusion of wetland, water bodies, existing continuous urban fabric and other marginal unsuitable land types. Conversions of the different types of land area to settlement areas are assumed to occur in the proportions under which the respective categories existed in 1990. For example, as 80 percent of the land is grassland, it is reasonable to assume that 80 percent of new buildings and road construction takes place on grasslands.

7.7.2 Carbon Stock Changes in Settlements

The assumption is made of complete removal of biomass in the year of the planning permission for buildings constructed or in the year of completion of road projects. The biomass loss from grassland and cropland is as per guidelines using Tier I. The relative loss of biomass from forest per hectare is large. Based on the carbon estimates in Section 7.3, the average biomass of forested lands in Ireland is of the order of 42 t C/ha. No account has been made of the potential increased carbon stock in biomass in urban areas. This may be a significant carbon sink, especially when one considers the policy of actively encouraging urban tree planting along new roads and in new housing developments, but no data are available.

7.8 5.F Other Land

7.8.1 Areas of Other Land

The category *5.F Other Land* includes all lands not classified under the categories 5.A through 5.E. It represents the difference between the sum for categories 5.A through 5.E and the total land area. A large part of *5.F Other Land* is not relevant in terms of potential for emissions or removals but for Ireland this category includes areas of natural grassland which are an available reserve for rough grazing but which are not grazed in the inventory year. As indicated above in section 7.5.1, when the demand for areas of pasture in a particular year is less than in the previous year, surplus areas of improved or unimproved pasture are allowed to revert to rough grazing, which are then not recorded as “in use” by CSO statistics. For area accounting purposes, such lands are assumed to be in transition and are assigned to category *5.F.2.3 Grassland Converted to Other Land* in a manner that maximises the area in *5.F.1 Other Land Remaining Other Land*.

7.8.2 Carbon Stock Changes in Other Land

The degradation of lands reverting to rough grazing not in use results in carbon losses from the soils. The soil classes are identified for *5.F.2.3 Grassland Converted to Other Land* in the same way as for other land-use categories. For mineral soils, SOC_{ref} is assigned according to Table 7.4 while Table 7.5 is used to apply the SOC adjustment factors and the carbon stock change is calculated using Equation 7.1. The default emission factor of 0.25 t C/ha is used to calculate carbon loss from organic soils.

7.9 Uncertainties in LULUCF

Detailed land-use datasets extending over a considerable time period are required in order to apply even the most basic Tier I methods of the IPCC good practice guidance to estimate emissions and removals of greenhouse gases in the land-based approach for the LULUCF sector. The analysis for the several land-use categories invariably means that datasets differing in terms of format, spatial resolution, reference years and other attributes need to be combined for national coverage of sources and sinks. It follows that a high degree of uncertainty is associated with the land area activity data in general. This is especially true of the conversion categories, which are in many cases the land-use categories having the greatest impact on carbon pools. Large uncertainties are also inherent in the parameters that determine carbon stock change factors and in the emission factors for N_2O as indicated by the wide error ranges given in the good practice guidance. It is also difficult to assess to what extent the given values for broad climatic regions are relevant to Irish circumstances. It may be concluded that the uncertainties in reported emissions and removals are large for the sector overall and their full evaluation in quantitative terms has not been possible for the current submission.

7.10 Recalculations in LULUCF

A number of revisions and updates, as listed below, have been made for the 2007 submission, which are described in the following paragraphs.

- a. Adjustment of CSO statistics, affecting 5.B Cropland and 5.C Grassland;
- b. Review of reported total land area of the Republic of Ireland;
- c. Revision of Forest Land and afforestation areas on organic soils;
- d. Revision of areas of Wetlands and Peatlands;
- e. Revised Approach for Agricultural Lime Application;
- f. Revision of 5.C.1 Land Converted to Grasslands;
- g. Revision of 5.E.2 Land Converted to Settlements

(a) Adjustment of CSO statistics, affecting 5.B Cropland and 5.C Grassland

A review of the Central Statistics Office (CSO) data for areas under agricultural use identified an apparent mismatch between total farmed area reported for the period 1990-1996 and that for the period 1997-2004 (Figure 7.1) The total net increase in area reported in agricultural usage in 1997 was approximately 89,000 ha. The increase was shared across most crops and grassland types and was contrary to the contemporary trends in the agricultural sector.

There was no change in the data compilation strategy or analysis methodology employed by the CSO at this time that would explain this increase.

It is difficult to assign a single cause to the reported increase in agricultural area, however a number of developments in methods of farm subsidy payments and administration at this time may have encouraged certain farmers to revise the area of utilised agricultural land that they reported to the CSO. The assumption made in this analysis is that the additional lands reported as utilised agricultural land reflect a change in perception by the farmers of what constituted utilised lands rather than an actual change in land management. In effect, the additional lands were always available, and were already managed as agricultural lands. The CSO statistics available since 1997 are interpreted as a more complete record of agricultural lands, with the proviso that there is some agricultural grade lands unutilised and therefore unreported. In the 2006 analysis of LULUCF, these areas of agricultural land were included in the “Other Land” category up to 1997. This was inappropriate as the lands are impacted by agricultural activities, albeit on an irregular basis. It is therefore appropriate to adjust the CSO data for the years up to 1997 to include these less utilised agricultural lands.




Figure 7.1: Estimated area of Agricultural Land in Ireland 1991-2005

The increase in farmed area reported for 2007 was reported against a background of a downward trend reported in years up to 1997 and again in the years afterwards (Figure 7.1). The average annual decrease in total farmed area from 1991-1996 was 20,000 ha (+/- 18,000 ha), or 16,800 ha annually from a linear fit trend. If farmers had used the older perception of land utilisation, the reported area for 1997 would have been of the order of 107,000 ha lower than given by CSO, showing a decrease of 16,000 ha from 1996. This represents an adjustment to reported areas of approximately 2.5 percent. This adjustment factor is applied *pro rata* backwards to 1990, as shown on Figure 7.1. The adjustment is provisional on the findings of future research analysis of the survey data collated at the time. Another effect of the adjustment is to create a greater continuity between area of land converted to forest lands and settlements and that lost from wetlands (and peatlands) and agricultural lands.

(b) Review of total land area of the Republic of Ireland

The total area of land and fresh water bodies as determined by the Ordnance Survey of Ireland (OSI) is 7.028 million ha. The total land area of 7.118 million ha used in the NIR 2006 was based on CORINE analysis for all land, fresh water and coastal zones. Upon closer examination, the discrepancy between the two area estimates was found to be entirely consistent with the inclusion of coastal zones at or below mean sea level (salt marshes, inter-tidal zones etc.) within CORINE. It was decided to continue using the CORINE value for total area so as to be able to account for land use in coastal areas, and possible land loss, in future analysis.

An analysis of the 90m digital terrain data for Ireland from the Shuttle Radar Topography Mission, SRTM-90, confirms the land area above sea level to be consistent with the OSI value, with no correction required, on a national scale, to account for terrain effects.

(c) Revision of Forest Land and Afforestation on Organic soils

The area and carbon loss due to plantation on organic soils have been updated to include those afforested areas which have been identified as a wetland soil type by IPCC analysis, but which CORINE identifies as another land cover type (usually grassland). In the previous analysis, these areas were categorised as mineral soils. The change is minor, affecting 0.75 percent of private afforestation areas (approximately 1,450 ha since 1990), and 0.6 percent of Coillte plantations (approximately 320 ha since 1990).

In the updated analysis, these lands are processed using the mineral soil methodology, but with a transition period of 4 years. The change in soil organic carbon is based on the default IPCC values for wetland soils, with the exception of afforestation areas on unimproved grasslands, which is treated in the same way as plantations on organic soils. The default emission from unimproved grassland on wetland soils is zero, which is inconsistent with recent research from Hargreaves (2004). This treatment has a knock-on effect in the area of lands lost from other land use categories. Afforested lands are now assumed to derive from Wetlands, rather than from Grasslands or Other Land.

A second update to in respect of Forest Land is the explicit inclusion of deforestation due to conversion of forest land to settlements. This manifests as a modest decrease in the area of category 5.A.1 Forest Land Remaining Forest Land. The biomass carbon loss is accounted for in 5.E Settlements, and not category 5.A. Care has been taken to avoid “double counting” of this carbon loss.

(d) Revision of areas of Wetlands and Peatlands

As noted above there is an additional decrease in wetlands areas lost to forestry due to the inclusion of those areas of afforestation on wetland soils as classified by soil type rather than by Corine land cover. Additional information has arisen as regards the typical period of recovery of biomass to wetland restoration areas. In the 2006 analysis it was assumed that full biomass was restored in the first year of planned restoration. Bord Na Mona (McNally, personal communication) suggest that a recovery period of 5 years is more typical. Therefore, the biomass storage on restored peatlands is assumed to be in transition for 5 years, with a linear growth rate of 0.6 t C/ha, as opposed to full restoration of 3.0 t C/ha in one year (Tomlinson, 2005). The effect of this change has been to smooth the stepwise shape of the carbon emissions curve for this sector. However, the increased rate at which drained wetland was brought into peat production in the period 2000-2005 has become more evident, with an associated increase in carbon emissions

(e) Revised Approach for Agricultural Lime Application

In the NIR 2006 no attempt was made to apportion emissions due to lime application between improved grassland and croplands. In the 2007 estimates, lime-spreading emissions have been divided in proportion to relative land areas between improved grassland and croplands. This equates to 10 percent of emissions from lime application being assigned to croplands and 90 percent to grasslands. It is assumed that no lime spreading occurs on unimproved grasslands. Some agriculture experts suggest that arable lands are much more intensive users of lime than grass based systems (Coulter, 2004), and therefore a greater proportion of lime usage should be attributed to cropland. However, more research is required to address this question fully.

(f) Revision to 5.C.1 Land Converted to Grasslands

The area of grasslands in transition from croplands was found to be underestimated in the 2006 analysis. In a given year, if there is a net decrease in the area of cropland, it is assumed that this prime agricultural land goes to improved grassland, with adjustments to account for lands being converted to Forest Land and Settlements. The default methodology suggests that lands should be considered “in transition” for a period of 20 years, during which time the soil tends towards an equilibrium state of carbon exchange to the atmosphere. Therefore, the applicable area of soil in transition to grasslands from croplands is the sum of areas converted over the previous 20 years, with the assumption (in this case) that no converted land reverts to cropland during this period.

In the case of cropland conversion to grassland this summation was omitted in error. Hence soils were considered in transition only for the year during which the conversion took place. The corrected estimates give an additional emission of carbon from the soil of approximately 275Gg C over the period 1990-2005.

The conversion of Other Land to Grassland is treated differently. It is assumed that Other Lands hold a reserve of natural, unimproved grassland, which are exploited for agricultural purposes as the need arises, and “abandoned” (i.e. not used in the inventory year) when there is surplus grassland. It is assumed that land parcels of this type are unsuited to exploitation for more than one year at a time, and therefore the carbon emission associated with degradation due to grazing animals occurs only in the year of conversion. Therefore a default 20-year transition period is not applied in the case of Other Land conversion to grassland.

Table 7.8 Mechanism for the calculation of area of land converted to settlements

Area of land converted to Settlement: Factors for dwellings (greenfield sites)					
	No. of Storeys	Adjustment for drives	Proportion requiring new access roads	Area of new road	Proportion of buildings on Greenfield sites
	A	D	R	nr	G
Bungalow	1	1.2	0 ¹	10*30 ²	1
Detached	2	1.15	0.5	10*25	0.75
Semi-detached	2	1.1	1	10*15	0.5
Terrace	2	1	0.5	10*10	0.25
Apartments	Urban				0.0
All values in this table are estimates.					
1: It is assumed bungalows are built alongside existing road networks					
2: Based in An Taisce estimate a loss of 30m of hedgerow for access to one off housing.					

(g) Revision of 5.E.2 Land Converted to Settlements

The 2006 estimate of the area of land converted to settlement was based on an estimate of building footprint. However, substantial additional areas of land will be required for new road and driveway access to these buildings. In the 2007 analysis, a mechanism has been developed to introduce these factors into the estimation of total area taken, as shown in Table 7.8.

$$Land\ to\ Settlement_{dwelling} = \sum_{dwelling\ type} N * G * \left(\frac{D * Floor\ Area}{S} + R * nr \right)$$

where N is the number of dwellings built in each dwelling category, D is a factor to allow for additional driveway and other paved areas, S is the number of storeys typical of each housing type, R is the proportion of housing of each type that might require new access roads, G is the proportion of house building on greenfield sites. The values shown in the table are speculative, but do give an indication of the additional land area required for housing beyond the building footprint (Figure 7.2).

The average floor area of new dwellings has increased over the last decade. All houses, except bungalows are assumed to be two storey. New access roads are assumed to be 10m wide, inclusive of pavement. The length of new road associated with each dwelling type is estimated based on the relative frontage of each house type. Additional information on the average floor area of “one-off “ and multi-unit developments is available from the CSO for 2003-2005 (“Construction and Housing in Ireland”, CSO Prn A6/1233). One-off housing is assumed to include both bungalow and detached houses.

The trends for increasing dwelling floor area seen in the 2003-2005 figures are consistent with the aggregate figures available for the period 1990-2005. Therefore an estimate of the average floor area for each broad type of dwelling can be extrapolated backwards to 1990, relative to the overall average floor area for all houses in that year. There has been strong growth in the land converted to settlement under all dwelling types except terraces (estimated for 1990-1994). There has been a very dramatic increase in semi-detached dwelling construction since 2001. This is due to a combination of increasing numbers and increasing floor area. The updated analysis estimates that approximately 12,600 ha (1990-2004), has been converted to settlement area, with complete soil sealing. In the previous analysis (NIR 2006) the estimate from 1990-2004 was c10,100 ha. An additional 1,800ha of new settlement is estimated to have occurred in 2005 (Figure 7.3).




Figure 7.2: Time Series of Land Area converted for Housing




Figure 7.3: Time Series of Land Area converted to Settlements

7.11 Improvements in LULUCF

The coverage of sources of emissions and removals by Ireland in LULUCF is complete for the years 1990-2005. Even though a rather simplified approach has had to be followed due to the level of information available, the assessment of emissions and removals according to the reporting requirements of Decision 13/CP.9 has identified a number of important CO₂ emission sources, in addition to the well known carbon sink in forests. Extensive further work has been conducted to improve data treatment for this submission. The inventory agency is continuing to collaborate with the bodies from which the key land-use and forestry datasets are obtained and has established formal arrangements for the provision of the data within the national system, in the same way as for other sectors. The agency's capacity on GIS is being expanded so that it may devote more resources to integrating the available datasets and to undertaking the level of spatial analysis needed to improve the land area activity data.

The results of an extensive new forest inventory are now available for use in the LULUCF inventory. COFORD is funding a new series of projects on climate change and forestry over the period from 2007 to 2012. The CARBWARE project will develop forest carbon stock change reporting tools and software, building on work begun in 2000, which used FIPS and Forest Service planting data. It will also access data from the new CARBiFOR II and other research projects, to refine estimates of carbon stock change for reporting purposes. Data from IFORIS and NFI will be needed to fulfil reporting requirements. An initial data provision and analysis framework has been developed for CARBWARE, which is being extensively enhanced to make available an integrated system that will meet the reporting needs of the Convention and the Kyoto Protocol.

Work on developing a single forest cover and attribute data set has been progressing in the Forest Service over the past two years. Most of the data set has been compiled, apart from a subset of grant and premium data that needs to have species attributes input manually. In its final version the full data set will include location, planting year, species area and open space area attributes, for all forest greater than 0.5 ha in area (with the post 1990 afforestation data for areas down to 0.1 ha). The Forest Service will have a system in place for access to and use of the data.

Chapter Eight

Waste

8.1 Overview of Waste Sector

The main activities giving rise to greenhouse gas emissions in the *Waste* sector are solid waste disposal in landfill sites, wastewater treatment and waste incineration (Table 8.1). The most important of these sources is solid waste disposal where CH₄ is the gas concerned. Landfills represent a key emission category in Ireland (Chapter Three) and the emission estimates are reasonably well quantified in current inventories. The treatment of wastewaters and sludge in anaerobic systems may also be an important source of CH₄. All wastewater treatment in Ireland is aerobic and no CH₄ emissions are produced. However, estimates of CH₄ emissions from sludge treatment are included. The N₂O emissions arising from the production of human sewage continue to be reported following the inclusion of first estimates for this source as part of the recalculations undertaken for the 2002 submission.

Unlike many other developed countries, Ireland has not used waste incineration as a waste management option to any significant extent to date. No incineration of municipal waste currently takes place and the burning of hospital wastes was discontinued around 1995. The practice is now mainly confined to the destruction of liquid vapours by a small number of chemical and pharmaceutical companies. The quantities of both greenhouse gases and indirect gases concerned may be negligible. The incineration of municipal waste will become an additional source of emissions for inclusion in annual inventories in the coming years following the grant of IPC licences for two incinerators by the EPA recently.

Table 8.1. Level 3 Category and Gas Coverage for Waste

Waste	CO ₂	CH ₄	N ₂ O
A. Solid Waste Disposal on Land			
1. Managed Waste Disposal on Land	NA	All	NA
2. Unmanaged Waste Disposal Sites	NA	All	NA
3. Other	NO	NO	NO
B. Wastewater Handling			
1. Industrial Wastewater	NA	All	NE
2. Domestic and Commercial Wastewater	NA	All	All
3. Other	NO	NO	NO
C. Waste Incineration	NE	NE	NE
D. Other	NO	NO	NO

All : all emission sources covered; NE : emissions not estimated; NO : activity not occurring; NA : not applicable (no emissions of the gas occur in the source category); IE : emissions included elsewhere

8.2 Solid Waste Disposal (SWD)

The anaerobic decomposition of organic matter in solid waste disposal sites (SWDS) is a major source of CH₄ in developed countries. The CH₄ production potential of SWDS in a particular year depends on the cumulative solid waste disposal over many previous years, the composition of the wastes and the level of management applied to the disposal sites concerned. Well managed deep landfills in which the wastes receive constant compaction and cover material have a much greater capacity for CH₄ production than shallow unmanaged sites or open dumps where aerobic conditions may dominate. Methane production within landfills occurs in a number of distinct phases with virtually all CH₄ usually being realised within a period of approximately 20 years.

8.2.1 Methodology for CH₄ Emissions from Solid Waste

The development of a national waste management strategy for Ireland (DELG, 1998) recognised the need for comprehensive analysis of the CH₄ production potential of landfills, particularly in view of the need to reduce the amount of municipal waste being placed in landfills. A modified form of the IPCC Tier 2 method was therefore adopted as the most appropriate basis on which to assess annual CH₄ emissions where reasonable predictions could be made for decreasing waste quantities into the future. The results obtained from this revised methodology were included as an important component of the recalculations reported in the 2002 submission. More in-depth analysis of the historical time series of solid waste disposal was undertaken in estimating the 2001 emissions from this source, necessitating further revision of the previous estimates for the years 1990 to 2000. The revised estimates were submitted in 2003. The same basic method continues to be used but further development and minor updating of the underlying activity data remains part of the ongoing work and assessment in relation to this source of CH₄ emissions. In this regard, sewage sludge placed in landfills was taken into account as an additional source of degradable organic matter for the estimation of emissions in 2004.

The approach underlying the quantification of CH₄ from solid waste disposal uses the relationship given in Figure 8.1 to describe the CH₄ production from all contributing solid waste deposited in landfills in a particular year. This relationship is based on a two-stage first-order model (Cossu et al, 1996) for landfill gas production, incorporating a lag period of one year before CH₄ generation commences, followed by active CH₄ production over 20 years. The estimates take account of a variable allocation of wastes between well-managed landfills, where the full CH₄ potential is realised, and shallow unmanaged landfills for which 40 percent of the potential CH₄ is assumed to be emitted. To estimate annual emissions for the years 1990 to 2005, the CH₄ potential of wastes landfilled in each year from 1969 (21 years prior to 1990) is first determined. These annual CH₄ potentials are then assigned as emissions over 20 subsequent years (with an initial lag of 1 year) according to the proportions depicted in Figure 8.1 and their cumulative contributions for the 20 year period give the total emissions for the end year in that period.

8.2.2 CH₄ Production Potential of Solid Wastes

The CH₄ production potential of solid wastes is determined by the amount of degradable organic carbon (DOC) in wastes, which in turn depends on the amount and composition of the waste material. The IPCC Guidelines use municipal solid waste (MSW), which usually refers to household and commercial refuse, as the basic parameter from which the amount of DOC is established for the purposes of estimating CH₄ potential. However, it is recognised that some industrial wastes, sewage sludge and street cleansings may also contribute to degradable organic matter in landfills and therefore they should be taken into account to the extent possible.

The EPA commenced the development of the National Waste Database in the early 1990s to address a severe lack of information on waste production and waste management practices in Ireland. The database is needed to support radical reform of national policy and legislation on waste pursuant to the Waste Management Act of 1996 and subsequent Government strategies on sustainable development (DELG, 1997) and waste management (DELG, 1998). National statistics generated from this database published on a three-year cycle and interim reports published on a yearly basis since 2001 by the EPA, are the primary basis for establishing the historical time-series of MSW placed in landfills in Ireland for the purpose of estimating CH₄ emissions from this source. These publications provide detailed descriptions of the methods employed to compile the waste database. The results of other less comprehensive surveys undertaken in previous years (Boyle, 1987, ERL, 1993, MCOS, 1994 and DOE, 1994) have also been used to some extent in compiling the MSW time-series.




Figure 8.1. Typical CH₄ Production Pattern in Solid Waste Disposal Sites

The National Waste Database reports for 1995 (Carey et al, 1996), 1998 (Crowe et al, 2000), 2001 (Meaney et al, 2003), and 2004 (Collins et al, 2005) and interim reports for 2002 (Collins et al, 2004a), 2003 (Collins et al, 2004b), 2005 (le Bolloch et al, 2006) were used to establish the MSW time-series up to 2005 and the corresponding annual CH₄ emission estimates. The time-series estimates given in the present submission also account for the inclusion of sewage sludge and are fully consistent over the period 1990-2005. The historical time series of wastes placed in solid waste disposal sites up to 2005, along with their associated DOC contents, used as the basis of CH₄ emission estimates from this source are provided in Table F.1 of Annex F. The following paragraphs describe the steps and assumptions made in developing these data from the available National Waste Database statistics:

- the waste material contributing to DOC includes MSW (household and commercial refuse) and street cleansings, as given in the National Waste Database reports together with sludge from municipal wastewater treatment that are deposited in landfills;
- the per-capita MSW generation rates indicated for 2004, 2001, 1998 and 1995, along with those implied by the earlier surveys, are used to assign the rate of MSW production in all years;

- similarly, the proportion of MSW that is placed in landfills in 1995, 1998, 2001, 2002, 2003, 2004 and 2005 is used to assign the corresponding value in other years;
- the per-capita MSW generation rate and the proportion of MSW that is placed in landfills are assumed to remain constant at 1 kg/cap/day and 75 percent, respectively prior to 1985;
- the amount of street cleansings is estimated from the ratio of street cleansings to MSW given by the 1995, 1998, 2001, 2002, 2003, 2004 and 2005 data;
- the waste constituents of MSW that contribute to DOC are organics, paper, textiles and the category other (fine elements, unclassified materials and wood wastes), as identified in the available breakdown for 1995, 1998, 2001, 2002, 2003, 2004 and 2005
- the IPCC default proportions of DOC are used for organics, paper and textiles (15, 40 and 40 percent, respectively);
- DOC contents of 25 percent and 15 percent have been assumed for street cleansings and the category other, respectively;
- the DOC contribution of sludge is determined from information on the BOD content, the BOD removal rate and the proportion of sludge disposed to landfill.

The potential CH₄ available from the annual DOC in SWDS, determined as described above, is estimated as follows;

- in accordance with the IPCC good practice guidance, 60 per cent of the total available DOC in solid waste is dissimilated on an equi-molar basis to CH₄ and CO₂;
- in the period 1990-1995, 60 percent of DOC is assigned a methane correction factor (MCF) of 1, on the basis that the MSW from all major population centres (60 percent of the population) is deposited in managed landfills (Carey et al, 1996) and the full CH₄ potential is ultimately realised;
- in the period 1990-1995, 40 percent of DOC is assigned a MCF of 0.4, on the assumption that 40 percent of MSW is placed in unmanaged SWDS of less than 5 m depth;

the MSW split between managed and unmanaged sites in 1969 is taken to be the reverse of that adopted for the years 1990-1995 and appropriate adjustment is made for the intervening years and for the years after 1995 to reflect a gradual increase for managed landfills. The MSW split adopted for 2005 is 0.95 for managed sites and 0.05 for unmanaged sites. This represents a major change from the 0.67 : 0.33 division used for 2004 and it has been made following discussions with waste experts who believe that almost all landfills in Ireland could be classified in the managed category as used by IPCC.

8.2.3 CH₄ Emissions from Solid Waste in 2005

The final estimates of CH₄ emissions from the IPCC source-category *6.1 Solid Waste Disposal on Land* are derived for the years 1990-2005 from the time-series data on CH₄ potential given in Table F.2 of Annex F, using the time-dependent rate of release shown in Figure 8.1. The emissions in a particular year are simply the cumulative contribution for that

year arising from managed landfills and from unmanaged landfills separately over the period of 21 years that ends in the year concerned.

Landfill gas has been recovered at a small number of landfill sites in Ireland since 1996. The amount of CH₄ utilised for electricity production at these sites is known from annual reports on renewable energy use. In this top-down analysis, the amount of CH₄ captured for energy use is estimated from the reported electricity production from this source in the national energy balance, assuming assigned percentage conversion efficiency factors. As part of the implementation of Directive 1999/31/EC (CEU, 1999) and reporting to the European Pollution Emissions Register (EPER), the EPA made estimates of CH₄ emissions from 65 individual landfills in Ireland for 2004 using either the LANDGEM model developed by the USEPA or the GASSIM model used in the UK. These were considered to be all the landfills that were producing CH₄ in any appreciable quantities in that year and the exercise was a repeat of that undertaken for 2001. The landfills identified which burn-off or flare CH₄ produced in 2004 were contacted by inspectors in early 2007 to provide data for methane flaring for 2005. The total CH₄ flared was then added to the total utilised for energy production to derive a total quantity of CH₄ that must be deducted from CH₄ production to obtain the emission estimates.

Figure 8.2 shows the trend in emissions from solid waste disposal over the period 1990-2005 achieved from the analysis described above and using the results of flaring and utilisation of landfill gas for the years 1996 to 2005. The revised estimates indicate CH₄ emissions of 63.431 Gg in 1990 from an annual average of 0.94 million tonnes of contributing municipal waste over the preceding period of 20 years. In 2005, emissions had increased to 77.056 Gg after recovery and flaring and the average contributing wastes were approximately 1.6 million tonnes annually.

It is evident from Figure 8.2 that emissions from solid waste disposal are increasing significantly even though a substantial part of CH₄ production is offset by flaring and utilisation. The implied emission factors for CH₄ production are typically 0.1 to 0.11 tCH₄/t MSW for managed landfills. These values may not be comparable to those for other Parties as the activity data reported for a particular year is the 20-year average MSW for the period contributing to emissions in that year. This gives an IEF approximately 30 percent higher than those based on MSW for the year concerned, e.g. range 0.066-0.080 compared to 0.101-0.115 over the period 1990-2005. Ireland has no waste incineration and a high proportion of waste consequently goes to landfill.




Figure 8.2. Methane Emissions from Solid Waste Disposal Sites 1990-2005

8.3 Emissions from Wastewater Handling

8.3.1. CH₄ Emissions from Wastewater and Sludge

It is assumed that in Ireland all wastewaters sent to municipal wastewater treatment plants are treated aerobically and that all wastewater treatment plants on-site of industries also operate aerobically. A third of the population, mostly found in rural areas, treat their wastewater using septic tanks (Smith et al, 2003). The temperature experienced in a septic tank is not optimum for methane-producing bacteria, therefore there is negligible CH₄ production. The wastewater treated by the septic tank is released to a percolation area or to surface waters. The commercial wastewater that does not go to municipal wastewater treatment plants is assumed to be treated aerobically on site before discharge to percolation areas or surface waters and again there are no CH₄ emissions. Consequently, there are no CH₄ emissions produced from wastewater handling and the notation key NO is used under sub-categories 6.B.1(a) and 6.B.2(a) of CRF Table 6.B.

National studies (O'Leary and Carty, 1998) indicate that 3 percent of sludge produced in both industrial wastewater and domestic and commercial wastewater handling, including septic tanks, is treated anaerobically. A portion of this sludge is spread on agricultural lands (12 percent in 1990 and 35 percent in 2005) where it contributes to the N₂O emissions from soils, as described in section 6.5.1. Some of the sludge is also sent to landfill (42 percent in 1990 and 31 percent in 2005) and is accounted for in the estimate of degradable organic carbon that generates CH₄ in landfills as outlined in section 8.2.1 of this chapter. The estimates of CH₄ emissions from wastewater and sludge are derived using the national statistics, country specific values and default values from the IPCC Guidelines listed in Table 8.2.

8.3.2 N₂O Emissions from Human Sewage

Emissions of N₂O from human sewage discharges reported under source category 6.B Wastewater Handling have been made following the IPCC methodology. This source of emissions was first included as part of the recalculation exercise undertaken for the 2002 submission and continues to be part of the inventory. In previous submissions, the body

weight and average protein intake of the population were taken as 80 kg and 0.75 g/kg body weight per day, respectively, to estimate annual protein consumption based on information provided by the Food Safety Authority of Ireland (FSAI, 1999). The 2003 in-country review of Ireland's 2003 submission identified that FAO statistics indicate a typical protein intake of about 114 g/capita/day for the population of Ireland, compared to the 60 g/capita/day suggested by the FSAI recommendations. Ireland adopted the FAO estimate of protein intake in the estimates for 2003 and the corresponding emissions in other years were recalculated on this basis for the purpose of the 2005 submission. The emissions in 2005 are estimated in the same way. The N₂O emissions are computed by taking the IPCC default proportion of 0.16 for the nitrogen content in protein and applying the default emission factor of 0.01 to obtain the quantity of nitrogen in sewage ultimately entering the atmosphere as N₂O.

Table 8.2 Parameters for Estimation of CH₄ Emissions from Wastewater and Sludge 2005

Parameter	Source	Value
Population Equivalent	National Statistic (Incl. Dom/Commercial and Some Industry)	6118.7
Kg BOD/1000 people/yr of wastewater	IPCC Guidelines; 0.06*1000*365	21,900
Total Organic Waste from Dom/Comm/Some Industry (Gg Dc/yr)	21900*PE (National Statistic)	163.85
Total amount of sludge (Tonnes of DS)	National Statistic for some years	43,912
Total Wet Sludge (Gg Dc/yr)/(20% solids)	Tonnes of DS * 100/20	219,559
Fraction of Sludge (wet weight) comes off Wastewater	Total Wet Sludge (M ³) /Total Wastewater (M ³)	0.0005
Bo Maximum Methane Producing Potential CH ₄ kg/kgBOD	Default IPCC	0.6
SBF is the Fraction of BOD that readily settles	Country Specific	0.4
Organic Content of Industrial Sludge (BOD kg/m ³)	0.4 of Ave.BOD of Treated Ind. WW- 150kg/m ³	60
Fraction of BOD in sludge degrades anaerobically (FTA)	Country Specific	3%
Emission Factor for Sludge (3% Anaerobic)	0.03*0.6	0.018

8.4 Recalculations in Waste

Recalculated estimates of emissions in the Waste sector are due to the following inventory revisions carried out in the 2007 reporting cycle for the years 1990-2005

- Improved estimates of paper, organic matter, textiles and other wastes as a proportion of total MSW sent to landfill sites were obtained for the years 2002-2004 inclusive;
- The National Waste Report 2004 (Collins et al, 2005) provided revised estimates of MSW arising for 2002 and 2003 and also data in relation to the total amount of industrial sludge produced in 2004;
- To better reflect the rapid change towards managed landfills the proportions of MSW deposited at managed and unmanaged sites were revised from 2001 onwards with a 0.95 : 0.05 split adopted for 2005;
- Revised landfill gas utilisation figures for energy production were supplied as part of the national energy balance. These revised estimates apply to the years 1996-2004 inclusive.

The combined effects of these revisions on estimates of emissions in the waste sector is quite small and they affect the years 2002-2004 only. The estimates for 2002 increased from 70.64 Gg CH₄ to 71.28 Gg CH₄ while the estimates decreased from 78.73 Gg CH₄ to 78.07 Gg CH₄ in 2003 and from 79.91 Gg CH₄ to 77.90 Gg CH₄ in 2004. The net effect of the recalculations outlined above on the inventory is presented in Table 8.3.

8.5 Improvements in Waste

Radical changes in waste management have taken place in Ireland through the EPA waste licensing system and Government initiatives designed to implement waste management policies that favour prevention, minimisation and recycling options at the expense of waste disposal on land. Many landfills have been closed down and new landfills operate only to

best practice in terms of management and pollution control and in their capacity to track waste streams, in accordance with Directive 1999/31/EC (CEU, 1999).

These changes have major implications for the evolution of the time-series of municipal solid waste as it is applied to the estimation of CH₄ emissions from landfill sites. Major assumptions and generalisations are inherent to the determination of these emissions using the approach described above that has been used now for many years. Some of these assumptions may need to be modified from time to time, as in the case of the split between managed and unmanaged landfills applied in this reporting cycle. The huge variety of landfill sites that is represented by calculations at the national scale and the lack of good historical data for the extended period that must be taken into account mean that the emissions baseline relative to this new waste management regime is already highly uncertain. It is difficult for the methodology to adequately reflect major changes relating to landfills in a robust and transparent manner that maintains consistency in the emissions time series.

In addition, further assessment of landfill gas production is needed on an individual site basis to compare results from top-down and bottom-up approaches currently being used. To this end, the inventory experts will continue to collaborate with colleagues in the EPA using available models to estimate landfill gas emissions for individual sites as part of bottom-up emissions reporting related to Directive 1999/31/EC and EPER.

Table 8.3. Recalculations for Waste 1996-2000 and 2002-2004

IPCC Source Category		1996	1997	1998	1999	2000	2002	2003	2004
Emissions 2006 Submission (Gg CO₂ eq)									
6.A.1 Managed Waste Disposal on Land	CH ₄	1254.38	1207.59	988.00	889.68	1088.82	1000.92	1117.76	1178.94
6.A.2 Unmanaged Waste Disposal Sites	CH ₄	368.84	376.79	385.50	397.00	413.17	454.47	479.40	498.96
6.B Wastewater Handling	CH ₄	14.73	18.13	20.52	21.31	22.23	23.48	23.85	24.15
6.B.2 Domestic & Commercial Wastewater	N ₂ O	112.44	115.82	118.20	121.56	126.70	127.27	129.27	130.20
Total		1750.40	1718.33	1512.23	1429.55	1650.92	1606.13	1750.28	1832.25
Emissions 2007 Submission (Gg CO₂ eq)									
6.A.1 Managed Waste Disposal on Land	CH ₄	1130.96	919.23	985.23	1009.03	1078.35	1042.48	1169.06	1174.12
6.A.2 Unmanaged Waste Disposal Sites	CH ₄	368.84	376.79	385.50	397.00	413.17	454.47	470.47	461.87
6.B Wastewater Handling	CH ₄	14.73	18.13	20.52	21.31	22.23	23.48	23.85	24.69
6.B.2 Domestic & Commercial Wastewater	N ₂ O	112.44	115.82	118.20	121.56	126.70	127.27	129.27	131.38
Total		1626.97	1429.97	1509.45	1548.90	1640.45	1647.69	1792.65	1792.06
% Change from Recalculation									
6.A.1 Managed Waste Disposal on Land	CH ₄	-9.84	-23.88	-0.28	13.41	-0.96	4.15	4.59	-0.41
6.A.2 Unmanaged Waste Disposal Sites	CH ₄	NO	NO	NO	NO	NO	NO	-1.86	-7.43
6.B Wastewater Handling	CH ₄	NO	NO	NO	NO	NO	NO	NO	2.24
6.B.2 Domestic & Commercial Wastewater	N ₂ O	NO	NO	NO	NO	NO	NO	NO	0.91
Total		-7.05	-16.78	-0.18	8.35	-0.63	2.59	2.42	-2.19

Chapter Nine

Recalculations and Improvements

9.1 Recalculation Issues

Increasing demands for more complete and more accurate estimates of greenhouse gas emissions means that the methodologies being used are subject to constant revision and refinement as inventory capacity is increased and better data become available. The general improvement in inventories over time may therefore introduce inconsistencies between the emissions estimates for recent years and those for years much earlier in the time-series. Recalculated estimates are often needed to eliminate these inconsistencies and to ensure that the inventories for all years in a time-series are directly comparable with respect to the sources and gases covered and that the methods, activity data and emission factors are applied in a transparent and consistent manner. In this way, the results can be used with greater confidence in identifying trends and in monitoring progress towards the commitments that have been defined with reference to emissions in the base year. The UNFCCC guidelines provide for the reporting of recalculations as part of the annual submissions from Annex I Parties. Justification for the recalculations are required, as well as explanations of the changes that have been made and the numerical values of the original and revised estimates must be compared to show the impact of the changes.

In 2006, the primary reasoning behind recalculations for the 2006 submission was the need to provide the best possible estimates of GHG emissions in the base year, in the knowledge that such estimates would become fixed from 2006 for the purposes of establishing the assigned amount under the Kyoto Protocol. The main focus of the work was therefore the revisions to the estimates of CO₂, CH₄ and N₂O for 1990, which involved a move from Tier 1 to Tier 2 methods in a number of key categories, and updating of the estimates of HFC, PFC and SF₆ for 1995. These revisions were made in parallel with the inventory compilation for the year 2004, which involved a range of additional improvements in methodologies and data across most sectors, along with the inclusion of some minor additional sources. Another key element of the work on recalculations in 2006 was the completion of the inventory for the LULUCF sector in accordance with the requirements of Decision 13/CP.9 for all years. For the present submission, the recalculations largely relate to the years 1990-2003 in order to achieve consistency with 1990 and 2004, particularly in the *Energy* sector. However, some other improvements are necessary, which in some cases affect all years.

9.2 Recalculations in the 1990-2004 Time Series

The foregoing chapters describe recalculations and improvements for the individual Level 1 source sectors of the inventory and they present the corresponding changes in emissions and removals within the sectors. This chapter summarises the recalculations and assesses their effect in relation to total national emissions to record the updates and the available emissions time-series as they appear in the 2007 submission. The information given here corresponds to that in Table 8 of the CRF for each year. The principal changes that result in recalculated estimates for the years 1990-2004 are

- (i) An expansion and updating of the energy balances for the years 1991-2003, consistent with those adopted for 1990 and 2004 in the 2006 submission, which facilitated more detailed treatment of *1.A.2 Manufacturing Industries and Construction* and the inclusion of a minor source under *1.A.1(c) Manufacture of Solid Fuels and Other Energy Industries*;
- (ii) The application of verified 2005 ETS data to update the estimates of process CO₂ emissions from cement and lime production where appropriate for previous years;
- (iii) The updating of UK population statistics used in the derivation of Irish per-capita emission factors for HFC, PFC and SF₆ under *2.F Consumption of Halocarbons and SF₆*.
- (iv) The updating of UK population statistics used in the derivation of Irish per-capita emission factors for sectors *3.A Paint Application* and *3.D Other Use of Solvents*.
- (v) Correction of a calculation error that was found in the 1990-2004 time-series estimates of the total NH₃-N lost, which is used in the calculation of F_{AW} (N input from manure after adjustment for volatilization) in category *4.D Agricultural Soils*;
- (vi) The emission factor for CH₄ emissions for laying hens in *4.B Manure Management* in the 2006 submission was that for table birds, which is now revised;
- (vii) Revisions to various components of the *LULUCF* sector to take account of improved and updated area totals for most land categories;
- (viii) revised treatment of agricultural lime application to account separately for grass and crops;
- (ix) improved estimates of the carbon loss for biomass due to the land use change associated with the construction of dwellings;
- (x) Improved estimates of the proportional breakdown of municipal solid waste into its constituent components and minor revisions to national statistics in *6.A Solid Waste Disposal on Land*;
- (xi) The relative proportions of municipal solid waste deposited at managed and unmanaged solid waste disposal sites was adjusted from 2001 to better reflect the move towards fully managed landfills *6.A Solid Waste Disposal on Land*;
- (xii) Revised landfill gas utilisation figures for energy production were supplied as part of the improvements to the national energy balances. These revised estimates have been applied to *6.A Solid Waste Disposal on Land*.

The original and revised numerical values of the emissions estimates for the years 1990-2004, along with the changes related to methods, activity data and emission factors are detailed in the respective Tables 8(a) and 8(b) of the CRF time-series. The trend in emissions over this period is discussed in Chapter Two.

9.3 Effect of Recalculations in the 1990-2004 Time Series

Tables 9.1 and 9.2 present the changes due to recalculations according to greenhouse gas and IPCC sector. The overall effect is a very minor increase in total emissions in most years. More significant changes are evident for some individual gases and sectors, although the largest proportional changes occur in the case of either sectors or gases that contribute a relatively small percentage of total emissions. The major change for gases in Table 9.1 is the decrease in N₂O emission estimates, which is approximately 3 percent in each year over the

period 1990-2004. This decrease reflects recalculations in *Agriculture* resulting from item (v) above, where they reflect the revised estimates of N₂O emissions from agricultural soils (Table 6.4). This decrease however is somewhat offset by a minor increase in CH₄ emissions as a result of recalculations for the poultry sector. There is an increase of up to 3 percent for CO₂ emissions without LULUCF due to the changes in the energy balances.

The largest sectoral change occurs in the case of *Land Use, Land Use Change and Forestry* where they reflect revised and improved estimates of agricultural land areas, updates to areas of forest land and areas of afforestation in addition to more detailed estimates of the carbon loss for biomass due to the construction of dwellings. The revised estimates for *Waste* over the years 1996–2004 vary from a decrease of approximately 17 percent in 1997 to increases in total emissions of approximately 2 percent in later years.

9.4 Inventory Improvements to Date

Consistent greenhouse gas inventories in Ireland are available for the years 1990 and 2005. The annual inventories are substantially complete with respect to the coverage of the six greenhouse gases and all IPCC source categories. The range of really important greenhouse gas emission sources in Ireland is quite small and most of the important elements of good practice are now taken into account in the current approach to estimating their emissions. In the past, the full implementation of sector-specific good practice guidance was constrained by a lack of resources and the scarcity of activity data and country-specific emission factors, which precluded the use of the recommended high-tier methodologies in several instances. The inventory agency has recently acquired additional staff resources and has been able to apply the results of several important outsourced projects that were targeted at producing the necessary information to apply more appropriate methods and emission factors in these areas during the 2007 reporting cycle. As a consequence, a large amount of work has been undertaken to recalculate previously reported inventories for the years 1990-2004.

Ireland recognises the need to deliver annual submissions in close conformity with the UNFCCC reporting guidelines on annual inventories to facilitate the work of expert review teams in conducting productive and efficient technical reviews of greenhouse gas inventories. Every attempt is made to participate in the UNFCCC review process and to facilitate the work of the UNFCCC secretariat, especially insofar as it impacts on the quality and transparency of the Irish estimates of emissions. The in-country review of Ireland's 2003 submission (UNFCCC, 2003b) was an important development in this regard. The majority of the recommendations in the 2003 review report have now been implemented, following the extensive improvements and recalculations conducted for the 2006 and 2007 submissions. As these improvements cover issues such as the development of an expanded national inventory report in line with the structure specified in the UNFCCC reporting guidelines, the complete coverage of the LULUCF sector according to the requirements of Decision 13/CP.9 and detailed work to ensure full consistency between the NIR information and the CRF tables, they also address the main findings of the more recent centralised reviews in 2004 and 2005. It may be stated therefore that the inventory material being submitted in 2007 broadly meets the principles of transparency, completeness, consistency, comparability and accuracy laid down in the UNFCCC reporting guidelines.

9.5 Inventory Improvements Going Forward

Further general improvements to greenhouse gas inventories during this period are taking place through consolidation and implementation of the national system, which is now established, and through application of formal QA/QC procedures that have been put into operation as an integral part of the system. The consultant's Scoping Report (Thistlethwaite *et al*, 2005) on Ireland's national system concluded that many of the essential aspects of a national system are in place in Ireland and that the principal outstanding requirements were to clarify the roles and responsibilities between EPA and DEHLG and to formalise arrangements with the key data providers. The Scoping Report proposed that a series of Memoranda of Understanding define the relationships between the inventory agency and key data providers, outlining the responsibilities that are conferred to the data providers under the NIS, including their involvement in the annual inventory review process. These Memoranda of Understanding have been agreed and they underpin the national system in Ireland in the years ahead. An updated national climate change strategy has been published providing a framework in which internal review of annual inventories will take place among all stakeholders to monitor progress on the strategy, thereby fulfilling another important requirement of national system implementation.

Formal QA/QC activities were lacking in Ireland's work on emission inventories up to 2005. The implementation of comprehensive QA/QC procedures in this reporting cycle according to the plan supporting the national inventory system represents an important step towards the general improvement in quality of Irish greenhouse gas inventories. The QA/QC elements include a plan and procedures for QA/QC in data selection and acquisition, data processing and reporting to comply with international requirements under Decision 280/2004/EC and the Kyoto Protocol. The plan provides guidance on and templates for appropriate quality checking, documentation and traceability, the selection of appropriate source data and calculation methodologies. It extends to peer review and expert review of inventory data and outlines the annual requirements of a continuous improvement programme for the inventory. Participation in the internal review mechanisms foreseen within the EU as part of the QA/QC plan developed for the EU inventory under Decision 280/2004/EC provides an opportunity to engage with other Member States in the examination and assessment of individual IPCC sectors and particular issues relating to methodologies and country-specific approaches that could bring mutual benefits to their greenhouse gas inventories. Ireland will continue to avail of this provision in the coming years.

Table 9.1 Recalculations by Gas 1990-2004

(a) Emissions by Gas 1990 –2004 in 2006 Submission (Gg CO₂eq)

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
Net CO ₂ emissions/removals	32,667.65	32,858.83	33,437.87	32,823.02	34,163.99	34,987.89	36,374.48	38,639.74	40,090.59	41,890.74	44,238.96	46,530.37	45,509.39	44,136.60	45,194.89
CO ₂ emissions (without LULUCF)	32,559.48	32,556.64	33,088.07	32,701.35	34,115.79	34,782.67	36,081.48	38,503.75	40,305.53	42,136.03	44,240.93	46,704.25	45,700.51	44,519.38	45,266.48
CH ₄	13,215.96	13,396.72	13,468.58	13,584.48	13,584.24	13,658.67	14,061.52	14,319.02	14,259.35	13,732.07	13,376.07	13,172.85	13,146.94	13,735.97	13,285.28
N ₂ O	9,801.99	9,632.70	9,654.21	9,779.97	10,086.57	10,279.24	10,361.68	10,320.58	10,905.14	11,034.23	10,521.43	10,054.19	9,565.64	9,399.85	9,243.07
HFCs	0.69	5.13	5.89	8.89	20.45	44.60	75.64	130.99	189.02	194.83	228.93	253.07	288.84	357.91	399.25
PFCs	0.09	0.09	0.09	0.09	75.38	75.38	103.09	130.82	61.87	195.93	305.41	295.98	212.40	228.79	196.37
SF ₆	35.40	36.38	37.36	38.33	81.85	82.83	102.06	132.10	94.24	68.96	55.91	69.43	70.22	118.59	70.01
Total (without CO₂ from LULUCF)	55,613.62	55,627.66	56,254.20	56,113.11	57,964.28	58,923.38	60,785.47	63,537.27	65,815.15	67,362.05	68,728.67	70,549.78	68,984.55	68,360.50	68,460.46

(b) Recalculated Emissions by Gas 1990 –2004 in 2007 Submission (Gg CO₂eq)

Net CO ₂ emissions/removals	32,673.69	33,672.17	33,607.34	33,586.49	34,666.43	35,662.63	37,447.07	38,851.78	40,551.58	42,137.21	44,992.27	47,361.95	45,776.82	44,847.48	45,552.30
CO ₂ emissions (without LULUCF)	32,552.88	33,399.53	33,286.13	33,430.00	34,683.79	35,480.90	37,139.01	38,622.65	40,688.06	42,288.97	44,883.98	47,342.95	45,902.78	45,146.17	45,746.89
CH ₄	13,286.51	13,474.80	13,555.44	13,667.18	13,667.56	13,722.62	14,031.64	14,102.74	14,331.89	13,919.70	13,438.10	13,260.61	13,268.48	13,881.23	13,338.46
N ₂ O	9,498.85	9,352.58	9,346.12	9,499.28	9,764.81	9,964.97	10,038.59	9,937.06	10,559.77	10,648.13	10,214.20	9,702.05	9,240.36	9,083.32	8,935.99
HFCs	0.69	5.14	5.90	8.90	20.52	44.89	76.23	132.09	189.01	194.82	228.95	251.50	276.55	349.52	384.04
PFCs	0.09	0.09	0.09	0.09	75.38	75.38	103.09	130.82	61.87	195.93	305.41	295.98	212.40	228.79	186.84
SF ₆	35.40	36.38	37.36	38.33	81.85	82.83	102.06	132.10	94.24	68.96	55.91	69.49	70.31	118.68	67.09
Total (without CO₂ from LULUCF)	55,374.43	56,268.52	56,231.04	56,643.80	58,293.92	59,371.58	61,490.61	63,057.46	65,924.84	67,316.51	69,126.54	70,922.59	68,970.89	68,807.72	68,659.31

(c) Percentage Change in Emissions by Gas 1990-2004

Net CO ₂ emissions/removals	0.02	2.48	0.51	2.33	1.47	1.93	2.95	0.55	1.15	0.59	1.70	1.79	0.59	1.61	0.79
CO ₂ emissions (without LULUCF)	-0.02	2.59	0.60	2.23	1.66	2.01	2.93	0.31	0.95	0.36	1.45	1.37	0.44	1.41	1.06
CH ₄	0.53	0.58	0.64	0.61	0.61	0.47	-0.21	-1.51	0.51	1.37	0.46	0.67	0.92	1.06	0.40
N ₂ O	-3.09	-2.91	-3.19	-2.87	-3.19	-3.06	-3.12	-3.72	-3.17	-3.50	-2.92	-3.50	-3.40	-3.37	-3.32
HFCs	0.00	0.19	0.17	0.11	0.34	0.65	0.78	0.84	-0.01	-0.01	0.01	-0.62	-4.25	-2.34	-3.81
PFCs	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	-4.85
SF ₆	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.09	0.13	0.08	-4.17
Total (without CO₂ from LULUCF)	-0.43	1.15	-0.04	0.95	0.57	0.76	1.16	-0.76	0.17	-0.07	0.58	0.53	-0.02	0.65	0.29

Table 9.2 Recalculations by IPCC Sector 1990-2004

(a) Emissions by IPCC Sector 1990 –2004 in 2006Submission (Gg CO₂eq)

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
Energy	31,665.36	31,757.08	32,387.91	32,008.14	33,269.36	34,002.81	35,287.99	37,493.27	39,512.58	41,500.00	43,097.80	45,402.54	44,507.99	44,346.10	44,400.91
Industrial Processes	3,166.43	2,866.26	2,781.45	2,746.11	3,117.66	3,062.75	3,204.72	3,656.26	3,494.73	3,560.03	4,186.65	4,294.30	3,734.39	3,050.92	3,169.72
Solvent and Other Product Use	80.94	82.82	83.04	83.50	84.40	86.19	86.87	87.28	88.33	85.01	80.32	79.55	77.20	75.71	74.47
Agriculture	19,240.15	19,425.74	19,460.76	19,688.71	19,856.60	20,083.07	20,455.48	20,582.13	21,207.28	20,787.46	19,712.98	19,296.04	19,058.84	19,137.49	18,981.72
LULUCF	108.17	302.20	349.80	121.67	48.20	205.22	293.01	135.99	-214.94	-245.29	-1.98	-173.88	-191.12	-382.78	-71.60
Waste	1,460.75	1,495.77	1,541.04	1,586.65	1,636.26	1,688.55	1,750.39	1,718.33	1,512.23	1,429.55	1,650.92	1,477.36	1,606.13	1,750.28	1,833.63
Total (without CO₂ from LULUCF)	55,721.79	55,929.86	56,604.00	56,234.78	58,012.48	59,128.60	61,078.47	63,673.26	65,600.21	67,116.76	68,726.69	70,375.90	68,793.43	67,977.72	68,388.86

(b) Recalculated Emissions by IPCC Sector 1990 –2004 in 2007 Submission (Gg CO₂eq)

Energy	31,603.37	32,588.14	32,538.75	32,739.47	33,802.76	34,676.83	36,324.52	37,520.10	39,808.17	41,528.02	43,683.48	45,944.03	44,634.23	44,913.64	44,814.39
Industrial Processes	3,166.43	2,866.26	2,781.45	2,746.12	3,117.73	3,063.04	3,205.31	3,657.35	3,494.73	3,560.02	4,186.67	4,292.79	3,722.18	3,041.57	3,147.14
Solvent and Other Product Use	80.86	82.74	82.96	83.42	84.32	86.11	86.79	87.20	88.06	84.75	80.07	79.66	77.37	75.96	75.70
Agriculture	19,063.03	19,235.61	19,286.82	19,488.13	19,652.85	19,857.05	20,247.02	20,362.83	21,024.43	20,594.83	19,535.86	19,128.74	18,889.41	18,983.90	18,830.02
LULUCF	120.81	272.64	321.21	156.48	-17.36	181.73	308.06	229.14	-136.47	-151.76	108.29	18.99	-125.96	-298.70	-194.58
Waste	1,460.75	1,495.77	1,541.04	1,586.65	1,636.26	1,688.55	1,626.97	1,429.97	1,509.45	1,548.90	1,640.45	1,477.36	1,647.69	1,792.65	1,792.06
Total (without CO₂ from LULUCF)	55,495.25	56,541.16	56,552.24	56,800.28	58,276.56	59,553.31	61,798.67	63,286.59	65,788.37	67,164.75	69,234.83	70,941.58	68,844.92	68,509.03	68,464.72

(c) Percentage Change in Emissions by Sector 1990-2004

Energy	-0.20	2.62	0.47	2.28	1.60	1.98	2.94	0.07	0.75	0.07	1.36	1.19	0.28	1.28	0.93
Industrial Processes	0.00	0.00	0.00	0.00	0.00	0.01	0.02	0.03	0.00	0.00	0.00	-0.04	-0.33	-0.31	-0.71
Solvent and Other Product Use	-0.10	-0.10	-0.10	-0.10	-0.09	-0.09	-0.09	-0.09	-0.31	-0.31	-0.31	0.14	0.22	0.33	1.65
Agriculture	-0.92	-0.98	-0.89	-1.02	-1.03	-1.13	-1.02	-1.07	-0.86	-0.93	-0.90	-0.87	-0.89	-0.80	-0.80
LULUCF	11.69	-9.78	-8.17	28.61	-136.02	-11.45	5.14	68.50	-36.51	-38.13	-5569.19	-110.92	-34.09	-21.97	171.76
Waste	0.00	0.00	0.00	0.00	0.00	0.00	-7.05	-16.78	-0.18	8.35	-0.63	0.00	2.59	2.42	-2.27
Total (without CO₂ from LULUCF)	-0.41	1.09	-0.09	1.01	0.46	0.72	1.18	-0.61	0.29	0.07	0.74	0.80	0.07	0.78	0.11

References

- Adams, M., Wagner, A., Goodwin, J., O'Leary, E., Creedon, M., McHugh, M and Gibson, C. 2005. Compiling Emission Inventories of HFCs, PFCs and SF₆ for inclusion in Ireland's greenhouse gas submissions to the EU under Decision 280/2004/EC under the United National Framework Convention on Climate Change. Netcen/ ED48411/R1, AEA Technology UK.
- Ahlvik, P. et al, 1997. COPERT II (Computer Programme to Calculate Emissions from Road Transport). Methodology and Emission Factors. European Environment Agency, Copenhagen.
- Boyle, O. C., 1987. National Database on Waste. An Foras Forbartha, Dublin.
- Carey, P. et al, 1996. National Waste Database Report 1998, Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- CEC (Council of the European Communities), 1991. Council Directive 91/676/EEC of 12 December 1991 concerning the protection of waters against pollution caused by nitrates from agricultural sources O.J. L167, July 1993.
- CEC (Council of the European Communities), 1993. Council Decision 93/389/EEC for a monitoring mechanism of Community CO₂ and other greenhouse gas emissions. O.J. L375 31 December 1991.
- CEC (Commission of the European Communities), 2004. Commission Decision 2004/156/EC of 29/01/2004 establishing guidelines for the monitoring and reporting of greenhouse gas emissions pursuant to Directive 2003/87/EC. O.J. L59, 26 February 2004.
- CEU (Council of the European Union), 1999. Council Directive 1999/31/EC on the landfill of waste. O.J. L182, July 1999.
- CEU (Council of the European Union), 2001. Council Directive 2001/80/EC of 23 October 2001 on the limitation of emissions of certain pollutants into the air from large combustion plants. O.J. L309, 27 November 2001.
- Collins, C. et al, 2004a. National Waste Database Interim Report 2002. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- Collins, C. et al, 2004b. National Waste Database Interim Report 2003. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- Collins, C. et al, 2005. National Waste Report 2004. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- Cossu, R., Andreottola, G. and Muntoni, A., 1996. Modelling Landfill Gas Production. In Landfilling of Waste Biogas. Edited by T. H. Christensen, R. Cossu and R. Stegman. E & FN Spon, London.
- Coulter, B.S. (Ed), 2004. Nutrient and trace element advice for grassland, tillage, vegetable and fruit crops. Teagasc, 19 Sandymount Avenue, Ballsbridge, Dublin 4.
- Crowe, M. et al, 2000. National Waste Database Report 1998, Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- Crowley, A. M. 2001. Prediction of net energy requirements of beef cattle. M.Agr.Sc. Thesis, National University of Ireland, University College Dublin, November 2001, 145 pp.
- Crowley, A. M., Keane, M.G., Agabriel, J. and O'Mara F. P. (2002). Prediction of net energy requirements of beef cattle. Proceedings of Agricultural Research Forum, 11 and 12 March 2002, Tullamore, p 19.
- Cruickshank, M.M., Tomlinson, R.W. and Trew, S. 2000. Application of CORINE land-cover mapping to estimate the carbon stored in the vegetation of Ireland. Journal of Environmental Management, 58, 269-287.

CTC (Clean Technology Centre), 2005. NMVOC Inventory for Ireland (SNAP Sector 6 Solvent and Other Product Use). Report to the Environmental Protection Agency.

DAF(Department of Agriculture and Food), 2006. CMMS Statistics Report 2005. Department of Agriculture and Food, Dublin. www.agriculture.gov.ie.

DELG (Department of the Environment and Local Government), 1997. Sustainable Development. A Strategy for Ireland. Department of the Environment and Local Government, Custom House, Dublin.

DELG (Department of the Environment and Local Government), 1998. A Policy Statement. Waste Management. Changing our Ways. Department of the Environment and Local Government, Custom House, Dublin.

DOE (Department of the Environment), 1992. The Environmental Protection Agency Act, 1992. SI No 7 of 1992. Department of the Environment, Custom House, Dublin.

DOE (Department of the Environment), 1994. Recycling for Ireland. A Strategy for Recycling Domestic and Commercial Waste. Department of the Environment, Custom House, Dublin.

DOE (Department of the Environment), 1997. IRELAND : Second National Communication under the UN Framework Convention on Climate Change. Department of the Environment, Custom House, Dublin.

EP and CEU (European Parliament and Council of the European Union), 2001. Directive 2001/81/EC of 23 October 2001 on national emissions ceilings for certain atmospheric pollutants. O.J. L309, 27 November 2001.

EP and CEU (European Parliament and Council of the European Union), 2003. Directive 2003/87/EC establishing a scheme for greenhouse gas emission allowance trading within the Community and amending Council Directive 96/61/EC. O.J. L275, 25 October 2003.

EP and CEU (European Parliament and Council of the European Union), 2004. Decision 280/2004/EC of the European Parliament and of the Council concerning a mechanism for monitoring Community greenhouse gas emissions and for implementing the Kyoto Protocol. O.J. L49, 19 February 2004.

ERL (Environmental Resources Limited), 1993. Towards a Recycling Strategy for Ireland. Department of the Environment, Custom House, Dublin.

Fehily Timoney and Company, 1985. Code of Good Practice for the Use of Biosolids in Agriculture. Guidelines for Farmers. Department of Environment and Local Government, Dublin.

Finn, J. et al, 2001. Estimation of Emissions of Non-Methane Volatile Organic Compounds from SNAP Sector 6 : Solvent and Other Product Use for Ireland in 1998. Prepared for the Environmental Protection Agency by the Clean Technology Centre, Cork Institute of Technology. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.

FSAI (Food Safety Authority of Ireland), 1999. Recommended Dietary Allowances for Ireland. Food Safety Authority of Ireland, Dublin. <http://www.fsai.ie/>

Gallagher, G., Hendrick, E. and Byrne, K. Preliminary Estimates of Biomass Carbon Stock Changes in Managed Forests in the Republic of Ireland over the period 1990-2000. In Irish Forestry Vol. 61 No 1, 2004.

Gardiner, M. J. and Radford, T., 1980. Soil Associations of Ireland and their Land Use Potential. Explanatory Bulletin to Soil Map of Ireland 1980, Soil Survey Bulletin No 36, An Foras Taluntais, Dublin.

Hamilton, G.J. 1975. Forest Mensuration Handbook. Forestry Commission Booklet 39, HMSO, London.

Hargreaves, K., Milne, R. and Cannell, M. 2004. Carbon balance of afforested peatland in Scotland. Forestry, 76, 299-317.

INRA, 1989. Ruminant Nutrition. Recommended Allowances and Feed Tables. R. Jarrige, Ed. John Libbey Eurotext, London and Paris.

- IPCC (Intergovernmental Panel on Climate Change), 1995. IPCC Guidelines for National Greenhouse Gas Inventories. OECD, Paris.
- IPCC (Intergovernmental Panel on Climate Change), 1997. Revised 1996 Guidelines for National Greenhouse Gas Inventories. OECD, Paris.
- IPCC (Intergovernmental Panel on Climate Change), 2000. IPCC Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. IGES (Institute for Global Environmental Strategies), Kanagawa, Japan.
- IPCC (Intergovernmental Panel on Climate Change), 2003. Good Practice Guidance for Land Use, Land-Use Change and Forestry. IGES (Institute for Global Environmental Strategies), Kanagawa, Japan.
- IUCC (Information Unit on Climate Change), 1998. The Kyoto Protocol to the Framework Convention on Climate Change. UNEP, Geneva.
- Le Bolloch, O. et al, 2006. National Waste Report 2005. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- Li, C., Frohling, S. and Harriss, R. 1994. Modelling carbon biogeochemistry in agricultural soils. *Global Biogeochemical Cycles*, 8, 237-254.
- Li, C., Narayanan, V. and Harriss, R.C. 1996. Model estimates of nitrous oxide emissions from agricultural lands in the United States. *Global Biogeochemical Cycles* 10, 297-306.
- Meaney, B. et al, 2003. National Waste Database Report 2001, Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- MCOS (M. C. O'Sullivan), 1994. Report on Solid Waste Recycling (Packaging). Department of the Environment, Custom House, Dublin.
- McCulloch, A., 2002. Comparison between the results of RTDI Projects L.S.-5.1.3(a) and L.S.-5.1.3(b): Emission Inventories of HFC, PFC and SF6 from Ireland. Note to the Environmental Protection Agency.
- McInnes, G. (editor), 1996. CORINAIR/EMEP Atmospheric Emission Inventory Guidebook. First Edition. European Environment Agency, Copenhagen.
- NEUT(Working Group on Nutrients and Eutrophication under the OSPAR Convention), 1999. Screening Procedure for Irish Coastal Waters with Regard to Eutrophication Status.
- O'Brien, P. 2007. Data Analysis and Estimation of Greenhouse Gas Emissions and Removals for the IPCC Sector Land Use Land-Use Change and Forestry in Ireland. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland (in preparation).
- O'Doherty, S. and McCulloch, A. 2002. Emission Inventories of HFC, PFC and SF6. RTDI Project L.S.-5.1.3(a) Final Report. Environmental Protection Agency, Johnstown Castle, Wexford.
- O'Leary, G. and Carty, G. Urban Waste Water Discharges in Ireland. A Report for the years 1996 and 1997.
- O'Leary, E., Finn, J. and Cunningham, D., 2002. Emission Inventories for HFCs, PFCs and SF6 for Ireland 1998. RTDI Project L.S.-5.1.3(b) Final Report. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- O'Leary, E., 2002. Emission Inventories for HFCs, PFCs and SF6 for Ireland 1995-2000. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.
- O'Mara, F. P., Caffrey, P. J. and Drennan, M. J. 1997. Net energy values of grass silage determined from comparative feeding trials. *Irish Journal of Agricultural and Food Research*, 36, 110.
- O'Mara, F. 2006. Development of Emission Factors for the Irish Cattle Herd. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.

Richardson, S. (editor), 1999. CORINAIR/EMEP Atmospheric Emission Inventory Guidebook. Second Edition. European Environment Agency, Copenhagen.

SBSTA (Subsidiary Body for Scientific and Technological Advice), 1999. Guidelines for the Preparation of National Communications by Parties included in Annex I to the Convention. Part I: UNFCCC Reporting Guidelines on Annual Inventories. FCCC/SBSTA/1999/6/Add.1. UNFCCC Secretariat, Bonn.

SBSTA (Subsidiary Body for Scientific and Technological Advice), 2002. Guidelines for the Preparation of National Communications by Parties included in Annex I to the Convention. Part I: UNFCCC Reporting Guidelines on Annual Inventories. FCCC/SBSTA/2002/L.5/Add.1, UNFCCC Secretariat, Bonn.

Smith, D., O'Neill, N., Doris, Y. and Moriarty, J., 2004. Urban Waste Water Discharges in Ireland. A Report for the Years 2002/2003. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.

Thistlethwaite, G., Baggott, S. L. and Goodwin, J., 2005. National Inventory System Scoping Report for the Greenhouse Gas Emission Inventory in Ireland. AEAT/ENV/R/1990 Issue 1, NETCEN, UK.

Tomlinson, R.W., Cruickshank, M.M., Cruickshank, J.G., Mallon, G., McStravick, A., Byrne, K., Collins, J., O'Toole, P., Farrell, E., Cunningham, D., Milne, R. and Walsh, M. 2005. Impact of land use and land-use change on carbon emission/fixation. Environmental Protection Agency, Johnstown Castle, Wexford, Ireland.

UNECE (United Nations Economic Commission for Europe), 1999. Protocol to the 1979 Convention on Long-Range Transboundary Air Pollution to Abate Acidification, Eutrophication and Ground Level Ozone. UNECE Secretariat, Geneva.

UNFCCC (United Nations Framework Convention on Climate Change). 2003a. FCCC/SBSTA/2003/L.22/Add.1. UNFCCC Secretariat, Bonn.

UNFCCC (United Nations Framework Convention on Climate Change). 2003b. Ireland : Report on the Individual Review of the Greenhouse Gas Inventory Submitted in 2003. FCCC/WEB/IRI(2)/2003/IRL (<http://unfccc.int/program/mis/ghg/indrev2003.html>). UNFCCC Secretariat, Bonn.

UNFCCC (United Nations Framework Convention on Climate Change). 2004 FCCC/SBSTA/2004/8 UNFCCC Secretariat, Bonn.

Yan, T., Agnew, R. E., Gordon, F. J. and Porter, M. G. (2000). Prediction of methane energy output in dairy and beef cattle offered grass silage-based diets. *Livestock Production Science*, 64: 253-263.

Glossary

Annex 1 Parties	Countries listed in Annex I to the United Nations Framework Convention on Climate Change
Base year	The year or period for which quantified emissions reduction or limits are established under the Kyoto Protocol.
BOD	Biochemical Oxygen Demand
CARBWARE	A national forest model to calculate carbon stock change and increment for forests
CFCs	Chlorofluorocarbon
CH₄	Methane
CHP	Combined Heat and Power.
CMMS	Cattle Movement and Monitoring System
CO	Carbon Monoxide
CO₂	Carbon Dioxide
CO₂ equivalent	The equivalent mass as CO ₂ of other greenhouse gases converted on the basis of their global warming potential (GWP)
COFORD	National Council for Forest Research and Development
Commitment Period	The years 2008 to 2012 inclusive for which quantified emissions reductions or limits, as well as other commitments, are established under the Kyoto Protocol
COP	Conference of the Parties
CORINAIR	Co-ordinated Information on the environment in the European /community-AIR. CORINAIR was one of several collaborative exercises initiated under the CORINE programme to harmonise the collection and dissemination of information on the environment in the EU.
CRF	Common Reporting Format
DAF	Department of Agriculture and Food
Decision 13/CP.9	Decision No 13 at Ninth Session of the COP in Milan, December 2003
Decision 18/CP.9	Decision No 18 at Ninth Session of the COP in Milan, December 2003
DEHLG	Department of Environment Heritage and Local Government
DNDC	DeNitrification-DeComposition, is a computer simulation model of carbon and nitrogen biogeochemistry in agri-ecosystems
EMEP	European Monitoring and Evaluation Programme, a co-operative programme for monitoring and evaluation of the long-range transmissions of air pollutants in Europe
Emission	(of a greenhouse gas). The release of greenhouse gases into the atmosphere.
Enteric Fermentation	The digestive process in ruminant animals (e.g cattle and sheep) where bacteria convert the feed to a usable form of energy for the animal, producing CH ₄ as a by product
EUROSTAT	Statistical Agency of the European Union
FAO	Food and Agriculture Organisation of the United Nations
FIPS	Forest Inventory and Planning System
Fluorinated Gases	HFCs, PFCs and SF ₆
Fossil Fuel	Peat, coal, oil and natural gas and associated derivatives
FTA	Fraction of BOD in sludge that degrades anaerobically
GasSim	A model that estimates landfill gas emissions from managed or unmanaged landfill sites (Golder Associates, UK)

GDP	Gross Domestic Product
Gg	Gigagram (10^9 g) = kilo tonne = 1,000 tonnes
Greenhouse Gas	A gas in the atmosphere that allows solar radiation through to the earth's surface, but traps some of the heat radiated back from the earth's surface
GWP	The cumulative warming over a specified time period, e.g. 100 years, resulting from a unit mass of a greenhouse gas emitted at the beginning of that time period, expressed relative to an absolute GWP of 1 for CO ₂
HCFCs	Hydrochlorofluorocarbon
HFCs	Hydrofluorocarbons
HGV	Heavy Goods Vehicle
IEA	International Energy Agency
IPC	Integrated Pollution Control
IPCC	Intergovernmental Panel on Climate Change
IUCC	Information Unit on Climate Change
kt	kilo tonne (1,000 tonnes)
Kyoto Protocol	The Protocol to the UNFCCC adopted by Decision 1/CP.3 under which industrialised countries agreed to reduce their combined greenhouse gas emissions in 1990 by at least 5 percent by the period 2008-2012
LandGEM	Landfill Gas Emissions model (USEPA)
Montreal Protocol	Protocol on substances that deplete the ozone layer
Mt	million tonnes or mega tonnes
N₂O	Nitrous Oxide
NIR	National Inventory Report
NMVOC	Non Methane Volatile Organic Compounds
NO_x	Nitrogen Oxides
NRA	National Roads Authority
OSPAR	Oslo and Paris Convention for the Protection of the Marine Environment
PFCs	Perfluorocarbons
SBSTA	Subsidiary Body for Scientific and Technological Advice
SEI	Sustainable Energy Ireland
SF₆	Sulphur Hexafluoride
Sink	The reservoir or pool in which sequestered carbon is stored; the process of sequestration
SO₂	Sulphur Dioxide
Teagasc	Irish Agriculture and Food Development Authority
TPER	Total Primary Energy Requirement
UNECE	United Nations Economic Commission for Europe
UNFCCC	United Nations Framework Convention on Climate Change
VOC	Volatile Organic Compounds

Annex A

Greenhouse Gases, GWP and IPCC Reporting Format

Table A.1 Greenhouse Gases and GWP Values

Greenhouse Gas	Chemical Formula	IPCC GWP (1995) ^a
Carbon Dioxide	CO ₂	1
Methane	CH ₄	21
Nitrous Oxide	N ₂ O	310
Hydrofluorocarbons (HFC)		
HFC-23	CHF ₃	11700
HFC-32	CH ₂ F ₂	650
HFC-41	CH ₃ F	150
HFC-43-10mee	C ₅ H ₂ F ₁₀	1300
HFC-125	C ₂ HF ₅	2800
HFC-134	C ₂ H ₂ F ₄ (CHF ₂ CHF ₂)	1000
HFC-134a	C ₂ H ₂ F ₄ (CH ₂ FCF ₃)	1300
HFC-152a	C ₂ H ₄ F ₂ (CH ₃ CHF ₂)	140
HFC-143	C ₂ H ₃ F ₃ (CHF ₂ CH ₂ F)	300
HFC-143a	C ₂ H ₃ F ₃ (CF ₃ CH ₃)	3800
HFC-227ea	C ₃ HF ₇	2900
HFC-236fa	C ₃ H ₂ F ₆	6300
HFC-245ca	C ₃ H ₃ F ₅	560
Perfluorocarbons(PFC)		
Perfluoromethane	CF ₄	6500
Perfluoroethane	C ₂ F ₆	9200
Perfluoropropane	C ₃ F ₈	7000
Perfluorobutane	C ₄ F ₁₀	7000
Perfluorocyclobutane	c-C ₄ F ₈	8700
Perfluoropentane	C ₅ F ₁₂	7500
Perfluorohexane	C ₆ F ₁₄	7400
Sulphur Hexafluoride	SF ₆	23900

(a) GWP (global warming potential) as provided by the IPCC in its Second Assessment Report

Table A.2 IPCC Reporting Format (Level 1 and Level 2)

IPCC SOURCE and SINK CATEGORIES	CO ₂	CH ₄	N ₂ O	HFC	PFC	SF ₆
1. Energy						
A. Fuel Combustion (Sectoral Approach)						
1. Energy Industries						
2. Manufacturing Industries and Construction						
3. Transport						
4. Other Sectors						
5. Other						
B. Fugitive Emissions from Fuels						
1. Solid Fuels						
2. Oil and Natural Gas						
2. Industrial Processes						
A. Mineral Products						
B. Chemical Industry						
C. Metal Production						
D. Other Production						
E. Production of Halocarbons and SF ₆						
F. Consumption of Halocarbons and SF ₆						
G. Other						
3. Solvent and Other Product Use						
A. Paint Application						
B. Degreasing and Dry Cleaning						
C. Chemical Products Manufacture & Processing						
D. Other						
4. Agriculture						
A. Enteric Fermentation						
B. Manure Management						
C. Rice Cultivation						
D. Agricultural Soils						
E. Prescribed Burning of Savannas						
F. Field Burning of Agricultural Residues						
G. Other						
5. Land-Use Change and Forestry						
A. Forestry						
B. Cropland						
C. Grassland						
D. Wetland						
E. Settlements						
F. Other Land						
G. Other						
6. Waste						
A. Solid Waste Disposal on Land						
B. Wastewater Handling						
C. Waste Incineration						
D. Other						
7. Other						
Memo Items:						
International Bunkers						
Multilateral Operations						
CO₂ Emissions from Biomass						

The grey cells indicate sources/sinks where no emissions/removals of the various gases are expected

Annex B

Expanded Energy Balance Sheets for 2005

Table B.1 Expanded Energy Balance Sheet 2005

2005	Units = ktoe	NACE	COAL	Bituminous Coal	Anthracite + Manuf Ovoids	Coke	Lignite	PEAT	Milled Peat	Sod Peat	BRIQUETTES	OIL	Crude	Refinery Gas	Gasoline	Kerosene	JET Kerosene
Indigenous Production			0	0				810	627	183		0					
Imports			1,884	1,822	37		25	0				10,690	3,346		1,251	378	1,178
Exports			4	0	4		0	10			10	1,361			181	5	0
Mar. Bunkers			0					0				106					
Stock Change			-47	-44	-3		-0	-24	-30	0	5	28			87	8	21
Primary Energy Supply (incl non-energy)			1,832	1,777	31	0	24	776	597	183	-5	9,250	3,385	0	1,158	381	1,199
Primary Energy Requirement (excl. non-energy)			1,832	1,777	31	0	24	776	597	183	-5	8,960	3,385	0	1,158	381	1,199
Transformation Input			1,416	1,416	0	0	0	610	610	0	0	4,164	3,384	8	0	0	0
Public Thermal Power Plants			1,410	1,410				504	504	0		766					
Combined Heat and Power Plants			6	6				7	7			8		8			
Pumped Storage Consumption																	
Briquetting Plants			0					99	99			0					
Oil Refineries & other energy sector			0					0				3,390	3,384				
Transformation Output			0	0	0	0	0	95	0	0	95	3,155	0	100	714	263	0
Public Thermal Power Plants			0					0				0					
Combined Heat and Power Plants - Electricity			0					0				0					
Combined Heat and Power Plants - Heat																	
Pumped Storage Generation																	
Briquetting Plants								95			95	0					
Oil Refineries								0				3,155		100	714	263	0
Exchanges and transfers			18	-11	29	0	0	0	0	0	0	-17	0	0	1	317	-316
Electricity																	
Heat																	
Other			18	-11	29							-17			1	317	-316
Own Use and Distribution Losses			0					15	15			137		92			
Available Final Energy Consumption			434	350	60	0	24	247	-27	183	91	8,087	1	0	1,872	961	883
Non-Energy Consumption			0	0	0	0	0	0	0	0	0	290	0	0	0	0	0
Final non-Energy Consumption (Feedstocks)			0					0				290					
Total Final Energy Consumption			435	351	60	0	24	274	0	183	91	8,144	0	0	1,821	917	857
Industry*			163	163	0			0	0	0	0	1,075	0	0	0	125	0
Non-Energy Mining	13-14		0	0				0				45				1	
Food, beverages and tobacco	15 - 16		16	16				0				124				16	
Textiles and textile products	17 - 18		8	8				0				10				1	
Wood and wood products	20		0	0				0				13				2	
Pulp, paper, publishing and printing	21 - 22		0	0				0				8				1	
Chemicals & man-made fibres	24		0	0				0				48				7	
Rubber and plastic products	25		0	0				0				6				0	
Other non-metallic mineral products	26		139	139				0				299				3	
Basic metals and fabricated metal products	27 - 28		0	0				0				426				90	
Machinery and equipment n.e.c.	29		0	0				0				8				1	
Electrical and optical equipment	30 - 33		0	0				0				67				2	
Transport equipment manufacture	34 - 35		0	0				0				3				0	
Other manufacturing	36 - 37, 11		0	0				0			0	18				2	
Transport			0	0	0	0	0	0	0	0	0	5,069	0	0	1,821	0	857
Road Freight			0					0				1,218					
Road Private Car			0					0				1,911			1,521		
Public Passenger Services			0					0				131			26		
Rail			0					0				52					
Domestic Aviation			0					0				43			0		43
International Aviation			0					0				814					814
Fuel Tourism			0					0				763			182		
Unspecified			0					0				137			91		
Residential			246	163	59		24	273		183	90	1,144			0	793	
Commercial/Public Services			26	24	1	0	1	0	0	0	0	587	0	0	0	0	0
Commercial Services			26	24	1		1	0				383				0	
Public Services			0					0		0	0	204					
Agricultural			0	0				0				270			0	0	
Statistical Difference			-2	-1	-0	0	0	-26	-27	1	0	-347	1	0	51	43	25

Table B.1 (continued) Expanded Energy Balance Sheet 2005

2005	Units = ktoe	NACE	Fueloil	LPG	Gasoil / Diesel /DERV	Petroleum Coke	Naphta	Bitumen	White Spirit	Lubricants	NATURAL GAS	RENEWABLES	HYDRO	Wind	Biomass	Landfill Gas	Biogas	Liquid Biofuel	Solar	Geothermal	ELECTRICITY	Heat	TOTAL
Indigenous Production											462	392	54	96	206	25	9	1	0	0			1 663
Imports			1,098	123	2 637	312	0	238	120	10	3,016	0									176		15,765
Exports			1,013	23	44	0	17	0	71	6	0	0									0		1,376
Mar. Bunkers			24		83						0	0											106
Stock Change			26	-3	-149	0	-1	0	0	0	0	-0						-0					-43
Primary Energy Supply (incl non-energy)			87	97	2,361	312	-18	238	48	4	3,477	391	54	96	206	25	9	1	0	0	176	0	15,903
Primary Energy Requirement (excl. non-energy)			87	97	2,361	312	-18	0	0	0	3,477	391	54	96	206	25	9	1	0	0	176	0	15,613
Transformation Input			697	0	75	0	0	0	0	0	2,044	30	0	0	2	25	2	0	0	0	60	0	8,323
Public Thermal Power Plants			697		68						1,958	25				25							4,663
Combined Heat and Power Plants			0	0	0						85	5			2		2						111
Pumped Storage Consumption																					46		46
Briquetting Plants											0												99
Oil Refineries & other energy sector					6						0										14		3,404
Transformation Output			893	63	1,117	0	5	0	0	0	0	11	0	0	1	9	1	0	0	0	2,073	0	5,335
Public Thermal Power Plants											9					9					1,991		2,000
Combined Heat and Power Plants - Electricity											2				1		1				53		55
Combined Heat and Power Plants - Heat											0												0
Pumped Storage Generation																					30		30
Briquetting Plants											0												95
Oil Refineries			893	63	1,117		5				0												3,155
Exchanges and transfers			13	0	-13	-18	0	0	0	0	0	-161	-54	-96	-1	-9	-1	0	0	0	161	0	1
Electricity												-161	-54	-96	-1	-9	-1				161		0
Heat																							0
Other			13		-13	-18					0												1
Own Use and Distribution Losses			31	8	6						72	0									280		503
Available Final Energy Consumption			265	152	3,383	293	-13	238	48	4	1,361	212	0	0	203	0	7	1	0	0	2,071	0	12,412
Non-Energy Consumption			0	0	0	0		238	48	4	0	0	0	0	0	0	0	0	0	0	0	0	290
Final non-Energy Consumption (Feedstocks)								238	48	4	0	0											290
Total Final Energy Consumption			480	146	3,643	278	1	0	0	0	1,340	211	0	0	203	0	7	1	0	0	2,094	0	12,499
Industry*			453	62	191	243	1	0	0	0	428	163	0	0	159	0	4	0	0	0	660	0	2,490
Non-Energy Mining	13-14		5	0	38	0					32	0									54		131
Food, beverages and tobacco	15 - 16		59	4	45	0					152	54			50		4				162		508
Textiles and textile products	17 - 18		3	1	5	0					1	0									8		27
Wood and wood products	20		6	0	5	0					3	109			109						28		154
Pulp, paper, publishing and printing	21 - 22		3	1	3	0					20	0									30		58
Chemicals & man-made fibres	24		24	1	16	0					107	0									97		252
Rubber and plastic products	25		1	0	5	0					8	0									30		45
Other non-metallic mineral products	26		10	3	40	243					48	0									52		538
Basic metals and fabricated metal products	27 - 28		327	2	7	0					11	0									41		478
Machinery and equipment n.e.c.	29		2	2	3	0					12	0									16		36
Electrical and optical equipment	30 - 33		6	47	13	0					24	0									108		200
Transport equipment manufacture	34 - 35		0	0	2	0					7	0									9		19
Other manufacturing	36 - 37, 1		7	1	7	0	1				2	0									25		45
Transport			17	5	2,369	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	5	0	5,075
Road Freight					1 218						0												1 218
Road Private Car				5	385						1							1					1 912
Public Passenger Services					105						0												131
Rail					52						0										5		57
Domestic Aviation											0												43
International Aviation											0												814
Fuel Tourism					581						0												763
Unspecified			17		29						0												137
Residential			0	70	246	35					522	44			44				0	0	646		2,874
Commercial/Public Services			10	10	567	0	0	0	0	0	390	3	0	0	0	0	3	0	0	0	728	0	1,735
Commercial Services			1	8	374						171	3			0		3				522		1,105
Public Services			9	3	192						219	0									206		630
Agricultural				0	270						0	0									55		325
Statistical Difference			-215	6	-260	16	-14	0	0	0	21	1	0	0	-0	0	0	0	0	-0	-24	0	-376

Annex C

Calculation Sheets for Energy 2005

Table C.1 Calculation Sheet for Emissions from Fuel Combustion 2005 (continued on following pages)

Sectoral Disaggregation of Fuel Combustion from National Energy Balance				Emission Factors			Emissions		
				CO ₂	CH ₄	N ₂ O	CO ₂	CH ₄	N ₂ O
		kTOE	TJ	kg/TJ	kg/TJ	kg/TJ	Gg	Mg	Mg
	1A1a Public Electricity								
1	Coal	1410.39	59050.32	93540	NO	14	5523.68	NO	827.00
2	Peat	503.80	21092.90	113120	NO	12	2386.00	NO	253.00
3	Fuel Oil and Gas Oil	766.00	32068.36	79920	NO	14	2562.77	NO	409.00
4	Natural Gas	1958.34	81991.68	56880	NO	3	4664.00	NO	246.00
5	Landfill Gas	24.94	1044.06	54940	NO	3	57.36	NO	3.00
	Public Electricity Total	4663.40	195247.32				15136.45	0	1738.00
	1A1b Refinery Fuel								
6	Refinery Gas	100.24	4196.88	55880	0	3	238.43	NO	12.59
7	Fuel Oil	30.53	1278.31	75801	0	10	95.11	NO	12.78
8	LPG	7.88	330.09	65083	2	3	59.88	0.66	0.99
9	Gasoil/Diesel/DERV	6.21	259.85	73403	2	10	17.80	0.52	2.60
	Refinery Total	144.86	6065.13				411.22	1.18	28.96
	1A1c Manufacture of Briquettes								
10	Peat	14.55	901.95	121540	50	5	109.62	45.10	4.51
	1A2a-1A2f Industry Fuel								
11	Bituminous Coals	169.53	7098.02	94600	100	3	671.47	709.80	21.29
12	Kerosene	124.56	5215.11	71400	5	10	372.36	26.08	52.15
13	Fuel Oil	453.05	18968.46	76000	0	10	1441.60	NO	189.68
14	LPG	61.95	2593.58	63700	2	3	165.21	5.19	7.78
15	Gasoil/Diesel/DERV	191.36	8012.03	73300	5	10	587.28	40.06	80.12
16	Pet Coke	242.85	10167.48	100800	50	12	1024.88	508.37	122.01
17	Naptha	1.05	44.00				NO	NO	NO
18	Natural Gas	500.75	20965.34	56800	2	3	1190.83	41.93	62.90
19	Biomass	159.43	6675.03	110000	30	4	734.25	200.25	26.70
20	Biogas	3.95	165.36	84200	0	0	13.92	0.00	0.00
	Industry Total	1908.48	79904.41				5453.64	1531.68	562.64
	1A3a Aviation								
21	Civil Aviation Kerosene		1509.57	71400	0.10	0.10	107.73	1.682	3.699
	1A3b Road Transport Fuel								
22	Gasoline	1821.15	76247.91	69960	25.0	11.4	5334.30	1908	873
23	Gasoil/Diesel/DERV	2315.985	96965.67	73300	2.4	5.7	7107.58	228	557
24	LPG	4.505	188.62	63700	5.3	0.0	12.02	1	NO
	Road Transport Total	4141.64	173402.20				12453.90	2137.00	1430.00

Table C.1 Calculation Sheet for Emissions from Fuel Combustion 2005 (continued from previous page)

Sectoral Disaggregation of Fuel Combustion from National Energy Balance				Emission Factors			Emissions		
				CO ₂	CH ₄	N ₂ O	CO ₂	CH ₄	N ₂ O
Sector/Fuel		kTOE	TJ	kg/TJ	kg/TJ	kg/TJ	Gg	Mg	Mg
1A3c-1A3e Other Transport Fuel									
25	Railway Diesel	51.597	2160.24	73300	5	30	158.35	10.80	64.81
26	Navigation Fuel Oil	16.743	701.01	76000	5	30	53.28	3.51	21.03
27	Navigation Gasoil	1.194	50	73300	5	30	3.67	0.25	1.50
28	Gas Distribution Use Gasoil	69.43	2907.05	56800	5	3	165.12	14.54	5.81
Other Transport Total		138.964	5818.304				380.41	29.09	93.15
1A4a Commercial/Institutional Fuel									
29	Bituminous Coal	24.49	1025.41	94600	100	12	97.00	102.54	12.30
30	Anthracite + Manufactured Ovoids	0.88	37.02	98260	100	12	3.64	3.70	0.44
31	Lignite	0.70	29.13	101200	50	12	2.95	1.46	0.35
32	Briquettes	0.46	19.12	98860	50	5	1.89	0.96	0.10
33	Fuel Oil	9.85	412.36	76000	5	10	31.34	2.06	4.12
34	LPG	10.14	424.40	63700		2	27.03	0.00	0.85
35	Gasoil / Diesel/ DERV	566.85	23732.93	73300	5	10	1739.62	118.66	237.33
36	Natural Gas	403.09	16876.49	56800	5	2	958.58	84.38	33.75
37	Biomass	0.17	7.18	110000	30	4	0.79	0.22	0.03
38	Biogas	2.97	124.38	84200	0	0	10.47	NO	NO
Commercial/Institutional Total		1019.60	42688.42				2862.06	313.98	289.28
1A4b Residential Fuel									
39	Bituminous Coal	163.24	6834.74	94600	100	12	646.57	683.47	82.02
40	Anthracite + Manufactured Ovoids	59.14	2476.15	98260	100	12	243.31	247.61	29.71
41	Lignite	23.53	985.32	101200	50	12	99.71	49.27	11.82
42	Sod Peat	182.79	7653.14	104000	50	5	795.93	382.66	38.27
43	Briquettes	90.30	3780.52	98860	50	5	373.74	189.03	18.90
44	Kerosene	792.76	33191.09	71400	5	10	2369.84	165.96	331.91
45	LPG	69.83	2923.67	63700		2	186.24	0.00	5.85
46	Gasoil / Diesel/ DERV	246.19	10307.37	73300	5	10	755.53	51.54	103.07
47	Petroleum Coke	35.01	1465.91	100800	50	12	147.76	73.30	17.59
48	Natural Gas	521.61	21838.69	56800	5	2	1240.44	109.19	43.68
49	Biomass	43.67	1828.47	110000	30	4	201.13	54.85	7.31
Residential Total		2228.08	93285.06				6859.07	2006.87	690.14
1A4c Agriculture Fuel									
50	Gasoil	269.99	11303.00	73300	5	10	828.51	56.515	113.03
Total Energy							44602.61	6123.10	4953.40

Table C.2 Emissions from Fuel Combustion Allocated by IPCC Source Category

	GREENHOUSE GAS SOURCE AND SINK CATEGORIES	AGGREGATE ACTIVITY DATA	IMPLIED EMISSION FACTORS			EMISSIONS		
			CO ₂	CH ₄	N ₂ O	CO ₂	CH ₄	N ₂ O
			(tTJ)	(kg/TJ)	(kg/TJ)	(Gg)		
A	I.A.1. Energy Industries	202214.40				15657.29	0.05	1.77
B	Solid Fuels	81045.17	98.95	0.56	13.38	8019.30	0.05	1.08
C	Liquid Fuels	38133.49	77.99	0.03	11.48	2973.98	0.00	0.44
D	Gaseous Fuels	81991.68	56.88	NO	3.00	4664.00	NO	0.25
E	Biomass	1044.06	54.94	NO	3.00	57.36	NO	0.00
F	I.A.2 Manufacturing Industries and Construction	79860.46				5453.71	1.54	0.56
G	Solid Fuels	7098.02	94.60	100.00	3.00	671.47	0.71	0.02
H	Liquid Fuels	44956.67	79.88	12.89	10.05	3591.34	0.58	0.45
I	Gaseous Fuels	20965.35	56.80	2.00	3.00	1190.90	0.04	0.06
J	Biomass	6840.43	110.00	30.00	4.00	752.45	0.21	0.03
K	I.A.3 Transport	180730.07				12942.05	2.17	1.53
L	Solid Fuels	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO	NE,NO
M	Liquid Fuels	177823.02	71.85	12.10	8.55	12776.92	2.15	1.52
N	Gaseous Fuels	2907.05	56.80	5.00	2.00	165.13	0.01	0.01
O	Biomass	NO	NO	NO	NO	NO	NO	NO
P	I.A.4 Other Sectors	147276.93				10549.80	2.38	1.09
Q	Solid Fuels	22840.55	99.15	72.71	8.49	2264.74	1.66	0.19
R	Liquid Fuels	83761.18	72.66	5.56	9.72	6085.92	0.47	0.81
S	Gaseous Fuels	38715.18	56.80	5.00	2.00	2199.15	0.19	0.08
T	Biomass	1960.02	108.36	28.10	3.75	212.39	0.06	0.01
U	I.A.5 Other (Not specified elsewhere) ⁽⁶⁾	NO				NO	NO	NO
V	I.A. Fuel Combustion	610081.86				44602.85	6.13	4.95
	Memo Items							
W	Air Bunkers	34387.48	71365.00	1.07	2.33	2454.07	36.00	80.00
X	Marine Bunkers	4454.32	73899.00			329.17	NE	NE
Y	CO2 from Biomass	9844.51	103.83			1022.20	NA	NA

Table C.3 Correspondence Between National Disaggregation of Sources and IPCC Combustion Source Categories

IPCC Source Category/Fuel Groups from Table C.2	National Disaggregated Sources from Table C.1
A 1.A.1 Energy Industries (A = B+C+D+E) B (a) Solid Fuels C (b) Liquid Fuels D (c) Gaseous Fuels E (d) Biomass F 1.A.2 Manufacturing Industries (F = G+H+I+J) G (a) Solid Fuels H (b) Liquid Fuels I (c) Gaseous Fuels J (d) Biomass K 1.A.3 Transport (K = L+M+N+O) L (a) Solid Fuels M (b) Liquid Fuels N (c) Gaseous Fuels O (d) Biomass P 1.A.4 Other Sectors (P = Q+R+S+T) Q (a) Solid Fuels R (b) Liquid Fuels S (c) Gaseous Fuels T (d) Biomass U 1.A.5 Other V 1.A Fuel Combustion (V = A+F+K+P+U)	1+2+10 3+7+8+9 4+6 5 11 12+13+14+15+16+17 18 19+20 NO 21+22+23+24+25+26+27 28 NO 29+30+31+32+39+40+41+42+43 33+34+35+44+45+46+47+50 36+48 37+38+49 NO

Table C.4 Emissions of CO₂ from the Reference Approach in 2005 [CRF 2005 Table 1.A(b); Biomass excluded]

FUEL TYPES			Unit	Production	Imports	Exports	International bunkers	Stock change	Apparent consumption	Conversion factor (TJ/Unit)	NCV/ GCV ⁽¹⁾	Apparent consumption (TJ)	Carbon emission factor (t C/TJ)	Carbon content (Gg C)	Carbon stored (Gg C)
Liquid Fossil	Primary Fuels	Crude Oil	kt	NA	2,894.00			-32.00	2,925.99	42.81	NCV	125,273.42	20.00	2,505.47	NA
		Orimulsion		NO	NO	NO		NO	NO	NO	NCV	NO	NO	NO	NO
		Natural Gas Liquids	ktoe	97.99	NA	NA		NA	97.99	41.87	NCV	4,102.65	17.73	72.73	NA
	Secondary Fuels	Gasoline	kt		1,190.00	122.00	1.01	-80.00	1,147.00	44.59	NCV	51,143.36	19.08	975.82	NA
		Jet Kerosene	kt		965.00		671.64	-68.00	361.35	44.10	NCV	15,935.54	19.47	310.31	NA
		Other Kerosene	kt		383.00	13.00	NA	2.00	368.00	44.20	NCV	16,264.08	19.47	316.69	NA
		Shale Oil			NO	NO		NO	NO	NO	NCV	NO	NO	NO	NO
		Gas / Diesel Oil	kt		2,510.00	124.00	101.00	-14.00	2,298.99	43.31	NCV	99,564.83	19.99	1,990.40	NA
		Residual Fuel Oil	kt		1,318.00	942.00	49.00	54.00	273.00	41.24	NCV	11,257.55	20.73	233.34	NE
		Liquefied Petroleum Gas (LPG)	kt		112.00	20.00			92.00	47.16	NCV	4,338.35	17.37	75.37	NA
		Ethane			NO	NO		NO	NO	NO	NCV	NO	NO	NO	NA
		Naphtha	ktoe		NE	27.33		-6.31	-21.02	41.87	NCV	-880.07	20.00	-17.60	NE
		Bitumen	ktoe		166.57	NO		NO	166.57	41.87	NCV	6,973.95	22.00	153.43	153.43
		Lubricants	ktoe		34.34	6.06			28.28	41.87	NCV	1,184.03	20.00	23.68	11.84
		Petroleum Coke	ktoe		293.18				293.18	41.87	NCV	12,274.86	27.49	337.45	NO
Refinery Feedstocks			NO	NO		NO	NO	NO	NCV	NO	NO	NO	NO		
Other Oil			NO	NO		NO	NO	NO	NCV	NO	NO	NO	NO		
Other Liquid Fossil											NO		NO	NO	
Other non-specified				NO	NO	NO	NO	NO	NO	NCV	NO	NO	NO	NO	
Liquid Fossil Totals											347,432.56		6,977.08	165.27	
Solid Fossil	Primary Fuels	Anthracite ⁽²⁾	ktoe	NO	41.90	5.35		-2.69	39.24	41.87	NCV	1,642.90	26.80	44.03	NO
		Coking Coal		NO	NO	NO		NO	NO	NO	NCV	NO	NO	NO	NA
		Other Bituminous Coal	ktoe	NE	1,663.44			47.21	1,616.23	41.87	NCV	67,668.32	25.80	1,745.84	NA
		Sub-bituminous Coal		NO	NO	NO	NO	NO	NO	NO	NCV	NO	NO	NO	NO
		Lignite	ktoe	NO	19.41			3.79	15.62	41.87	NCV	653.98	27.60	18.05	NO
		Oil Shale		NO	NO	NO		NO	NO	NO	NCV	NO	NO	NO	NO
	Peat		889.35				314.28	575.07	41.87	NCV	24,077.03	33.26	800.71	NA	
	Secondary Fuels	BKB ⁽³⁾ and Patent Fuel			NA	10.19		-8.86	-1.33	41.87	NCV	-55.68	26.96	-1.50	NA
Coke Oven/Gas Coke				NO	NO		NO	NO	NO	NCV	NO	NO	NO	NO	
Other Solid Fossil											NO		NO	NO	
Other non-specified				NO	NO	NO	NO	NO	NO	NCV	NO	NO	NO	NO	
Solid Fossil Totals											93,986.54		2,607.13	NA,NO	
Gaseous Fossil		Natural Gas (Dry)	ktoe	690.07	2,965.35			2.42	3,653.00	41.87	NCV	152,943.80	15.48	2,368.03	NE
Other Gaseous Fossil												NO		NO	NO
Other non-specified				NO	NO	NO	NO	NO	NO	NCV	NO	NO	NO	NO	NO
Gaseous Fossil Totals												152,943.80		2,368.03	NE,NO
Total												594,362.90		11,952.23	165.27
Biomass total												8,851.31		253.00	NO
		Solid Biomass	ktoe	183.97					183.97	41.87	NCV	7,702.46	30.00	231.07	NO
		Liquid Biomass		NO	NO	NO		NO	NO	NO	NCV	NO	NO	NO	NO
		Gas Biomass	ktoe	27.44					27.44	41.87	NCV	1,148.86	19.09	21.93	NO

Table C.5 Comparison of Results from Sectoral Approach and Reference Approach for 2005 [CRF 2005 Table 1.A(c)]

FUEL TYPES	REFERENCE APPROACH			SECTORAL APPROACH ⁽¹⁾		DIFFERENCE ⁽²⁾	
	Apparent energy consumption ⁽³⁾	Apparent energy consumption (excluding non-energy use and feedstocks) ⁽⁴⁾	CO ₂ emissions	Energy consumption	CO ₂ emissions	Energy consumption	CO ₂ emissions
	(PJ)	(PJ)	(Gg)	(PJ)	(Gg)	(%)	(%)
Liquid Fuels (excluding international bunkers)	347.43	NE	24,976.64	332.04	24,419.74	4.63	2.28
Solid Fuels (excluding international bunkers) ⁽⁵⁾	93.99	NE	9,559.46	94.90	9,562.92	-0.97	-0.04
Gaseous Fuels	152.94	NE	8,682.77	152.23	8,633.96	0.47	0.57
Other ⁽³⁾	NO	NE	NO	NA,NO	NA,NO		
<i>Total⁽⁵⁾</i>	<i>594.36</i>	<i>NE</i>	<i>43,218.88</i>	<i>588.02</i>	<i>42,616.62</i>	<i>2.62</i>	<i>1.41</i>

Annex D

Activity Data for Agriculture,

Methane Emission Factors

and

Input Data for the calculation of Nitrous Oxide emissions from Agricultural sSils

Year 1990-2005

Table D.1 (a) Activity Data for Agriculture

Housing and Storage (1000's)

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Total Cattle	5,969.10	6,100.40	6,147.30	6,236.40	6,263.90	6,343.80	6,450.70	6,661.00	6,881.60	6,951.80	6,557.70	6,330.10	6,408.10	6,332.90	6,223.50	6,211.50
Dairy Cows	1,341.60	1,322.20	1,288.00	1,246.20	1,248.30	1,233.00	1,220.80	1,215.60	1,201.40	1,198.80	1,173.80	1,152.80	1,148.00	1,128.70	1,135.70	1,121.80
All Other Cattle(exclud.Dairy Cows)	4,627.50	4,778.20	4,859.30	4,990.20	5,015.60	5,110.80	5,229.90	5,445.40	5,680.20	5,753.00	5,383.90	5,177.30	5,260.10	5,204.20	5,087.80	5,089.70
<i>Other Cows</i>	659.20	729.40	784.00	916.70	936.60	968.70	1,004.60	1,083.40	1,163.80	1,196.20	1,166.80	1,155.20	1,159.70	1,150.80	1,144.20	1,150.80
<i>Dairy Heifers</i>	172.30	185.30	182.10	198.70	193.70	209.40	235.40	243.80	244.00	223.80	210.40	202.90	206.20	215.80	225.60	238.00
<i>Other Heifers</i>	100.00	91.30	91.70	117.30	121.40	107.20	129.20	139.00	154.00	128.80	125.20	140.40	147.50	141.60	140.90	143.60
<i>Cattle <1 yrs</i>	1,436.20	1,477.00	1,491.00	1,472.30	1,564.90	1,556.50	1,631.40	1,735.00	1,828.60	1,789.60	1,648.90	1,689.90	1,879.40	1,805.70	1,751.10	1,746.00
<i>Cattle 1-2 yrs</i>	1,311.70	1,347.60	1,399.30	1,379.00	1,361.40	1,403.90	1,380.20	1,424.60	1,481.90	1,548.90	1,446.40	1,269.30	1,329.00	1,363.80	1,319.90	1,253.40
<i>Cattle >2yrs</i>	922.60	920.10	881.50	873.80	803.10	829.10	810.50	778.00	763.10	818.90	738.70	669.80	485.30	471.20	449.50	499.20
<i>Bulls</i>	25.50	27.50	29.70	32.40	34.50	36.00	38.60	41.60	44.80	46.80	47.50	49.80	53.00	55.30	56.60	58.70
Total Sheep	8,020.98	8,483.65	8,735.75	8,977.22	8,559.06	8,363.83	8,329.04	8,050.87	8,572.21	8,547.15	7,957.34	7,615.85	6,925.31	6,722.29	6,902.84	6,599.73
<i>Ewes Lowland</i>	2,396.60	2,542.54	2,621.99	2,576.45	2,511.11	2,426.99	2,369.07	2,389.75	3,056.41	2,936.12	2,814.25	2,704.31	2,637.25	2,552.34	2,463.79	2,590.08
<i>Ewes Upland</i>	1,960.85	2,080.26	2,145.26	2,108.00	2,054.54	1,985.72	1,938.33	1,955.25	1,309.89	1,258.34	1,206.11	1,158.99	1,130.25	1,093.86	1,055.91	647.52
<i>Rams</i>	116.85	122.55	126.45	125.05	122.05	120.00	113.15	115.50	115.70	113.25	110.65	106.55	104.65	102.35	100.00	96.40
<i>Other Sheep>1</i>	298.38	174.80	161.35	179.22	194.86	205.33	192.19	215.37	245.21	218.35	204.74	182.20	184.46	205.74	199.44	179.73
<i>Lambs</i>	3,248.30	3,563.50	3,680.70	3,988.50	3,676.50	3,625.80	3,716.30	3,375.00	3,845.00	4,021.10	3,621.60	3,463.80	2,868.70	2,768.00	3,083.70	3,086.00
Pigs	1,220.85	1,325.95	1,405.50	1,503.55	1,513.40	1,546.15	1,642.70	1,708.25	1,809.75	1,774.85	1,731.50	1,756.05	1,775.55	1,713.35	1,701.55	1,679.55
<i>Gilts in Pig</i>	20.80	22.30	25.60	22.75	21.45	24.00	24.50	26.85	25.60	24.85	23.15	24.05	20.05	20.00	21.30	19.05
<i>Gilts not yet Served</i>	11.85	14.00	14.75	14.60	15.05	17.40	16.85	17.70	18.70	16.20	16.80	17.70	19.55	17.80	18.90	19.15
<i>Sows in Pig</i>	83.35	91.05	96.30	100.55	98.70	99.90	103.15	107.95	109.10	108.60	110.40	108.45	110.00	103.95	102.40	100.10
<i>Other Sows for Breeding</i>	30.30	31.20	33.85	32.30	29.60	30.95	35.80	37.05	38.00	37.70	35.10	39.20	32.85	23.15	30.45	33.65
<i>Boars</i>	6.25	6.65	6.55	6.35	5.65	5.30	5.10	5.10	4.75	4.20	4.00	3.55	3.30	3.00	2.75	2.60
<i>Pigs 20 Kg +</i>	749.15	802.65	836.50	904.95	917.65	951.95	1,015.75	1,063.90	1,144.35	1,094.10	1,037.90	1,033.65	1,049.30	1,029.25	1,018.55	1,006.95
<i>Pigs Under 20 Kg</i>	319.15	358.10	391.95	422.10	425.30	416.65	441.55	449.75	469.25	489.25	504.15	529.55	540.50	516.25	507.20	498.10
Poultry	11,446.33	12,371.71	12,946.57	12,746.41	13,708.55	14,112.70	15,049.62	15,221.04	15,349.21	15,142.98	15,338.00	15,673.15	15,193.32	15,796.47	16,588.77	16,057.10
<i>Layer</i>	1,868.25	1,800.00	2,231.00	1,831.50	1,730.00	1,370.50	1,701.00	1,580.00	1,558.50	1,537.00	1,572.00	1,676.00	1,613.00	1,906.60	1,906.27	1,950.00
<i>Broiler</i>	8,035.13	8,904.90	9,066.82	9,522.47	10,392.54	11,092.18	11,729.88	12,096.34	12,286.79	12,200.11	12,426.10	12,628.89	12,321.96	12,672.21	13,212.60	12,817.70
<i>Turkey</i>	1,542.95	1,666.80	1,648.76	1,392.44	1,586.01	1,650.02	1,618.74	1,544.70	1,503.92	1,405.87	1,339.91	1,368.26	1,258.35	1,217.66	1,469.90	1,326.98
Horses	53.50	63.10	65.10	66.20	67.00	68.00	69.90	71.90	72.80	75.50	69.90	71.00	72.60	70.40	72.80	72.80
Mules	7.30	7.30	8.00	8.50	7.80	7.00	7.60	7.10	7.50	7.30	5.00	4.90	4.70	5.80	5.70	5.70
Goats	17.40	17.40	17.80	17.60	16.10	15.60	14.90	15.20	15.10	13.50	8.10	7.80	7.70	7.60	7.50	7.50
Fertiliser (1000's kg/N)	379,311	370,121	358,302	377,985	404,811	428,826	416,918	380,350	431,999	442,916	407,598	368,667	363,513	388,080	362,525	352,165

Table D.1 (b) Activity Data for Agriculture

Pasture (1000's)

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
total Cattle	6,816.10	6,912.10	6,951.40	6,981.70	6,996.70	7,034.00	7,313.50	7,532.80	7,639.90	7,387.00	7,037.40	7,049.70	6,992.20	6,967.00	7,015.70	6,887.90
Dairy Cows	1,359.70	1,330.80	1,277.90	1,263.50	1,260.60	1,256.20	1,266.40	1,251.70	1,233.80	1,200.60	1,177.50	1,182.50	1,164.10	1,155.60	1,156.10	1,117.20
All Other Cattle(exclud.Dairy Cows)	5,456.40	5,581.30	5,673.50	5,718.20	5,736.10	5,777.80	6,047.10	6,281.10	6,406.10	6,186.40	5,859.90	5,867.20	5,828.10	5,811.40	5,859.60	5,770.70
Other Cows	731.30	817.30	889.10	979.70	1,011.00	1,039.10	1,112.70	1,201.90	1,247.90	1,217.30	1,187.00	1,196.80	1,154.20	1,187.30	1,207.10	1,231.00
Dairy Heifers	158.60	129.70	174.50	187.90	203.90	224.10	231.40	243.90	228.80	213.60	206.50	198.30	230.70	215.80	229.60	227.50
Other Heifers	68.60	50.10	94.50	111.50	101.50	117.20	128.80	143.50	126.70	116.50	125.10	132.80	143.20	137.00	139.60	140.90
Cattle <1 yrs	1,716.10	1,764.70	1,694.50	1,737.50	1,736.20	1,746.10	1,852.10	1,938.20	1,965.00	1,820.70	1,751.90	1,824.40	1,799.30	1,728.50	1,771.40	1,687.30
Cattle 1-2 yrs	1,663.10	1,692.00	1,637.60	1,587.00	1,586.10	1,586.10	1,639.40	1,717.00	1,782.60	1,706.00	1,517.10	1,515.00	1,593.20	1,577.20	1,534.80	1,509.50
Cattle >2yrs	1,092.60	1,098.80	1,151.80	1,077.90	1,057.70	1,022.90	1,036.20	985.70	1,002.10	1,057.70	1,016.20	941.10	844.70	901.50	910.60	905.60
Bulls	26.10	28.70	31.50	36.70	39.70	42.30	46.50	50.90	53.00	54.60	56.10	58.80	62.80	64.10	66.50	68.90
Total Sheep	8,020.98	8,483.65	8,735.75	8,977.22	8,559.06	8,363.83	8,329.04	8,050.87	8,572.21	8,547.15	7,957.34	7,615.85	6,925.31	6,722.29	6,902.84	6,599.73
Lowland Ewes	2,396.60	2,542.54	2,621.99	2,576.45	2,511.11	2,426.99	2,369.07	2,389.75	3,056.41	2,936.12	2,814.25	2,704.31	2,637.25	2,552.34	2,463.79	2,590.08
Upland Ewes	1,960.85	2,080.26	2,145.26	2,108.00	2,054.54	1,985.72	1,938.33	1,955.25	1,309.89	1,258.34	1,206.11	1,158.99	1,130.25	1,093.86	1,055.91	647.52
Rams	116.85	122.55	126.45	125.05	122.05	120.00	113.15	115.50	115.70	113.25	110.65	106.55	104.65	102.35	100.00	96.40
Other Sheep>1	298.38	174.80	161.35	179.22	194.86	205.33	192.19	215.37	245.21	218.35	204.74	182.20	184.46	205.74	199.44	179.73
Lambs	3,248.30	3,563.50	3,680.70	3,988.50	3,676.50	3,625.80	3,716.30	3,375.00	3,845.00	4,021.10	3,621.60	3,463.80	2,868.70	2,768.00	3,083.70	3,086.00
Pigs	1,220.85	1,325.95	1,405.50	1,503.60	1,513.40	1,546.15	1,642.70	1,708.30	1,809.75	1,774.90	1,731.50	1,756.15	1,775.55	1,713.40	1,701.55	1,679.60
Gilts in Pig	20.80	22.30	25.60	22.75	21.45	24.00	24.50	26.85	25.60	24.85	23.15	24.05	20.05	20.00	21.30	19.05
Gilts not yet Served	11.85	14.00	14.75	14.60	15.05	17.40	16.85	17.70	18.70	16.20	16.80	17.70	19.55	17.80	18.90	19.15
Sows in Pig	83.35	91.05	96.30	100.55	98.70	99.90	103.15	107.95	109.10	108.60	110.40	108.45	110.00	103.95	102.40	100.10
Other Sows for Breeding	30.30	31.20	33.85	32.30	29.60	30.95	35.80	37.05	38.00	37.70	35.10	39.20	32.85	23.15	30.45	33.65
Boars	6.25	6.65	6.55	6.35	5.65	5.30	5.10	5.10	4.75	4.20	4.00	3.55	3.30	3.00	2.75	2.60
Pigs 20 Kg +	749.15	802.65	836.50	904.95	917.65	951.95	1,015.75	1,063.90	1,144.35	1,094.10	1,037.90	1,033.65	1,049.30	1,029.25	1,018.55	1,006.95
Pigs Under 20 Kg	319.15	358.10	391.95	422.10	425.30	416.65	441.55	449.75	469.25	489.25	504.15	529.55	540.50	516.25	507.20	498.10
Poultry	11,446.33	12,371.71	12,946.57	12,746.41	13,708.55	14,112.70	15,049.62	15,221.04	15,349.21	15,142.98	15,338.00	15,673.15	15,193.32	15,796.47	16,588.77	16,094.68
Layer	1,868.25	1,800.00	2,231.00	1,831.50	1,730.00	1,370.50	1,701.00	1,580.00	1,558.50	1,537.00	1,572.00	1,676.00	1,613.00	1,906.60	1,906.27	1,950.00
Broiler	8,035.13	8,904.90	9,066.82	9,522.47	10,392.54	11,092.18	11,729.88	12,096.34	12,286.79	12,200.11	12,426.10	12,628.89	12,321.96	12,672.21	13,212.60	12,817.70
Turkey	1,542.95	1,666.80	1,648.76	1,392.44	1,586.01	1,650.02	1,618.74	1,544.70	1,503.92	1,405.87	1,339.91	1,368.26	1,258.35	1,217.66	1,469.90	1,326.98
Horses	53.50	63.10	65.10	66.20	67.00	68.00	69.90	71.90	72.80	75.50	69.90	71.00	72.60	70.40	72.80	72.80
Mules	7.30	7.30	8.00	8.50	7.80	7.00	7.60	7.10	7.50	7.30	5.00	4.90	4.70	5.80	5.70	5.70
Goats	17.40	17.40	17.80	17.60	16.10	15.60	14.90	15.20	15.10	13.50	8.10	7.80	7.70	7.60	7.50	7.50
Fertiliser(kgs N)	379,311	370,121	358,302	377,985	404,811	428,826	416,918	380,350	431,999	442,916	407,598	368,667	363,513	388,080	362,525	352,165

Table D.2 (a) Emission factors used in the calculation of CH₄ emissions from enteric fermentation (kg/head/year)

Animal Category	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Dairy cows	101.38	101.95	102.52	103.09	103.66	104.23	104.80	105.38	105.95	106.52	107.09	107.66	108.23	108.80	108.50	107.22
Beef cows(Suckler Cows)	74.03	74.04	74.06	74.07	74.08	74.10	74.11	74.12	74.13	74.15	74.16	74.17	74.19	74.20	74.41	74.12
Dairy heifers	51.82	51.69	51.56	51.44	51.31	51.18	51.05	50.93	50.80	50.67	50.54	50.42	50.29	50.16	50.03	49.90
Beef heifers	55.42	55.28	55.14	55.00	54.85	54.71	54.57	54.43	54.29	54.15	54.00	53.86	53.72	53.58	53.44	53.30
Cattle <1 year	28.76	28.75	28.75	28.74	28.74	28.73	28.73	28.72	28.72	28.71	28.71	28.70	28.70	28.70	28.80	28.79
Cattle 1-2years	57.88	57.47	57.05	56.64	56.22	55.81	55.39	54.98	54.56	54.15	53.73	53.32	52.90	52.49	51.88	52.30
Cattle >2 year	38.37	37.60	36.83	36.06	35.29	34.52	33.75	32.98	32.21	31.44	30.67	29.90	29.13	28.37	28.85	30.07
Bulls for breeding	86.38	86.01	85.61	85.23	84.86	84.49	84.12	83.75	83.38	83.01	82.63	82.26	81.89	81.55	81.21	80.87
Male cattle																
< 1 year	30.46	30.39	30.32	30.25	30.17	30.10	30.03	29.96	29.89	29.82	29.74	29.67	29.60	29.53	29.70	29.72
1 - 2 years	62.22	62.08	61.94	61.79	61.65	61.51	61.37	61.22	61.08	60.94	60.80	60.65	60.51	60.37	59.26	58.96
> 2 years*	55.08	53.48	51.88	50.28	48.68	47.08	45.48	43.87	42.27	40.67	39.07	37.47	35.87	34.27	35.23	37.67
Female cattle																
< 1 year	27.05	27.11	27.17	27.24	27.30	27.36	27.42	27.49	27.55	27.61	27.67	27.74	27.80	27.86	27.89	27.86
1 - 2 years	53.54	52.85	52.16	51.48	50.79	50.10	49.41	48.73	48.04	47.35	46.66	45.98	45.29	44.60	44.50	45.64
> 2 years*	21.65	21.71	21.77	21.84	21.90	21.96	22.02	22.09	22.15	22.21	22.27	22.34	22.40	22.46	22.46	22.46
Sheep																
Lowland Sheep	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00
Upland Sheep	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00
Rams	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00
Sheep > yrs	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00
Lambs	3.38	3.38	3.38	3.38	3.38	3.38	3.38	3.38	3.38	3.38	3.38	3.38	3.38	3.38	3.38	3.38
Horses	18.00	18.00	18.00	18.00	18.00	18.00	18.00	18.00	18.00	18.00	18.00	18.00	18.00	18.00	18.00	18.00
Mules	10.00	10.00	10.00	10.00	10.00	10.00	10.00	10.00	10.00	10.00	10.00	10.00	10.00	10.00	10.00	10.00
Goats	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00
Pigs	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50
Gilts in Pig	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50
Gilts not yet Served	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98	0.98
Sows in Pig	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50
Other Sows for Breeding	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50
Boars	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50
Pigs > 20 Kg	0.44	0.44	0.44	0.44	0.44	0.44	0.44	0.44	0.44	0.44	0.44	0.44	0.44	0.44	0.44	0.44
Pigs < 20 Kg	0.11	0.11	0.11	0.11	0.11	0.11	0.11	0.11	0.11	0.11	0.11	0.11	0.11	0.11	0.11	0.11
Poultry	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE

* Note: This value is low because this category of animal only live part of their third year.

Table D.2 (b) Emission factors used in the calculation of CH₄ emissions from manure management (kg/head/yr)

Animal Category	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Dairy cows	21.57	21.49	21.41	21.33	21.25	21.17	21.09	21.01	20.93	20.85	20.77	20.69	20.61	20.53	20.53	20.43
Beef cows(Suckler Cows)	14.02	14.01	14.00	13.99	13.98	13.97	13.96	13.96	13.95	13.94	13.93	13.92	13.91	13.90	13.90	13.86
Dairy heifers	13.40	13.21	13.02	12.83	12.64	12.45	12.26	12.07	11.88	11.69	11.50	11.31	11.12	10.93	10.74	10.55
Beef heifers	15.61	15.40	15.19	14.98	14.77	14.56	14.35	14.13	13.92	13.71	13.50	13.29	13.08	12.87	12.66	12.45
Cattle <1 year	9.26	9.19	9.13	9.06	8.99	8.93	8.86	8.79	8.72	8.66	8.59	8.52	8.46	8.39	8.44	8.45
Cattle 1-2years	15.71	15.41	15.11	14.81	14.51	14.20	13.90	13.60	13.30	13.00	12.70	12.40	12.10	11.80	11.43	11.73
Cattle >2 year	2.45	2.33	2.21	2.09	1.98	1.86	1.74	1.62	1.50	1.38	1.27	1.15	1.03	0.91	0.97	1.13
Bulls for breeding	23.79	23.42	23.05	22.67	22.30	21.93	21.56	21.18	20.81	20.44	20.07	19.69	19.32	18.95	18.58	18.21
Male cattle																
< 1 year	9.73	9.64	9.54	9.45	9.35	9.26	9.16	9.07	8.97	8.88	8.78	8.69	8.59	8.50	8.58	8.60
1 - 2 years	16.68	16.49	16.31	16.12	15.93	15.75	15.56	15.37	15.18	15.00	14.81	14.62	14.44	14.25	13.75	13.82
> 2 years*	4.57	4.33	4.09	3.86	3.62	3.38	3.14	2.91	2.67	2.43	2.19	1.96	1.72	1.48	1.60	1.91
Female cattle																
< 1 year	8.79	8.75	8.71	8.67	8.63	8.59	8.55	8.52	8.48	8.44	8.40	8.36	8.32	8.28	8.30	8.29
1 - 2 years	14.74	14.32	13.91	13.49	13.08	12.66	12.25	11.83	11.42	11.00	10.59	10.17	9.76	9.34	9.11	9.63
> 2 years*	0.33	0.33	0.33	0.33	0.33	0.33	0.33	0.34	0.34	0.34	0.34	0.34	0.34	0.34	0.34	0.34
Sheep																
Lowland Sheep	0.21	0.21	0.21	0.21	0.21	0.21	0.21	0.21	0.21	0.21	0.21	0.21	0.21	0.21	0.21	0.21
Upland Sheep	0.22	0.22	0.22	0.22	0.22	0.22	0.22	0.22	0.22	0.22	0.22	0.22	0.22	0.22	0.22	0.22
Rams	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20
Sheep >1 yrs	0.21	0.21	0.21	0.21	0.21	0.21	0.21	0.21	0.21	0.21	0.21	0.21	0.21	0.21	0.21	0.21
Lambs	0.11	0.11	0.11	0.11	0.11	0.11	0.11	0.11	0.11	0.11	0.11	0.11	0.11	0.11	0.11	0.11
Horses	1.40	1.40	1.40	1.40	1.40	1.40	1.40	1.40	1.40	1.40	1.40	1.40	1.40	1.40	1.40	1.40
Mules	0.76	0.76	0.76	0.76	0.76	0.76	0.76	0.76	0.76	0.76	0.76	0.76	0.76	0.76	0.76	0.76
Goats	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12
Pigs																
Gilts in Pig	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46
Gilts not yet Served	17.17	17.17	17.17	17.17	17.17	17.17	17.17	17.17	17.17	17.17	17.17	17.17	17.17	17.17	17.17	17.17
Sows in Pig	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46
Other Sows for Breeding	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46
Boars	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46	21.46
Pigs > 20 Kg	12.88	12.88	12.88	12.88	12.88	12.88	12.88	12.88	12.88	12.88	12.88	12.88	12.88	12.88	12.88	12.88
Pigs < 20 Kg	8.58	8.58	8.58	8.58	8.58	8.58	8.58	8.58	8.58	8.58	8.58	8.58	8.58	8.58	8.58	8.58
Poultry																
Layers	2.28	2.28	2.28	2.28	2.28	2.28	2.28	2.28	2.28	2.28	2.28	2.28	2.28	2.28	2.28	2.28
Broilers	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08
Turkeys	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.27	0.08	0.08

* Note: This value is low because this category of animal only live part of their third year.

Table D.3 Input Data for the calculation of N₂O emissions from Agricultural Soils

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Frac _{GASF}	0.014	0.014	0.016	0.016	0.015	0.014	0.019	0.021	0.018	0.018	0.021	0.020	0.017	0.014	0.016	0.016
Frac _{GRAZ}	0.658	0.657	0.659	0.656	0.656	0.653	0.656	0.656	0.655	0.648	0.648	0.653	0.652	0.654	0.656	0.654
Frac _{GASH1}	0.489	0.487	0.490	0.492	0.492	0.492	0.492	0.493	0.493	0.494	0.493	0.492	0.492	0.493	0.492	0.493
Frac _{GASH2}	0.043	0.042	0.041	0.041	0.041	0.040	0.040	0.040	0.040	0.040	0.040	0.038	0.038	0.038	0.038	0.037
Frac _{GASH}	0.195	0.195	0.194	0.196	0.196	0.197	0.196	0.196	0.196	0.200	0.200	0.195	0.196	0.195	0.194	0.195
Frac _{LEACH}	0.100	0.100	0.100	0.100	0.100	0.100	0.100	0.100	0.100	0.100	0.100	0.100	0.100	0.100	0.100	0.100
F _{SN} (tonnes/year)	374,027	364,792	352,647	372,051	398,673	422,648	408,806	372,451	424,071	434,967	398,911	361,348	357,490	382,602	356,796	346,395
F _{AM} (tonnes/year)	75,524	77,516	77,731	78,292	78,195	79,019	80,075	81,841	83,946	84,286	80,678	78,107	77,513	76,667	76,238	75,649
F _S (tonnes/year)	105.954	109.229	112.817	116.150	119.252	120.912	133.556	200.853	369.625	387.009	394.296	413.930	572.911	602.662	666.410	701.578
F _{BN}	0.010	0.009	0.009	0.017	0.017	0.013	0.014	0.013	0.022	0.012	0.005	0.006	0.005	0.009	0.010	0.012
F _{CR} (tonnes/year)	19,692	19,001	19,335	16,201	17,934	19,439	20,848	19,969	19,919	21,128	21,683	19,742	17,757	19,962	23,568	18,034
F _{AW} (tonnes/year)	75,630	77,625	77,844	78,409	78,314	79,140	80,209	82,041	84,316	84,673	81,072	78,521	78,086	77,270	76,904	76,351

Annex E

Activity Data for the Carbon Stock Change Estimates of LULUCF Category 5.A Forest Land

Years 1990-2005

Table E.1 Activity Data and Carbon Stock Change Estimates for LULUCF Category 5.A Forest Land

A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	Q	R	S	T
Year	Affor	Refor	Felling	Cleared and Unclassified	Forest Cover	Total Forest	Harvest Volume	Harvest Carbon Stock	Affor Carbon Stock	Carbon Stock in young forests	Carbon Stock in Mature Forests	Carbon Stock in Total Forests	Total Carbon Stock Change	5.A.1 Forest Land Remaining Forest Land	5.A.2 Land Converted to Forest Land	Cropland Converted to Forest Land	Grassland Converted to Forest Land	Wetland Converted to Forest Land	Other Land Converted to Forest Land
	ha	ha	ha	ha	ha	ha	1000m ³	Gg C	Gg C	Gg C	Gg C	Gg C	Gg C	Gg C	Gg C	Gg C	Gg C	Gg C	Gg C
1990	15817	3882	4203	111528	370160	481688	1676	508.83	273.03	1937.06	13948.18	15885.24	215.831	271.155	-55.323	-5.532	-16.597	-31.534	-1.660
1991	19147	4203	4063	127246	373728	500974	1769	537.07	330.51	1853.03	14259.29	16112.32	227.089	311.119	-84.029	-8.403	-25.209	-47.897	-2.521
1992	16699	4063	4621	140883	376237	517120	2083	632.42	288.25	1801.56	14496.45	16298.01	185.684	237.158	-51.474	-5.147	-15.442	-29.340	-1.544
1993	15998	4621	4816	151985	380940	534918	2100	637.60	276.15	1776.17	14730.81	16506.97	208.965	234.357	-25.392	-2.539	-7.618	-14.473	-0.762
1994	19459	4816	5447	165792	386598	552390	2287	694.30	335.89	1818.06	14888.74	16706.80	199.826	157.932	41.894	4.189	12.568	23.880	1.257
1995	23710	5447	6203	180777	395331	576108	2382	723.28	409.27	1920.35	15005.91	16926.26	219.460	117.168	102.292	10.229	30.688	58.307	3.069
1996	20981	6203	7090	190089	407000	597089	2465	748.35	362.17	2044.93	15100.13	17145.05	218.794	94.217	124.577	12.458	37.373	71.009	3.737
1997	11434	7090	7185	189009	419514	608523	2322	705.11	197.37	2231.13	15164.68	17395.81	250.753	64.552	186.201	18.620	55.860	106.134	5.586
1998	12928	7185	7924	186511	434940	621451	2638	800.90	223.16	2376.63	15301.01	17677.64	281.829	136.331	145.498	14.550	43.649	82.934	4.365
1999	12668	7924	7747	186164	447955	634119	2777	843.12	218.67	2517.10	15455.78	17972.88	295.248	154.774	140.475	14.047	42.142	80.071	4.214
2000	15695	9238	8677	189917	459897	649814	3008	913.36	270.92	2746.15	15491.77	18237.92	265.034	35.990	229.044	22.904	68.713	130.555	6.871
2001	15465	8755	9132	190239	475501	665740	2836	861.01	266.95	3045.39	15504.83	18550.22	312.300	13.054	299.247	29.925	89.774	170.571	8.977
2002	15054	9258	10286	186422	494926	681348	2911	883.69	259.86	3295.12	15597.80	18892.92	342.699	92.976	249.723	24.972	74.917	142.342	7.492
2003	9097	10313	9289	177624	513338	690962	3000	910.80	157.03	3388.48	15928.79	19317.27	424.353	330.985	93.368	9.337	28.010	53.220	2.801
2004	9739	9289	9828	178667	522436	701103	2846	864.19	168.11	3513.85	16125.37	19639.23	321.955	196.584	125.371	12.537	37.611	71.461	3.761
2005	10096	9928	9781	178431	533150	711581	2942	893.19	174.27	3642.04	16353.13	19995.17	355.945	227.760	128.185	12.818	38.455	73.065	3.846

K The value 3,642.04 Gg is the afforestation carbon stock for the years 1986 to 1999 inclusive, similarly 3,513.85 is the afforestation carbon stock for the years 1985 to 1998 inclusive, etc

N Carbon stock change after harvest (corresponding in 2005 to difference between carbon stocks of 19,995.17 Gg in 2005 and 19,639.23 Gg in 2004)

P Carbon stock change for young forests (corresponding in 2005 to difference between carbon stocks of 3,642.04 Gg in 2005 and 3,513.85 Gg in 2004)

Q, R, S, T The total for 5.A.2 (column P) is split as Cropland – 0.10; Grassland – 0.30; Wetland – 0.57; Settlements – 0.00; Other Land – 0.0

DETERMINATION OF TIME-SERIES FOREST AREAS USING 1995 BASE YEAR DATA

The assumptions use to assign areas to the three different categories were:

1. Afforested and reforested areas 7 years and over, defined as cleared/unclassified in FIPS move each year into the young crops category. Areas were derived from Coillte felling and Forest Service planting records.
2. Five percent of the young crop category moves each year into the mature category. This means that there is a full turn-over of these crops every 20 years.
3. Mature crops are clearfelled and these areas come back to the cleared/unclassified category.
4. For the purposes of the model clearfell is defined as Coillte felling plus an arbitrary 200 ha of private felling.
5. The reforestation is derived from the clearfell area of the previous year.
6. The process works forward or back from FIPS base year 1995.

YOUNG CROPS

General rule for years before 1995:

Current year = (Current year+1) ha. - (afforestation [current year + 1 - minimum age for young trees] + reforestation [current year + 1 - minimum age for young trees])*(Category % related to planting) + (Current year+ 1)*Accretion Rate

Example: 1993 ha. = 1994 ha. - (afforestation 1987 + reforestation 1987)*species % + 1994 ha.*0.05

Example: 1994 ha. = 1995 ha. - (afforestation 1988 + reforestation 1988)*species % + 1995 ha.*0.05

1995 ha. = FIPS ha. For 1995 for a given category

General rule for years after 1995:

Current year = (Current year -1) ha. + (afforestation [current year - minimum age for young trees] + reforestation [current year - minimum age for young trees])*(Category % related to planting) - (Current year - 1)*Accretion Rate

Example: 1996 ha. = 1995 ha. + (afforestation 1989 + reforestation 1989)*species % - 1995 ha.*0.05

Example: 1997 ha. = 1996 ha. + (afforestation 1990 + reforestation 1990)*species % - 1996 ha.*0.05

MATURE CROPS

General Rule for years before 1995:

Current Year = (Current Year + 1)ha - ([Current Year + 1] Young Trees)ha*(Accretion Rate)+ ([Current Year + 1 Felling]ha * [Category % in Felling])

Example: 1993 ha. = 1994 ha. - 1994 'young' ha * 0.5 + 1994 Felling ha * Category % in Felling

Example: 1994 ha. = 1995 ha. - 1995 'young' ha. * 0.5 + 1995 Felling ha * Category % in Felling

1995 ha. = FIPS ha. For 1995 for a given category

General Rule for years after 1995:

Current Year = (Current Year - 1)ha + ([Current Year - 1] Young Trees)ha*(Accretion Rate) - ([Current Year Felling]ha * [Category % in Felling])

Example: 1996 ha. = 1995 ha. + 1995 'young' ha. * 0.5 - 1996 Felling ha * Category % in Felling

Example: 1997 ha. = 1996 ha. + 1996 'young' ha. * 0.5 - 1997 Felling ha * Category % in Felling

CLEARED/UNCLASSIFIED AREAS

The category cleared/unclassified represents total identified forest area by Forest Service less covered forest as located by remote sensing and classified in FIPS. This would include felled areas in which forest cover had not been established, recent plantings not yet classified and other productive unforested sites. This category is assumed not to store carbon.

General Rule for years before 1995:

Current Year= (Current Year + 1 ha) - Afforestation[Current Year +1] - Felling[Current Year + 1] + ([Current Year + 1 - minimum age for young trees]Afforestation) + ([Current Year + 1 - minimum age for young trees]Reforestation)

Example:

1994 ha. = 1995 ha. - 1995 Afforestation -1995 Felling + 1988 Afforestation + 1988 Reforestation

General Rule for years after 1995:

Current Year= (Current Year - 1 ha) + Afforestation[Current Year]+ Felling[Current Year] - ([Current Year - minimum age for young trees]Afforestation) - ([Current Year - minimum age for young trees]Reforestation)

Example:

1996 ha. = 1995 ha. + 1996 Afforestation +1996 Felling - 1989 Afforestation - 1989 Reforestation

The minimum age for young trees is 7 in all examples:

Accretion rate represents the movement of young categories into mature categories on the basis that a given percentage per annum reaches a given age. For example here (minimum age of 7 years assumed for young plantations and 25 years for mature plantations) the percentage is calculated as $[1/(25-7)]$ or 0.056%.

Annex F

Activity Data for 6.A Solid Waste Disposal on Land

Years 1968 – 2005

Table F.1 Time Series of Solid Waste Disposal and Composition 1968-2005

Year	Pop	MSW Prod Rate kg/cap/day	MSW Production tonnes	MSW to SWDS %	MSW to SWDS tonnes	Street Cleansing tonnes	MSW Organic %	MSW Paper %	MSW Textiles %	MSW Other %	MSW Organic tonnes	MSW Paper tonnes	MSW Textiles tonnes	MSW Other tonnes	DOC in Sludge tonnes	DOC in MSW tonnes
A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	Q
1968	2,912,200	1.00	1,062,953	0.75	797,215	21,259	0.36	0.28	0.03	0.19	286,997	223,220	23,916	151,471	13,564	183,503
1969	2,925,200	1.00	1,067,698	0.75	800,774	21,354	0.36	0.28	0.03	0.19	288,278	224,217	24,023	152,147	13,624	184,322
1970	2,949,900	1.00	1,076,714	0.75	807,535	21,534	0.36	0.28	0.03	0.19	290,713	226,110	24,226	153,432	13,739	185,879
1971	2,978,200	1.00	1,087,043	0.75	815,282	21,741	0.36	0.28	0.03	0.19	293,502	228,279	24,458	154,904	13,871	187,662
1972	3,024,400	1.00	1,103,906	0.75	827,930	22,078	0.36	0.28	0.03	0.19	298,055	231,820	24,838	157,307	14,086	190,573
1973	3,073,000	1.00	1,121,645	0.75	841,234	22,433	0.36	0.28	0.03	0.19	302,844	235,545	25,237	159,834	14,312	193,635
1974	3,123,900	1.00	1,140,224	0.75	855,168	22,804	0.36	0.28	0.03	0.19	307,860	239,447	25,655	162,482	14,549	196,843
1975	3,177,200	1.00	1,159,678	0.75	869,759	23,194	0.36	0.28	0.03	0.19	313,113	243,532	26,093	165,254	14,798	200,201
1976	3,227,800	1.00	1,178,147	0.75	883,610	23,563	0.36	0.28	0.03	0.19	318,100	247,411	26,508	167,886	15,033	203,390
1977	3,271,900	1.00	1,194,244	0.75	895,683	23,885	0.36	0.28	0.03	0.19	322,446	250,791	26,870	170,180	15,239	206,168
1978	3,314,000	1.00	1,209,610	0.75	907,208	24,192	0.36	0.28	0.03	0.19	326,595	254,018	27,216	172,369	15,435	208,821
1979	3,368,200	1.00	1,229,393	0.75	922,045	24,588	0.36	0.28	0.03	0.19	331,936	258,173	27,661	175,189	15,687	212,237
1980	3,401,000	1.00	1,241,365	0.75	931,024	24,827	0.36	0.28	0.03	0.19	335,169	260,687	27,931	176,895	15,840	214,303
1981	3,443,400	1.00	1,256,841	0.75	942,631	25,137	0.36	0.28	0.03	0.19	339,347	263,937	28,279	179,100	16,038	216,975
1982	3,480,000	1.00	1,270,200	0.75	952,650	25,404	0.36	0.28	0.03	0.19	342,954	266,742	28,580	181,004	16,208	219,281
1983	3,504,000	1.00	1,278,960	0.75	959,220	25,579	0.36	0.28	0.03	0.19	345,319	268,582	28,777	182,252	16,320	220,794
1984	3,529,000	1.00	1,283,556	0.75	962,667	25,671	0.36	0.28	0.03	0.19	346,560	269,547	28,880	182,907	16,436	221,645
1985	3,540,000	1.02	1,317,942	0.75	988,457	32,949	0.36	0.28	0.03	0.19	355,844	276,768	29,654	187,807	16,487	228,841
1986	3,540,600	1.04	1,344,012	0.75	1,008,009	33,600	0.35	0.28	0.03	0.19	352,803	282,242	30,240	191,522	16,490	231,532
1987	3,546,500	1.06	1,372,141	0.75	1,029,106	34,304	0.35	0.28	0.03	0.19	360,187	288,150	30,873	195,530	16,518	236,060
1988	3,530,700	1.09	1,404,689	0.75	1,053,517	35,117	0.34	0.28	0.03	0.19	358,196	294,985	31,606	200,168	16,444	239,614
1989	3,509,500	1.12	1,434,684	0.75	1,076,013	35,867	0.34	0.29	0.03	0.19	365,844	312,044	32,280	204,442	16,345	248,585
1990	3,505,800	1.15	1,471,560	0.77	1,133,101	36,789	0.34	0.29	0.03	0.19	385,254	328,599	33,993	215,289	16,328	260,644
1991	3,525,700	1.19	1,531,388	0.77	1,179,169	38,285	0.33	0.29	0.03	0.19	389,126	341,959	35,375	224,042	16,421	268,901
1992	3,554,500	1.24	1,608,767	0.78	1,254,838	40,219	0.33	0.30	0.03	0.18	414,097	376,451	31,371	225,871	16,555	285,734
1993	3,574,100	1.29	1,680,000	0.78	1,310,400	42,000	0.32	0.30	0.03	0.18	419,328	393,120	32,760	235,872	16,646	295,778
1994	3,585,900	1.33	1,740,775	0.80	1,392,620	43,519	0.32	0.30	0.02	0.18	445,638	417,786	27,852	250,672	16,701	310,283
1995	3,601,300	1.37	1,801,441	0.77	1,385,439	46,791	0.32	0.31	0.02	0.18	442,271	425,373	27,724	244,122	16,727	312,623
1996	3,626,100	1.41	1,866,172	0.80	1,492,938	55,985	0.32	0.31	0.02	0.18	476,588	458,379	29,875	263,064	16,750	336,995
1997	3,664,300	1.44	1,925,956	0.83	1,588,914	48,149	0.30	0.31	0.02	0.18	476,674	492,563	31,778	286,004	20,873	357,048
1998	3,703,100	1.46	1,975,653	0.85	1,685,766	80,999	0.27	0.32	0.02	0.19	455,204	547,850	36,142	323,463	23,745	394,392
1999	3,741,600	1.62	2,212,408	0.82	1,814,175	66,372	0.27	0.34	0.02	0.19	489,878	616,819	38,895	348,102	26,766	431,342
2000	3,789,500	1.77	2,448,206	0.79	1,934,083	73,446	0.28	0.36	0.03	0.19	541,543	696,270	48,352	371,110	30,083	483,192
2001	3,847,200	1.87	2,625,566	0.76	1,992,050	78,469	0.28	0.32	0.03	0.17	555,926	638,109	56,014	346,619	29,513	462,161
2002	3,917,200	1.86	2,654,812	0.72	1,901,864	65,573	0.28	0.31	0.03	0.18	541,125	583,430	55,998	344,080	21,147	426,092
2003	3,978,900	1.96	2,846,107	0.64	1,832,625	71,779	0.28	0.31	0.03	0.18	519,590	566,361	53,567	329,548	21,480	414,767
2004	4,043,800	2.01	2,964,905	0.61	1,818,536	69,661	0.37	0.25	0.08	0.11	667,081	446,310	146,987	206,469	22,188	407,954
2005	4,130,700	1.85	2,782,005	0.65	1,796,463	58,677	0.37	0.25	0.08	0.11	657,794	443,664	144,176	192,116	34,300	411,592

$$Q = G*0.25 + L*0.15 + M*0.4 + N*0.4 + O*0.15 + P$$

D = Total MSW – street cleanings

Table F.2 Potential CH₄ Production from Solid Waste 1968-2005

<i>Year</i>	<i>DOC in MSW tonnes</i>	<i>DOC Managed SWDS %</i>	<i>DOC Unmanaged SWDS %</i>	<i>Fraction DOC Dissimilated</i>	<i>Fraction CH₄ in Landfill</i>	<i>MCF Managed SWDS</i>	<i>MCF Unmanaged SWDS</i>	<i>Pot CH₄ Managed SWDS tonnes</i>	<i>Pot CH₄ Unmanaged SWDS tonnes</i>	<i>Pot CH₄ Total SWDS tonnes</i>
A	B	C	D	E	F	G	H	I	J	K
1968	183503	0.40	0.60	0.60	0.50	1.00	0.40	29360	17616	46977
1969	184322	0.40	0.60	0.60	0.50	1.00	0.40	29492	17695	47187
1970	185879	0.40	0.60	0.60	0.50	1.00	0.40	29741	17844	47585
1971	187662	0.41	0.59	0.60	0.50	1.00	0.40	30777	17715	48492
1972	190573	0.42	0.58	0.60	0.50	1.00	0.40	32016	17685	49701
1973	193635	0.43	0.57	0.60	0.50	1.00	0.40	33305	17660	50965
1974	196843	0.44	0.56	0.60	0.50	1.00	0.40	34644	17637	52281
1975	200201	0.45	0.55	0.60	0.50	1.00	0.40	36036	17618	53654
1976	203390	0.46	0.54	0.60	0.50	1.00	0.40	37424	17573	54997
1977	206168	0.47	0.53	0.60	0.50	1.00	0.40	38760	17483	56243
1978	208821	0.48	0.52	0.60	0.50	1.00	0.40	40094	17374	57468
1979	212237	0.49	0.51	0.60	0.50	1.00	0.40	41598	17318	58917
1980	214303	0.50	0.50	0.60	0.50	1.00	0.40	42861	17144	60005
1981	216975	0.51	0.49	0.60	0.50	1.00	0.40	44263	17011	61274
1982	219281	0.52	0.48	0.60	0.50	1.00	0.40	45610	16841	62451
1983	220794	0.53	0.47	0.60	0.50	1.00	0.40	46808	16604	63412
1984	221645	0.54	0.46	0.60	0.50	1.00	0.40	47875	16313	64188
1985	228841	0.55	0.45	0.60	0.50	1.00	0.40	50345	16477	66822
1986	231532	0.56	0.44	0.60	0.50	1.00	0.40	51863	16300	68163
1987	236060	0.57	0.43	0.60	0.50	1.00	0.40	53822	16241	70063
1988	239614	0.58	0.42	0.60	0.50	1.00	0.40	55590	16102	71693
1989	248585	0.59	0.41	0.60	0.50	1.00	0.40	58666	16307	74973
1990	260644	0.60	0.40	0.60	0.50	1.00	0.40	62555	16681	79236
1991	268901	0.60	0.40	0.60	0.50	1.00	0.40	64536	17210	81746
1992	285734	0.60	0.40	0.60	0.50	1.00	0.40	68576	18287	86863
1993	295778	0.60	0.40	0.60	0.50	1.00	0.40	70987	18930	89917
1994	310283	0.60	0.40	0.60	0.50	1.00	0.40	74468	19858	94326
1995	312623	0.60	0.40	0.60	0.50	1.00	0.40	75030	20008	95037
1996	336995	0.60	0.40	0.60	0.50	1.00	0.40	80879	21568	102447
1997	357048	0.60	0.40	0.60	0.50	1.00	0.40	85692	22851	108543
1998	394392	0.61	0.39	0.60	0.50	1.00	0.40	96232	24610	120842
1999	431342	0.62	0.38	0.60	0.50	1.00	0.40	106973	26226	133198
2000	483192	0.63	0.37	0.60	0.50	1.00	0.40	121764	28605	150369
2001	462161	0.69	0.31	0.60	0.50	1.00	0.40	128296	22627	150923
2002	426092	0.76	0.24	0.60	0.50	1.00	0.40	129191	16498	145689
2003	414767	0.82	0.18	0.60	0.50	1.00	0.40	136375	11813	148188
2004	407954	0.89	0.11	0.60	0.50	1.00	0.40	144579	7441	152020
2005	411592	0.95	0.05	0.60	0.50	1.00	0.40	156405	3293	159698

E from GPG

$$I = B * C * E * F * G * 16 / 12$$

G and H from IPCC Guidelines

$$J = B * D * E * F * H * 16 / 12$$

$$K = I + J$$